

Library of Congress Classification: R5-920, RA421-790.95

DETERMINING SANITATION RISK INDEX FOR MAKASSAR AND SINJAI ISLANDS USING SECOND CONFIRMATORY FACTOR ANALYSIS (2CFA)

Andy Susilawaty¹, Bambang Widjanarko Otok², Ambo Tuwo³, Anwar Daud⁴, Armyn Nurdin⁵

¹Department of Public Health, School of Health Science,
State Islamic University of Alauddin, Makassar,

²Laboratory of Environmental and Health Statistic,
Sepuluh Nopember Institute of Technology (ITS), Surabaya,

³Department of Fisheries, Faculty of Fisheries and Marine Science, Hasanuddin University, Makassar,

⁴Department of Environmental Health, School of Public Health, Hasanuddin University, Makassar,

⁵Faculty of Medicine, Hasanuddin University, Makassar (INDONESIA)

E-mails: andisusilawaty.uin@gmail.com, bambang_wo@statistika.its.ac.id

DOI: 10.7813/2075-4124.2015/7-1/A.22

Received: 05 Nov, 2014

Accepted: 17 Jan, 2015

ABSTRACT

The Indonesian archipelago (archipelago) has small islands. Small islands have a wide range of vulnerabilities due to its small size, both physically and socio-economic. In addition, the utilization of natural resources that are prone to bring greater conflict between economic development and environmental consequences. Priority environment is very important for small islands and coastal areas alone, which includes the balance of population and resources there are very important in the future will come. Given the standpoint of environmental health risks that have the potential for significant risk to human health and sanitation among other relevant indicator berish and drinking water sources, and ekskreta wastewater, solid waste, unhealthy behaviors, it is necessary to study on the risk index of sanitation in the islands Napier and Sinjai through approach Confirmatory Factor Analysis (CFA). The results showed that the risk of sanitation is fit model based on criteria CMIN / DF and RMSEA. The highest risk index sanitation is on some islands, namely Samolona, Barrang Lompo, Bird Loe, Kodongareng, Kambuno and Lumu-Lumu. So there needs to be the socialization of latrine ownership, ownership SPAL, use of clean and drinking water sources are not protected, the scarcity of clean water and drinking, physical quality of water does not qualify and the distance to the water source pollutant source ≤ 10 meters.

Key words: Islands, sanitation, clean water and drinking, and ekskreta wastewater, solid waste, behavior is unhealthy, CFA, Index

1. INTRODUCTION

Coastal areas and small islands have a wide range of distinctive characteristics which include characteristic ecological, geographic, demographic and socio-economic. Some ecological and environmental characteristics such as size relatively small, finite and vulnerable resource, risk of natural disasters (hurricanes, tsunamis, sea level rise, etc.), ecological instability, marine biodiversity richness and also the vulnerability of the species due to the influence of changes in activity land [1].

The present publication that has the characteristics of small islands such geographical location relatively isolated and surrounded by the sea which gives a high ratio between the sea than land space, this region also has a meeting of land and sea territory vast thereby increasing the fragility of coastal ecosystems [2]. The demand for management of coastal areas and the absence of interior hinterland areas or terrestrial central core area which basically is far from the sea that makes the planning and management of coastal resources is equated with national planning and management, the dominance of the sea and its use for delivery make these areas highly vulnerable against hazards associated with international shipping and waste disposal, as well as the limited area of land than sea make the islands are particularly vulnerable to global environmental phenomena such as sea level rise [3].

Several socio-economic characteristics of small islands, as shown by Tiempo (1993) [2], covering economic openness extreme example of external relations of the flow of trade, aid and investment technology, they are also highly dependent on foreign trade of the countries more large and has less influence on trade rules that run, which makes them very open economy.

In general, they also have a low economic resilience in recovering from the economic crisis and generally there is a close relationship between economic development and environmental assets as well as the limitations and the difficulty of adjusting the local population with job skills required.

Methods Confirmatory Factor Analysis (CFA) with maximum likelihood approach is used to identify indicators of environmental health risks that can measure four dimensions [4]. This approach requires a multivariate normal distribution of data [5, 6]. Observation units used are 16 islands in Makassar and Sinjai involving 15 indicators. From the results obtained scores 2CFA fit factor and then to calculate the risk index Sanitation.

2. LITERATURE REVIEW

2.1. Sanitation

The first thing to look at is the definition of the various islands and islets to have a common view in the subject matter covered. There are many definitions of small islands that have been established by various organizations and institutions both locally and globally [3]. Defines the island as a naturally formed area of land, surrounded by water, which is above water at high tide raises many interpretations because it involves the term "ground" to be declared clearly what it means. In the definition it can be said that the solid rock, which looks over the water at high tide should be considered as an island [7]. On the other hand, mangrove forests in the sea may look like an island, but there is no ground at high tide or low tide [3].

Society of small islands is faced with various problems caused by the location and their environment. Sanitation is a dominant issue. The discussion of sanitation in this paper refers to the safe disposal of human waste (excreta) [8]. Sanitation problem is more complex in the crowded slums are usually located in low-lying areas such as rivers, coastal areas and swamps than people in sparsely populated regions in rural areas [9, 14].

Poor sanitation, especially in densely populated areas, means exposure to an unpleasant environment and risk of the spread of infectious diseases through: contact with water, into the food chain and a breeding ground for insects. Lack of sanitation can also threaten the ecological balance of the environment when other species come into contact with contaminated water [10]. Where shallow groundwater, bacteriological quality and chemical quality rapidly deteriorated as a result of poor excreta disposal in areas of concentrated populations such as villages. Community Tonga - Lifuka Ha'apai is an example of contamination of the water resources of this land [8, 11].

In small islands ecologically sensitive, sanitation and safe waste disposal is closely related to water supply problems. As populations increase, so does the problem of water supply and sewage disposal, if the supply of fresh water is limited, especially on the island of coral (coral), should not be completely contaminated [9]. Such pollution is the cause of the cholera outbreak in urban Kiribati, a small island in the South Pacific, in 1977, and triggered the construction of toilets that use directly discharged into the open sea [12].

Lack of sanitation facilities to dispose of human waste, produces a high probability that the residents of coastal communities vulnerable to fecal-oral infection that is transmitted through consumption of contaminated food and drink [13]. The microorganisms that cause these infections are found in the feces of infected people or animals from defecating in the open by livestock or people who do not have a toilet [15]. This contaminated surface water that can infect humans through contamination on their hands, their equipment, or their drinking water supply. Children are particularly exposed to infection when playing or bathing in water [12].

2.2. Confirmatory Factor Analysis (CFA)

CFA is a method to test how well the measured variables can represent constructs or factors preconceived [5]. CFA can be divided into two, namely First-Order and Second-Order CFA. In First-Order a latent variable is measured by several indicators that can be measured directly, this is the model equation [4, 6].

$$x = \Lambda_x \xi + \delta \tag{1}$$

With the covariance matrix of x , written as a function θ and represented as $\Sigma(\theta)$

$$\Sigma(\theta) = \Lambda_x \Phi \Lambda_x' + \Theta_\delta \tag{2}$$

Where x is the observation variable, Λ is the factor loading matrix, ξ is a latent variable, and δ is the matrix of measurement errors Φ is the covariance matrix of the latent variable ξ and Θ_δ is the covariance matrix of measurement error δ .




Fig. 1. First-Order CFA




Fig. 2. Second-Order CFA

2.3. Index

Determination of the Index is based on the CFA is as follows:

$$I_R = X' F_j \times 100 \tag{3}$$

with,
 X = data indicator and F = factor score of latent variable

3. RESEARCH METHODOLOGY

The data used is data on the environmental health risks in Makasar dan Sinjai islands. The sampling method used simple random sampling [16]. The variables used in this research include five latent variables and 13 indicators. Source water and drinking (X1) consists of 6 indicators of Coverage use unprotected water sources (%) (X1.1), Coverage use unprotected sources of drinking water (%) (X1.2), scarcity of clean water (%) (X1.3), scarcity of drinking water (%) (X1.4), the physical quality of water does not qualify (%) (X1.5) and the distance to the water source pollutant source ≤ 10 meters (%) (X1.6), wastewater and ekskreta (X2) consists of two indicators, namely not have latrines (%) (X2.1), Not having SPAL (%) (X2.2), solid waste (X3) is composed of two indicators are not having trash (%) (X3.1), household garbage disposal to vacant land / sea (%) (X3.2); and unhealthy behaviors (X4) consists of three indicators are not doing important CTPS at 5 time (%) (X4.1), Behavior defecation (%) (X4.2), not sorting garbage (%) (X4.3) as well as the risk of sanitation (X) Following the conceptual framework of sanitary risk.




Fig. 3. Conceptual Framework Research

Second CFA analysis performed following steps.

- Multivariate normal testing manifest data
- Standardize the data
- Determining the measurement model
- Getting a parameter estimator measurement model with maximum likelihood approach
- Getting the score factor
- Getting a risk index for each latent variable

4. RESULTS AND DISCUSSION

4.1. Description of Research Variables

Characteristics islands Makasar and Sinjai based on minimum value, maximum, average and standard deviation of each indicator are presented in Table and Figure below.

Table 1. Descriptive Indicators Makasar and Sinjai Islands

Indicator	Minimum	Maximum	Mean	StDev
X _{1,1}	5	100	58.69	31.12
X _{1,2}	0	15	2.25	4.01
X _{1,3}	0	100	39.88	39.63
X _{1,4}	0	100	58.44	33.7
X _{1,5}	0	44	9.38	12.14
X _{1,6}	10	92	44.13	30.87
X _{2,1}	0	85	56.75	22.45
X _{2,2}	38	100	83.31	17.92
X _{3,1}	10	97	66.81	26.21
X _{3,2}	1	100	83.75	30.89
X _{4,1}	54	100	93.38	11.9
X _{4,2}	0	74	50.13	20.22
X _{4,3}	73	100	95.31	7.21

As for seeing the characteristics of outlier data can be viewed via the boxplot in Figure below.




Fig. 4. Boxplot Indicators Research

Table 1 shows the average percentage of the smallest shows that the majority of households in the islands have used the scope of use unprotected sources of drinking water (%) (X2.1) of 2.25 percent with a standard deviation 4.01 percent, using the physical quality of water does not qualify (%) (x1.5) for 9.38 percent with a standard deviation 12.14 percent. This suggests that there are outliers in the variable archipelago because heterogeneous and supported also in Figure 4, ie islands Lumu-Lumu, Samalona and Lae-lae. In addition to the largest average percentage shows that the majority of households in the islands do not sort the waste (%) (X4.3) of 95.31 percent with a standard deviation 7.21 percent, not doing CTPS at 5 critical time (%) (X4.1) of 93.38 percent with a standard deviation 11.9 percent. This suggests that unhealthy behaviors in the islands homogeneous, but when seen in Figure 4, there are islands of outliers, Bonetambung and Katindoang.

4.2. Modeling 2CFA Sanitary Risks

CFA modeling requires the covariance-based multivariate normal distribution of data. This method will produce a good parameter estimates if the data meet the assumption of multivariate normal. Previous normal multivariate testing will be done to the data.

- H₀: Multivariate normal distribution of data
- H₁: The data is not normally distributed multivariate

Table 2. Multivariate Normal Testing

Indicator	skew	c.r.	kurtosis	c.r.
X1.1	-.500	-.817	-.910	-.743
X1.2	2.255	3.682	4.497	3.672
X1.3	.340	.555	-1.667	-1.361
X1.4	-.557	-.909	-.954	-.779
X1.5	1.969	3.216	2.828	2.309
X1.6	.222	.363	-1.623	-1.325
X2.1	-1.024	-1.672	.503	.411
X2.2	-1.328	-2.169	.875	.715
X3.1	-.883	-1.443	-.292	-.238
X3.2	-1.804	-2.946	1.718	1.403
X4.1	-2.470	-4.034	5.634	4.600
X4.2	-.992	-1.619	.371	.303
X4.3	-2.005	-3.274	3.717	3.035
Multivariate			.693	.070

Table 2, shows that the value of CR Multivariate.07 located between the value -1.96 up to 1.96, the multivariate normal distribution of data. Furthermore 2CFA modeling presented in Figure below.




Fig. 5. Sanitation Risk Measurement Model

Based on Figure 5, shows that the two (2) the criteria used to judge the worth / absence of a model turned out to declare Both are CMIN/DF and RMSEA, as well as the three criteria stated quite well that probability, TLI and CFA. It can be said that the measurement model for 2CFA acceptable, which means there is a match between the model with the data.

4.3. Sanitation Risk Index

To obtain a sanitary risk index first sought first factor scores of each latent variable, then according to the equation (3) is produced Sanitation Risk Index are presented in the following table.

Table 3. Latent Variable Index

Island	Index				
	PTS	LP	LC	SA	Sanitation Risk
Langkai	0.935	0.845	0.987	0.892	1.030
Lumu-lumu	1.045	1.008	1.234	1.053	1.808
Bonetambung	1.177	1.051	1.090	1.056	0.615
Barrang Caddi	1.050	1.274	0.437	0.878	-1.349
Kodingareng	0.926	0.891	1.166	1.086	1.956
Barrang Lompo	0.818	0.927	1.195	1.171	2.675
Samalona	0.495	0.758	1.008	0.971	3.145
Lae-lae	0.765	0.825	0.899	0.932	1.459
Burung Loe	1.073	0.737	1.469	1.166	2.397
Katindoang	1.172	1.135	0.902	0.941	-0.096
Kodingareng	1.077	1.094	0.889	1.020	0.286
Liang-liang	1.203	1.156	0.865	0.922	-0.383
Kambuno	0.764	0.584	1.151	0.880	2.247
Kanalo 1	1.238	1.399	1.129	1.117	0.908
Kanalo 2	1.055	0.799	1.131	0.939	1.039
Batang Lampe	1.207	1.516	0.446	0.977	-1.738

Table 3 shows that the largest sanitary risk index respectively - also on islands Samolona, Barrang Lompo, Bird Loe, Kodingareng, Kambuno and Lumu-Lumu. This is because most of the islands Samolona and Kambuno, do not have latrines (%) (X2.1), Not having SPAL (%) (X2.2) and is also supported by LC Index - his. For Barrang islands Lompo, Bird Loe also largely not have latrines (%) (X2.1), Not having SPAL (%) (X2.2), Coverage Using clean water sources are not protected (%) (X1.1), Coverage use unprotected sources of drinking water (%) (X1.2), scarcity of clean water (%) (X1.3), scarcity of drinking water (%) (X1.4), the physical quality of water does not qualify (%) (x1.5) and the distance to the water source pollutant source ≤ 10 meters (%) (X1.6), it is also supported by LC index and index SA.

5. CONCLUSION

The results show that the approach 2CFA sanitary risk measurement model is fit model, and the dominant indicator in the form of sanitary risk respectively are solid waste, unhealthy behaviors, wastewater and ekskreta, as well as a source of clean water and drinking. The highest risk index sanitation on some islands, namely Samolona, Barrang Lompo, Bird Loe, Kodingareng, Kambuno and Lumu-Lumu. So there needs to be the socialization of latrine ownership, ownership SPAL, use of clean and drinking water sources are not protected, the scarcity of clean water and drinking, physical quality of water does not qualify and the distance to the water source pollutant source ≤ 10 meters.

REFERENCES

1. Department of Economic and Social Affairs. 2010. Trend in Sustainable Development: Small Island Developing States (SIDS). Division for Sustainable Development United Nations.
2. Tiempo. (1993). Small island characteristics and their constraints on sustainable development. Online Tiempo Climate Cyberlibrary [Online]. Available at: <http://www.cru.uea.ac.uk/cru/tiempo/issue10/characts.htm>. [Diakses 2 Februari 2014].
3. Undang-Undang Republik Indonesia Nomor 27. (2007). *Pengelolaan Wilayah Pesisir dan Pulau-Pulau Kecil*. Jakarta: UUD RI.
4. N. Rusdi Hidayat, Suhadak, Darminto, Siti Ragil Handayani, Bambang Widjanarko Otok, (2014)., Measurement Model of Service Quality, Regional Tax Regulations, Taxpayer Satisfaction Level, Behavior and Compliance Using Confirmatory Factor Analysis. *World Applied Sciences Journal* 29 (1): 56-61, ISSN 1818-4952 © IDOSI Publications, 2014
5. Hair, J. F., Black, W. C., Babin, B. J., dan Anderson, R. E. (2010). *Multivariate Data Analysis 7th*. Pearson Prentice Hall.
6. Bollen, K. (1989). *Structural Equations With Latent Variables*. New York: John Wiley & Sons, Inc.
7. Calado H, Quentela A, Porteiro J. (2007). Integrated Coastal Zone Management Strategies on Small Islands. *Journal of Coastal Research*, Special Issue50: 125-129.
8. Carr, R, ed. 2001. Excreta-related infections and the role of sanitation in the control of transmission. In: Bartram LFaJ. *Water Quality: Guidelines, Standards and Health*. London: IWA Publishing. p. 90 - 107.
9. Navarro, R. G. (2011). Improving sanitation in coastal communities with special reference to Puerto Princesa, Palawan Province, Philippines. Student Research. School of Architecture. Montreal: McGill University. [Online]. Available at: <http://www.mcgill.ca/mchg/student/sanitation/>. [Diakses 4 Februari 2014].
10. UNESCO. (2008). Sanitation Facts and Figures.
11. Furness, L. (1996). Assessment of spacing between sanitation facilities and wells on Lifuka Island, Kingdom of Tonga. Groundwater Pollution Study. Report on the UNESCO IHP Project 3.
12. Marjoram, T. (1983). Pipes and pits under the palms: water supply and sanitation in the South Pacific. *Waterlines*. Vol. 2. No. 1. p.16.
13. Endang Setiawati, Suprihanto Notodarmojo, Prayatni Soewondo, Agus Jatnika Effendi, Bambang Widjanarko Otok, (2013), Infrastructure development strategy for sustainable wastewater system by using SEM Method

- (Case study Setiabudi and Tebet Districts, South Jakarta), *The 3rd International Conference on Sustainable Future for Human Security SUSTAIN 2012, Procedia Environmental Sciences 17 (2013) 685 – 692, Elsevier.*
14. Daud, A. (2007). Aspek Kesehatan Penyediaan Air Bersih. Makassar: CV. Healthy and Sanitation.
 15. Sandy, C & Ouano, E.A.R.(1990). *Surface Water Drainage in Urban Area. In Poor Die Young: Housing and Health in Third World Cities.* London: Earthscan Publications. p.159
 16. Santoso, I.B and Otok, B.W (2014). Determination of sample size for evaluation Greenspace using the cumulative concentration Levels of carbon dioxide in ambient air. *International Journal of Academic Research.* part A; 2014; 6(1), 161-165.