

Antioxidant Activities of Ethyl Acetic Extract of Endophytic Fungi from *Caesalpinia sappan* L. and *Eucheuma* sp.

by Mashuri Masri

Submission date: 09-Apr-2019 09:58AM (UTC+0700)

Submission ID: 1108666281

File name: Endophytic_Fungi_from_Caesalpinia_sappan_L_and_Eucheuma_sp.pdf (205.07K)

Word count: 3335

Character count: 18268

Antioxidant Activities of Ethyl Acetic Extract of Endophytic Fungi from *Caesalpinia sappan* L. and *Eucheuma* sp.

H HAFSAN*, EKA SUKMAWATY, MASHURI MASRI, ISNA RASDIANAH AZIZ, SITI LATIFAH WULANDARI

Department of Biology, Universitas Islam Negeri Alauddin Makassar, Gowa, Indonesia. 92113

E-mail:hafsan.bio@uin-alauddin.ac.id, (m): +628114448781

Received: 11.06.18, Revised: 11.07.18, Accepted: 11.08.18

ABSTRACT

Antioxidants act as an antidote to free radicals in the body and prevent various diseases caused by oxidative stress. Endophytic fungi currently assuming as a promising source of antioxidant compounds to replace synthetic antioxidants. This study was aimed to determine the antioxidant activity of endophytic fungi isolated from sappan wood (*Caesalpinia sappan* L.) and Macro algae *Eucheuma* sp., to determine the antioxidant activity of the consortium both of fungal isolates. Antioxidant activity was determined by free radical reduction method using DPPH. The results showed that five isolates of endophytic fungi isolated from sappan wood were *Curvularia lunata*, *Penicillium* sp., *Chrysosporium* sp., *Trichoderma* sp. and *Geotrichum candidum*. One isolate from *Eucheuma* sp., namely *Aspergillus* sp. The highest antioxidant activity was found in *Trichoderma* sp. and *Aspergillus* sp. which IC_{50} point were 46.354 and 38.64 respectively. The consortium was able to increase the antioxidant activity of single isolates which the lowest IC_{50} .

Keywords: antioxidant, *Caesalpinia sappan* L., DPPH, *Eucheuma* sp., endophytic fungi

INTRODUCTION

Free radicals such as Reactive Oxygen Species (ROS) are produced in living organisms either from normal essential metabolic processes or from external sources e.g. x-ray exposure, air pollutants, and industrial chemicals[1]. As unpaired electrons, free radicals are very reactive, their excess production can cause damage to all biological macromolecules such as DNA, proteins, and lipids, lead to cell damage and then manifest of pathological conditions[2]. Severe oxidative damage from free radicals is associated with various diseases including cancer, inflammation, aging, and neurodegenerative diseases [3][4]. Antioxidants function as a defense factor against free radicals in the body[5]. The use of synthetic antioxidants to prevent free radical damage causes side effects such as damage DNA and protein as well as are toxic. Therefore, that natural antioxidants and free radical reducers are needed[6]–[8]. Endophytic fungi are spread out nearly 300,000 plant species and carry out most of their life cycles on healthy plant tissues without causing disease on their host [9][10]. At present, endophytes are seen as source both of extraordinary and ecofriendly bioactive natural products. This group of organisms produces bioactive compounds that may provide defense for host plants against pathogens, and some of these compounds are useful for new drugs[11]. A large number of new active compounds can be obtained from endophytic fungi such as anticancer and antioxidants [12][13]. Sappan wood (*Caesalpinia sappan* L.) is a plant that is widely found in Indonesia and empirically has been known to have many healing properties. This bark is often

consumed by the public as a health drink. Sappan wood contains compounds of brazilin ($C_{16}H_{14}O_5$), sappanin ($C_{12}H_{12}O_4$), brazilin, and essential oils such as D- α -phellandrene, gallic acid, and resin. Sappan wood has a higher antioxidant than commercial antioxidants (BHT and BHA) so that it has the potential as an antidote to free radicals[14]. Likewise marine species such as Macroalgae is known to produce diverse active compounds. Macroalgae *Eucheuma* sp. used as an alternative microbial algae symbiont to be able to take advantage from it and become one of the most effective solutions. However, a lack of information regarding the antioxidant activity of endophytic fungi on *Eucheuma* sp. needs a research study to determine it. Considering that both plants come from different habitats, it is interesting to know the changes in antioxidant activity when endophytic fungi from both are grown in one medium.

Material and Methods

Isolation of endophytic fungi

Samples of *Caesalpinia sappan* L. and *Eucheuma* sp. as the source of the isolate were washed and then sterilized the surface by soaking in 70% alcohol hypochlorite solution for 1 minute, and then rinsed with sterile distilled water. Following this step the leaves were crushed using mortar and pestle and spread on PDA media. While the stem samples were cut into small pieces and sown directly in PDA media. Incubation was carried out for one week at ambient temperature. Colonies growing from plant tissues were further isolated and purified. Collections of isolates were stored in a refrigerator before further

testing.

Identification of endophytic fungi

The fungus culture is taken aseptically using an aseptic needle then placed on top of a glass of objects that have been dripped with aquadest, dried with tissue, next step to obtain its microscopic characteristics, viewed under a microscope. Identification was done by matching the characteristics of fungi that obtained from observations with Barnett and hunter identification book [15].

Extraction of fermentation results

About 100 ml of isolates-fermented was separated from biomass with filtrate in Erlenmeyer. Biomass was dried first with an oven. The filtrate was extracted 2 times with 50 ml of ethyl acetate using a separating funnel, and then aerated until there was separation between the ethyl acetate and water phases. The ethyl acetate phase was collected then dried. The results of extraction of the dried filtrate were then tested for antioxidant activity.

Test of antioxidant activity extract of filtrate

Antioxidant activity was determined by free radical reduction methods using DPPH [16]. Test solution and positive control were put into each test tube and then added 1 ml of DPPH solution then added to 5 mL of methanol analysis pro, then homogenized with a blank solution, test solution and positive control were incubated for 30 minutes immediately. Absorbance equipped with 517 nm using a UV-vis spectrophotometer. Percentage of obstacles is calculated by the formula: percentage obstacle = (absorptive absorption-sample absorption) / (blank uptake) x 100% IC_{50} is calculated by entering 50% resistance in the linear regression equation. The IC_{50} is a value that shows the concentration of the test sample needed to reduce 50% of DPPH free radicals. Endophytic fungi from both plants were then made a consortium. The consortium is carried out which combines fungi which produce the highest and lowest antioxidant activity.

Making the growth curve of endophytic fungus consortium

Growth curves were made based on the results of fermented dry biomass. The workings of creating a growth curve are first cutting the filter paper with a circular shape and weighing it with an analytic balance sheet, and then the weighed filter paper was used to separate biomass and filtrate. Biomass obtained in the oven for 24 hours and recalculated after filter paper at room temperature. The results of the initial scales without biomass filter paper reduced by paper scales with biomass already in the oven are the weight of biomass inserted into the growth curve. This is applied from day 2 to day 20 to determine the phases that occurs during fungus fermentation.

Results and Discussion

The isolation results of endophytic fungi from *Caesalpinia sappan* L. were obtained from five fungi isolates consisting of *Curvularia lunata*, *Penicillium*

sp., *Chrysosporium* sp., *Trichoderma* sp., and *Geotrichum candidum*. While the isolation results from *Eucheuma* sp. obtained one isolate *Aspergillus* sp. Identification was applied both macroscopically and microscopically. Macroscopic observation consist of observing surface color, reverse color, texture, topography, exudate droplets, radial lines, and concentric circles. Whereas microscopic observations include observing the conidia and hyphae shape. Macroscopic and microscopic observations as seen in table Figure 1. Furthermore, endophytic fungus isolates were fermented in GDP media which were substituted with chloramphenicol. According to Pelczar and Chan [17], PBD is commonly used for fungi cultivation because it contains rich nutrient derived from potato and dextrose decoction which induces fungal sporulation, dye production, and fertile fungus growth. Fermentation was carried out for 7-10 days to get the results of the metabolism of the fungus to be extracted. On the seventh day, the growth of mycelia in the fluid and color of GDP began to change. This is presumably due to the association between the biosynthesis of secondary metabolites and the sporulation process in endophytic fungi. Sporulation results in the formation of secondary metabolites. Therefore, extraction was carried out using ethyl acetate solvent. Ethyl acetate is a semi-polar organic solvent. This selection is expected to be able to extract more polar and non-polar components and easily be separated by polar GDP liquid. The extract obtained was concentrated with a vacuum evaporator and tested for antioxidant activity using DPPH method with UV-vis ray spectrophotometry. Test results are present in Table 1. Among several antioxidant tests, the DPPH free radical scavenging test is the most widely used test and considering the most accurate screening method used to evaluate antioxidant activity. In addition, DPPH is not affected by metals and enzyme inhibition [18]. The reaction visualized as a change in color from purple to yellow. *Trichoderma* sp. isolates from sappan wood showed very strong antioxidant activity with IC_{50} point by 46.354 as well as endophytic fungi isolates *Aspergillus* sp. isolated from *Eucheuma* sp. with IC_{50} point by 38.64. However, both isolates are still lower than vitamin C as a control of it. Natural antioxidants derived from plants are associated with secondary metabolites, especially phenolic and flavonoids. Previous research on qualitative testing of ethanol extract from endophytic fungi reported flavonoids in extracts [19]. Some fungi have also been reported to produce compounds that have potential as antioxidant [20] such as *Aspergillus nidulans* with *A. oryzae* [21] and *A. niger* with *Fusarium oxysporum* [22]. Furthermore, a consortium was conducted between endophytic fungi isolates with very strong antioxidant activity and fungal isolates whose activity was low. The consorted isolates were *Trichoderma* sp. with *Chrysosporium* sp. and *Aspergillus* sp. and *Chrysosporium* sp. Antagonistic

test used to see the ability of endophytic fungi to live together on PDA media before being fermented in GDP media. The results of the antagonist test showed that the endophytic fungi isolate *Trichoderma* sp. with *Chrysosporium* sp. does not show antagonistic

interaction. Whereas in the consortium *Aspergillus* sp. and *Chrysosporium* sp. shows the hyphae of *Aspergillus* sp. dominate so that *Chrysosporium* sp. cannot grow properly (Figure 2).




Figure 1. Macroscopic and microscopic observation of samples, 1a: *Curvularia lunata* colonies in petri dishes; 1b: conidia *Curvularia lunata*. 2a: *Penicillium* sp. colonies in petri dishes; 2b: conidia *Penicillium* sp. 3a: *Chrysosporium* sp. colonies in petri dishes, 3b: conidia *Chrysosporium* sp. 4a: colonies of *Trichoderma* sp. in petri dishes; 4b: conidia *Trichoderma* sp. 5a: colony of *Geotrichum candidum* in petri dish; 5b: conidia *Geotrichum candidum*. 6a: colonies of *Aspergillus* sp. in petri dishes; 5b: conidia *Aspergillus* sp. (1-5: *Caesalpinia sappan* L. endophytic fungi; 6: *Eucheuma* sp. endophytic fungi)




Figure 2. The consortium of a: *Trichoderma* sp. with *Chrysosporium* sp. b: *Aspergillus* sp. and *Chrysosporium* sp.

Table 1. IC₅₀ values of endophytic fungal isolates

No.	Fungus Isolates	IC ₅₀ point	Remarks
1	<i>Curvularia lunata</i>	91.926	strong
2	<i>Penicillium</i> sp.	161.60	weak
3	<i>Chrysosporium</i> sp.	219.25	weak
4	<i>Trichoderma</i> sp.	46.354	very strong
5	<i>Geotrichum candidum</i>	219.81	weak
6	<i>Aspergillus</i> sp.	38.64	very strong

*IC₅₀ Vitamin C: 3.03 (very strong)

The growth curve of the endophytic fungus consortium was made to see the growth rate of endophytic fungus consortium and determine the growth phases of endophytic fungi and look for the

stationary phase of endophytic fungus consortium to see the optimum production time of its secondary metabolites (Figure 3).




Figure 3.The growth curve of endophytic fungus consortium. a:*Trichoderma* sp. with *Chrysosporium* sp.; b:*Aspergillus* sp. and *Chrysosporium* sp.

According to Srikandaceet *al.*[23]in the stationary phase, it is assuming that endophytic fungus cells are more resistant to extreme conditions such as cold and hot temperatures, secondary chemical and metabolic radiation produced by the fungus itself. Limitations of the main source of synthesis include sugar as a carbon source and protein as a source of amino acids or nitrogen. This can cause the release

of substances resulting from the catabolism process which are secondary metabolic. The antioxidant activity of the two consortia was tested at the end of the stationary period. The antioxidant activity of IC_{50} *Trichoderma* sp. with *Chrysosporium* sp. is very strong with IC_{50} of 47.807, while IC_{50} is 50.318. *Aspergillus* sp. with *Chrysosporium* sp. and relatively strong (Table 2).

Table 2.Antioxidant activity of endophytic fungus consortium

No.	Fungus consortium	IC_{50} point	remark
1	<i>Trichoderma</i> sp.+ <i>Chrysosporium</i> sp.	47.807	strong
2	<i>Aspergillus</i> sp. + <i>Chrysosporium</i> sp.	50.318	weak

Trichoderma sp. consortium with *Chrysosporium* sp. is not different from the antioxidant activity of single isolate *Trichoderma* sp. which is classified as very strong. Although there was a decrease in IC_{50} point compared to single isolates. However, this consortium was able to increase the antioxidant activity and IC_{50} of *Chrysosporium* sp. single isolates which previously were classified as weak. This is possible because the two isolates of the fungus can

live side by side without showing antagonistic interactions as shown in Figure 2. This also indicates that there is no competition for secondary metabolites produced by the two isolates. Unlike the consortium of *Aspergillus* sp. with *Chrysosporium* sp. which showed a decrease in antioxidant activity compared to the single isolate *Aspergillus* sp., although this consortium can increase the antioxidant activity of single isolates of *Chrysosporium* sp.

According to Septiana et al. [24] this digression was due to the interaction of endophytic fungi that were not optimal in producing secondary metabolites that act as antioxidants. Based on the test results of antioxidant activity produced from secondary metabolites, all consortiums of endophytic fungal isolates can be utilized as one of the strengths of endophytic fungi usage in the field of health microbiology, especially as a substitute for synthetic drugs for diseases caused by free radicals.

Conclusion

Five isolates of endophytic fungi were isolated from sappan wood (*Caesalpinia sappan* L.) consist of *Curvularia lunata*, *Penicillium* sp., *Chrysosporium* sp., *Trichoderma* sp. and *Geotrichum candidum*. In addition, one isolate from *Eucheuma* sp. namely *Aspergillus* sp. The highest antioxidant activity was found in *Trichoderma* sp. and *Aspergillus* sp. with IC_{50} point were 46.354 and 38.64, respectively. The consortium was able to increase the antioxidant activity of single isolates which had have the lowest IC_{50} .

References

1. M. Valko, D. Leibfriz, J. Moncol, M. T. D. Cronin, M. Mazur, and J. Telser, "Free radicals and antioxidants in normal physiological functions and human disease," *Int. J. Biochem. Cell Biol.*, vol. 39, no. 1, pp. 44–84, 2007.
2. B. Halliwell, "Role of free radicals in the neurodegenerative diseases: therapeutic implications for antioxidant treatment," *Drugs Aging*, vol. 7, no. 1, p. 67, 2001.
3. S. J. Hur, S. Y. Lee, Y. C. Kim, I. Choi, and G. B. Kim, "Effect of fermentation on the antioxidant activity in plant-based foods," *Food Chem.*, vol. 160, pp. 346–356, 2014.
4. F. Hecht, C. F. Pessoa, L. B. Gentile, D. Rosenthal, D. P. Carvalho, and R. S. Fortunato, "The role of oxidative stress on breast cancer development and therapy," *Tumor Biol.*, vol. 37, no. 4, pp. 4281–4291, 2016.
5. D. S. Arora and P. Chandra, "Assay of antioxidant potential of two *Aspergillus* isolates by different methods under various physio-chemical conditions," *Brazilian J. Microbiol.*, vol. 41, no. 3, pp. 765–777, 2010.
6. C. F. Chi, B. Wang, Y. Y. Deng, Y. M. Wang, S. G. Deng, and J. Y. Ma, "Isolation and characterization of three antioxidant pentapeptides from protein hydrolysate of monkfish (*Lophius litulon*) muscle," *Food Res. Int.*, 2014.
7. B. Wang, Y. D. Gong, Z. R. Li, D. Yu, C. F. Chi, and J. Y. Ma, "Isolation and characterisation of five novel antioxidant peptides from ethanol-soluble proteins hydrolysate of spotless smoothhound (*Mustelus griseus*) muscle," *J. Funct. Foods*, vol. 55, pp. 222–228, 2014.
8. N. Radulović, V. Stankov-Jovanović, G. Stojanović, A. Šmelcerović, M. Spittler, and Y. Asakawa, "Screening of in vitro antimicrobial and antioxidant activity of nine *Hypericum* species from the Balkans," *Food Chem.*, 2007.
9. R. X. Tan and W. X. Zou, "Endophytes: A rich source of functional metabolites," *Nat. Prod. Rep.*, vol. 18, pp. 448–459, 2001.
10. A. H. Aly, A. Debbab, and P. Proksch, "Fungal endophytes: Unique plant inhabitants with great promises," *Appl. Microbiol. Biotechnol.*, vol. 90, no. 6, pp. 1829–1845, 2011.
11. R. Shameem, M. Lacouture, and S. Wu, "Incidence and risk of high-grade stomatitis with mTOR inhibitors in cancer patients," *Cancer Invest.*, vol. 54, no. 1, pp. 70–77, 2015.
12. J. Bérdy, "Thoughts and facts about antibiotics: Where we are now and where we are heading," *J. Antibiot. (Tokyo)*, vol. 103, no. 1, pp. 15–21, 2012.
13. B. F. Pan, X. Su, B. Hu, N. Yang, Q. Chen, and W. Wu, "*Fusarium redolens* 6WBY3, an endophytic fungus isolated from *Fritillaria unibracteata* var. *wabuensis*, produces peimisine and imperialine-3 β -d-glucoside," *Fitoterapia*, vol. 103, pp. 213–221, 2015.
14. S. Badami, S. Moorkoth, S. R. Rai, E. Kannan, and S. Bhojraj, "Antioxidant Activity of *Caesalpinia sappan* Heartwood," *Biol. Pharm. Bull.*, vol. 26, no. 11, pp. 1534–1537, 2003.
15. P. I. Physiology et al., "Illustrated Genera of Imperfect Fungi," in *Transactions of the British Mycological Society*, Minneapolis, Burgess Publishing Co., 1955, p. 225.
16. V. Saraswaty, C. Risdian, T. A. Budiwati, and T. Mozef, "Aktivitas antioksidan dari kombinasi ekstrak etanol kulit manggis, daun sirsak dan daun sirih merah," in *Proceeding Seminar Ilmu Pengetahuan Teknik*, Bogor: LIPI Indonesia, 2013, pp. 196–200.
17. M. J. Pelczar, E. C. Chan, and N. R. Krieg, "Microbiology. 5th ed," *Tata Mc Graw Hill Publ.*, 1993.
18. M. Yadav, A. Yadav, and J. P. Yadav, "In vitro antioxidant activity and total phenolic content of endophytic fungi isolated from *Eugenia jambolana* Lam.," *Asian Pac. J. Trop. Med.*, vol. 7, no. 1, pp. 256–261, 2014.
19. V. Nagda, A. Gajbhiye, and D. Kumar, "Isolation and characterization of endophytic fungi from *Calotropis procera* for their antioxidant activity," *Asian J. Pharm. Clin. Res.*, vol. 10, no. 3, pp. 254–258, 2017.
20. T. Y. Song and G. C. Yen, "Antioxidant Properties of *Antrodia camphorata* in Submerged Culture," *J. Agric. Food Chem.*, vol. 50, no. 11, pp. 3322–3327, 2002.
21. Y. Lu, C. Chen, H. Chen, J. Zhang, and W. Chen, "Isolation and identification of endophytic fungi from *Actinidia macrosperma* and investigation of their bioactivities," *Evidence-Based Complement. Altern. Med.*, vol. 2012, no. 2012, pp. 1–8, 2012.
22. M. Govindappa, N. Bharath, H. B. Shruthi, and G. Santoyo, "In vitro Antioxidant Activity and Phytochemical Screening of Endophytic Extracts of *Crotalaria pallida*," *Free Radicals Antioxidants*, vol. 1, no. 3, pp. 79–86, 2011.

23. Y. Srikandace, Y. Hapsari, and P. Simanjuntak, "Seleksi Mikroba Endofit Curcuma Zedoaria dalam Memproduksi Senyawa Kimia Antimikroba," *J. Ilmu Kefarmasian Indones.*, vol. 5, no. 2, pp. 77–84, 2007.
24. E. Septiana, Bustanussalam., F. Rachman, Y. Hapsari, and P. Simanjuntak, "Potensi Ekstrak Kapang Endofit Asal Rimpang Kunyit sebagai Antimalaria dan Antioksidan," *J. Kefarmasian Indones.*, vol. 7, no. 1, pp. 1–9, 2017.

Antioxidant Activities of Ethyl Acetic Extract of Endophytic Fungi from *Caesalpinia sappan* L. and *Eucheuma* sp.

ORIGINALITY REPORT

6%

SIMILARITY INDEX

1%

INTERNET SOURCES

6%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Aliki Papadopoulou, Konstantinos Petrotos, Dimitrios Stagos, Konstantinos Gerasopoulos et al. "Enhancement of Antioxidant Mechanisms and Reduction of Oxidative Stress in Chickens after the Administration of Drinking Water Enriched with Polyphenolic Powder from Olive Mill Waste Waters", *Oxidative Medicine and Cellular Longevity*, 2017

2%

Publication
- 2

M. Murali, C. Mahendra, P. Hema, N. Rajashekar, A. Nataraju, M. S. Sudarshana, K. N. Amruthesh. "Molecular profiling and bioactive potential of an endophytic fungus isolated from : a medicinal plant ", *Pharmaceutical Biology*, 2017

1%

Publication
- 3

"In vitro antioxidant and antibacterial activity of endophytic fungi isolated from *Mussaenda luteola*", *Journal of Applied Pharmaceutical Science*, 2017

1%



journals.plos.org
Internet Source

1%

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%