



# THE 5th INTERNATIONAL CONFERENCE ON GREEN TECHNOLOGY

“Science and Technology in Islamic Perspective: Synergy and Theirs Contribution to Empower Nations”



## PROCEEDING

Issue: 5 / 2015

7th-8th NOVEMBER 2014  
Soekarno Hatta Building, 5th floor  
Maulana Malik Ibrahim State Islamic University  
of Malang

**Scientific.Net**  
Materials Science and Engineering

**ULUL ALBAB**  
Jurnal Teknik Kimia

**PENELITIAN HAYATI**  
Jurnal of Biological Researches

**ALCHEMY**  
Journal of Chemistry

**CAUCHY**  
Jurnal Matematika Murni dan Aplikatif

**Journal of Islamic Architecture**

**NEUTRINO**

**JURNAL BIOLOGI**  
*et - Husak*

**MATICS**



**PROCEEDING OF THE 5<sup>th</sup> INTERNATIONAL CONFERENCE  
ON GREEN TECHNOLOGY**

*Science and Technology in Islamic Perspective:  
Synergy and Their Contribution to Empower Nations*

**Copyright © 2015**

**REVIEWER**

Prof. Drs. Sutiman B. Sumitro, SU., D.Sc  
Assoc. Prof. Dr. Akira Kikuchi  
Assoc. Prof. Nangkula Utaberta, Ph.D  
Sr. Dr Md. Azree Othuman Mydin  
Dr. Agus Salim, M.Si  
Dr. drh. Bayyinatul Muchtaromah, M.Si  
Dr. H. Zainuddin, M.A.  
Dr. Eko Budi Minarno, M.Pd  
Dr. Cahyo Crysdiyan, MCS  
Dr. Sri Harini, M.Si  
Dr. Agung Sedayu, M.T

**EDITOR**

M. Jamhuri, M.Si  
Fachrur Rozi, M.Si  
Ismail

**FACULTY OF SCIENCE AND TECHONOLY  
MAULANA MALIK IBRAHIM STATE ISLAMIC UNIVERSITY OF MALANG  
MALANG**

© Hak Cipta Dilindungi oleh Undang-undang  
All Right Reserved

# CONTENT

## **SCIENCE AND ISLAM**

*Science of Islam*

Roikhan Mochamad Aziz

## **NATURAL SCIENCE**

*Screening of Antagonistic Activity of Endofitic Fungi from Toona Sinensis Against Candida Albicans*

Anggita R. Hafsari, Isma Asterina

*Sublethal Effect of Three Types of Herbicides to Growth and Reproduction of Common Earthworm (Lumbricus rubellus)*

Ida Kinasih, Astuti Kusumorini, Purnawarman

*The Potential of Protective Trees as CO<sub>2</sub> Absorber and Carbon Storage in Malang Highway*

Bagus Setiawan and Dwi Suheriyanto

*Moringa Leaves Extract Versus Glucose Levels and Cholesterol Blood Total*

Saleh Hidayat, Susi Dewiyeti, Donni Yusuf, and Sintiya Maisaroh

*Clove Oil and Neem Seed Extract as Botanical Fungicides Control Rust Disease (Phakopsora pachyrhizi) on Soybean*

Sumartini

*Lipid Production and Growth Performance of Chlorella sp. Grown in Tofu Waste Water*

Evika Sandi Savitri, Romaidi, Ruri Siti Resmisari, Siti Farichah, Hasanuddin

*Microcosm Scale Models of Petroleum Bioremediation in Mangrove Areas Using Indigenous Hydrocarbonoclastic Bacteria*

Hary Widjajanti, Moh. Rasyid Ridho, Munawar

*Correlation and Path Analysis Between Yield and Yield-Related Traits of Twelve Selected Genotypes of Orange-Fleshed Sweet Potato*

Wiwit Rahajeng and St. A. Rahayuningsih

*Performance of Methylobacterium spp. for Soybean Seed Invigoration*

Ratri Tri Hapsari, Eny Widajati, Selly Salma, Maryati Sari

*Ethnobotany Parasite Tea (Scurulla atropurpurea (Bl.) Dans.) at Wonorejo Village Community, District Lawang Malang.*

Furqon Al Muzakki and Siti Na'imatun Niswah

*Growth of Black Soldier Fly (Hermetia illucens) Larvae and Mealworm (Tenebrio molitor) during Bioconversion of Rice Straw and Cassava Peel*

Ramadhani Eka Putra, Rizal Jam Jam, Ateng Supriatna, Robert Manurung, and Ida Kinasih



*Response Pigments Anthocyanin from Local Resources of Various Yoghurt Quality*

Moch. Wachid, Zaidur Rizaq, Elfi Anis Saati and Warkoyo

*Isolation and Characterization of Lactic Acid Bacteria from Dangke Waste Production*

Hafsan Muhammad Halifah Mustami

*Preparation, Modification and Characterization of TiO<sub>2</sub>/Activated Natural Zeolite Malang Composite in Weight Ratio 4:9 (w/w)*

Suci Amalia, Susi Nurul Khalifah, Aisyah Erlynataand Mohammad Yusuf

*Solid State Reaction-Sonication Synthesis and Characterization of Vanadium (V) Doped TiO<sub>2</sub> Photocatalyst*

Desy Maulina, Rachmawati Ningsih, Nur Aini

*The Determination of Reaction Order and Activation Energy of Glucose Isomerization in Presence of Malang Natural Zeolite Impregnated by Tin with Temperature Variation*

Pamungkas Dwi Putri W. , Suci Amalia, Susi Nurul Kholifah

*Identification of Chlorellasp. Microalgae Fatty Acid by Gas Chromatography-Mass Spectrophotometer*

A. Ghanaim Fasya., Diah Kumalasari, Vera Susanti, Anik Kholifatuz Zahra', Tri Kustono Adi, Ahmad Hanapi, Anik Maunatin

*Fructooligosaccharides Yacon Tubers [Smallanthus sonchifolia (Poepp.et Endl.) H.Robinson] for Lowering Blood Triglycerides and Total Cholesterol: Length of Storage and Boiling Time Variations*

Prima Retno Wikandari, Leny Yuanita

*Stigmasterol From the Bark of Aglaia Glabrata (Meliaceae)*

Supriadin A. and Supratman U.

*Effect of Cryoprotectant and Explant Type to Recovery of Two Accessions of Dioscorea Alata L. After In Vitro Vitrification*

Ragapadmi Purnamaningsih, Surya Diantina and Sri Hutami

*Phylogenetic Analysis of Endophytic, Pathogenic, and Saprobic Fusaria Indigenous Indonesia based on Internal Transcribed Spacer and Partial Elongation Factor Gene Regions*

Feni Tunarsih, Gayuh Rahayu, Iman Hidayat

*Cytotoxic Activity Extract of Lophatherum Gracile Brongn Leaves and Roots using BSLT (Brine Shrimp Lethality Test) and Identification of The Active Compounds*

Elok Kamilah Hayati, Roihatul Muti'ah, Alfin Hilda Rizqiyah, Selina Purwita Sari

*Extraction of Gelatin from Broiler Chicken Bone using Acid*

Akyunul Jannah and Anik Maunatin

*Primers and Temperature Annealing Optimization of PCR Amplification Process In Resistance Gene Analog (RGA) from Some Banana Cultivars (Musa sp.)*

Siti Mar'atush Sholihah, Evika Sandi Savitri

*Cryopreservation of In Vitro Shoot Tips Dioscorea Esculenta with Vitrification Method*

Surya Diantina, Ragapadmi Purnamaningsih, Sri Hutami

*The Antioxidant Activity and Sensory Quality Quality Characteristic of Hojje Fruit Peel Essential Oil Chicken Sausage Formulation*

Dede Sukandar, Anna Muawanah, Eka Rizki Amelia dan Khilda Fithri Aryani

*Toxicity Test on Artemia salina Shrimp Larvae of Microalgae Chlorella sp. Resltded from Tauge Extract Medium*

Suci Amaliyah, A. Ghanaim Fasya and Romaidi

*Lead (Pb) Analysis in Packaged Candies Using Various Destruction Solution by Atomic Absorbtion Spectroscopy*

Siti Nurul Afifah', Diana Candra Dewi, A. Ghanaim Fasya, and Rachmawati Ningsih

*Phenylalanin and Phenylpiruvate Analysis Using Modified Silica Identification Reagent*

Diana Candra Dewi, M.Si, drg Arief Suryadinata, dr Ana Rahmawati

## **APPLIED SCIENCE AND TECHNOLOGY**

*Optimal Control Wind Energy Conversion System: Classical Mppt Vs Mppt Using Wind Turbulance*

Ahmad Nadhir

*Characterization of Crystal Structure of Hydrogen Storage Keratin-Mg Using Rietveld Analysis*

Azizah Fii Ahliha, Erna Hastuti

*Tannins as Protectans of Protein from Several Source of Feed Materials*

Dini Hardini and Siti Istiana

*Metallothionein Expression on the Gill of Fresh Water Mussel (Anadonta woodiana) Exposed Pb Sub Cronic Level*

Hartati Kartikaningsih, Asus Maisar Suryanto, Diana Arfiati

*Agronomic Changes and Tolerance of Soybean Genotypes to Waterlogged Soil*

Ayda Krisnawati, M. Muchlish Adie, and Didik Harnowo

*Acclimatize And Propagation Orchid Species In Tissue Culture*

Titiek Purbiati and Melia Puspitasari

*Recognition of Plate Characters using Morphology and Correlation Coefficient*

Bain Khusnul Khotimah, Haryanto, Sigit Susanto Putro



*Design of Brake Lining Pads of Non-Asbestos Composite Materials from Coconut Coir Fiber and Aluminium Powder as Filler*

Eko Marsyahyo, Eko Yohanes S., Yafhed Octavianus, Virginia C.W.

*Sugarcane Growth Response to Organic Input and Nitrogen Fertilization on an Inceptisol with Inoculation of *Pontoscolex corethrurus**

Nurhidayati and Abdul Basit

*Non-Linear Analysis Of Hollow Reinforced Concrete Column Square Cross-Section With Various Load Eccentricity And Concrete Strength*

Pingkan Nuryanti, Djoko Sulityo, Bambang Suhendro

*Development of Micro-hydro to Supply Clean Water in The Karangsono Village, Malang Districts*

Machmud, Irfan, Suwignyo

*Evaluation of Resistance of Soybean Genotypes Against Pod Borer, *Etiella zinckenella* Tr. to Support Environmentally Friendly Pest Control*

Marida Santi YIB dan Yusmani Prayogo

*Study of Free Radicals Absorption in Cigarette Smoke on Biofilter Composite Membranes Made of Tobacco and Moringo Leaf*

Nurun Nayiroh, Erika Rani and Agus Mulyono

*The Study of Some Shallot Varieties Production by Liquid Organic Fertilizer Application*

Nurul Istiqomah dan Amik Krismawati

*Management Information System for Parking Application With Barcode Technology*

Rika Yunitarini

*Rosella Processing Technology Introduction in Supporting Diversification of Food Product Development in Donomulyo District, Malang Regency*

Ita Yustina and Farid R. Abadi

*Characteristic of Soybean Maturity and Its Role as Seed Yield Determinant*

M. Muchlish Adie, Ayda Krisnawati, and Suyamto

*STAR: Story-Telling Augmented Reality for Islamic Childhood Education*

Ari Kusumaningsih, Arik Kurniawati

*Information System Arrangement for Tree Management of The Green Lane Road*

Bambang Sulistyantara, Prita Indah Pratiwi

*Introduction To Increase Rice Production Through Superior New Varieties of Rice Fields in The Madiun Regency*

Amik Krismawati and Nurul Istiqomah

*Contribution of Environmental Impact Analysis Study (AMDAL) to The Implementation of Green Development Process*

Titien Setiyo Rini, Miftahul Huda, Agus Purwito

*Effect of Fertilization in Monoculture and Intercropping Systems of Corn with Legumes in The Lowland*

Zainal Arifin, Indriana RD and Nurul Istiqomah

*“Tutup” Flowers ((Macaranga Tanarius (L.) Mull. Arg)) As A Preservative For Fresh Fish*

Fungki Sri Rejeki, Endang Retno Wedowati, and Diana Puspitasari

*Study of Soil Tillage and Mulch to Growth of Weed and Yield of Two Soybean Variety*

Zainal Arifin and Indriana RD

*Horenso Organic-Aquaponic System Using Fish Metabolism Residue In Various Planting Media*

Cecep Hidayat, Farhan Ferdian Ahmad Taofik, Sofiya Hasani

*Product Quality Assessment of Preserved Sweet Potato (Ipoemoea batatas) in Malang Regency, East Java*

Farid R. Abadi dan Ita Yustina

## **MATHEMATICS AND COMPUTATION TECHNOLOGY**

*Simulation Model Of Immune System and Mycobacterium Tuberculosis Infection*

Usman Pagalay, Khairul Umam, and Nurhasanah

*Stability of Jensen Functional Equation on Restricted Domain*

Hairur Rahman, Hilwin Nisa' and Zukhrufun Nadhifa

*An Application of  $\alpha$ -Cut Fuzzy X – S Control Charts Using Fuzzy Trapezoidal Number*

Fitriatuz Zakiyah, Dia Kusumawati, and Fachrur Rozi

*Run Lenght Matrix Feature Extraction and Local Binary Pattern For Face Expression Recognition*

Cucun Very Angkoso

*Electronic Data Interchange For Portal E-Commerce Community By Using Optimistic And Pesimistic Representational State Transfer*

Hermawan and Zamrony Hamzah

*Arabic Letters Pattern Recognition Using Web Camera*

Totok Chamidy, Ririen Kusumawati

*Steganography and Cryptography in Video For Information Security*

Wiwin Styorini, Hendrawan

*Semantic Web Based Search Engine Using Rabin Karp Algorithms in Encyclopedia of Indonesian Herbal Medicine*

Ludvi Andini and A'la Syauqi

*Immersive Tools Bicycle for Human Heart Health Based on Environment Green Technology*

Muhammad Faisal, Yunifa M. Arif, Fahrul Kurniawan, Fressy Nugroho, Hani Nurhayati

**ARCHITECTURE**

*Re-Construction of Taneyan Lanjhang: Observe The Meaning of The Settlement In Madura-Sumenep*

Pudji PratitisWismantara

*Identification of Important Factors Implementation of Green Construction Specifications*

Miftahul Huda and Titien Setiyo Rini

*Stabilization Expansive Soil with Waste of Sand Onyx*

Agus Tugas Sudjianto, Candra Adityaand Muhamad Cakrawala

*Study on The Benefit of Vertical Greenery on Office Building at DKI Jakarta (Case Study at Bank BTPN)*

Yozi Fitri Yeni, Indung Sitti Fatimah, Pingkan Nuryanti

*The Development of Expertise Certification Test Model Based on Regional Model of Competency Standards (RMCS) and Project Based Learning in The Board of Development and Construction Services*

Tri Kuncoro

*Science and Technology In Islamic Perspective: Strengthen National Resilience and Unity of NKRI (An Appropriate Step by Green and Multipurpose Terminal Development)*

Agung Sedayu

**PHARMACY AND MEDICINE**

*Profile Number of Mice Takizoit after Treatment With Alkaloid Fraction of Alstonia scholaris Leaves*

Abdul Hakim

*The Comparison between Ethanol Extract Activity and Ethyl Acetate Pulai Stem Bark (Alstonia scholaris, (l.) R. Br) as Antitoxoplasmosis*

Roudlotul Nadlifah, Begum Fauziyah, Mufasirin

*Effectiveness Test on Alkaloid Extract from Pulai Root (alstonia scholaris l. R.br.) In Increasing the Life of Mice Infected Toxoplasma Gondii Parasite*

Q. Rahmawati; Begum Fauziyah; Mufasirin



## ISOLATION AND CHARACTERIZATION OF LACTIC ACID BACTERIA FROM DANGKE WASTE PRODUCTION

Hafsan, Muhammad Halifah Mustami

Faculty of Science and Technology, Alauddin Islamic State University Makassar, Gowa,  
92113, South Sulawesi, Indonesia Tel: +62-85216478782

Corresponding Author E-mail: [hafsahbio@yahoo.com](mailto:hafsahbio@yahoo.com)

### ABSTRACT

This research was aimed to investigate the characteristics of lactic acid bacteria isolated from Dangke waste production, known as whey. Whey obtained from home industry of Dangke in Enrekang City, Enrekang District South Sulawesi. Isolation and characterization of lactic acid bacteria has been conducted in the Laboratory of Microbiology, Department of Biology, Faculty of Science and Technology UIN Alauddin Makassar, take place in January until March 2014. Isolation and characterization of lactic acid bacteria has been conducted in the Laboratory of Microbiology, Department of Biology, Faculty of Science and Technology UIN Alauddin Makassar. From the whey obtained two isolates of lactic acid bacteria, both Gram-positive rod-shaped bacterium was able to grow on medium MRSA, have colonies round shape, flat edge, convex surface and milky white. Based on biochemical tests, both isolates are non-motile and able to fermenting carbohydrates, especially glucose and sucrose. Negative result on the test indole, lysine test, citrate test, urease test is indicated by both isolates. Both Isolate (A1 and A2) showed positive result on the test methyl red, whereas Voges Proskauer test of the both isolates showed negative results as did the catalase test, all isolates could not catalyze H<sub>2</sub>O<sub>2</sub>. The difference between the observations results of the characteristics both isolates lies in the carbohydrate lactose and maltose test. The difference between the results of observations of the characteristics of both isolates lies in the carbohydrate lactose and maltose test. In addition, both cell lines also showed differences, isolates A1 rod-shaped isolates shorter than isolates A2. Both isolates showed characteristics of lactic acid bacteria that is strongly suspected as the genus *Lactobacillus*.

Keywords: Lactic Acid Bacteria, whey, Dangke, *Lactobacillus*

### INTRODUCTION

Lactic acid bacteria (LAB) are a group of bacteria that have the ability of produce lactic acid, either as the sole product nor as the primary products in the metabolism of carbohydrates. BAL widely used and researched as probiotics for humans, animals on the mainland and aquatic animals. It is based on the fact that BAL is the normal flora in the digestive tract with ability of tolerance to acidic conditions and bile salt (bile) in the digestive tract and be able to convert lactose to acetic acid, thus lowering the pH of the digestive tract and naturally prevent colonization of pathogenic bacteria (Watson *et al.*, 2008). Capabilities of BAL in maintaining digestive balance is also supported by its ability to produce antimicrobial substances such as organic acids, diacetyl, hydrogen peroxide, and bacteriocins that can suppress the growth of pathogenic bacteria. In addition, lactic acid bacteria have the ability of to stick to intestinal epithelial cells, as well to an increase in the cell as an indication secreted peyer plates immunoglobulin (IgA), ie an reaction of the formation immune to bacterial infections.

Hafsan, M.H. Mustami

Lactic acid bacteria shown to suppress the growth of pathogenic and spoilage bacteria so it can maintain the natural balance of microflora in the intestine. Lactic acid bacteria are generally present in the intestines of humans and animals is helpful in countering various diseases that usual entered through the digestive tract. It has been reported also by Trisna *et al.* (2012) that probiotics can improve the health of livestock, increase egg production, as well can eliminate the reservoir properties of AI (avian influenza) in poultry. Also been conducted isolation of lactic acid bacteria from chicken intestine performed by Mutmainnah (2013), where there are several isolates and qualify as a probiotic characteristics. Superior properties owned by BAL is very widely used in the processing and preservation of food (livestock, marine products, vegetables and fruits). Lactic acid bacteria commonly found in processed food products, either animal products such as meat and fermented fish, fermented milk, nor the fermented vegetable products such as vegetables and fruits (Suharsono, 2010).

Availability of lactic acid bacteria from various sources of domestic was still requires exploration of lactic acid bacteria to improve the collection of lactic acid bacteria isolates that have potential as probiotics. Lactic acid bacteria can be obtained by utilizing the resources that contain lactic acid bacteria. Dangke as one of the distinctive dairy products in South Sulawesi, the process leaves waste in the form of whey. Nutrient content of the whey which was still high can be utilized by bacteria as a nutritional source of growth. Whey contains about 55% of total nutrients from milk such as lactose, proteins, fats, vitamins, and mineral salts. Such conditions it is possible for BAL to make it as a medium to grow and evolve (Pato *et al.*, 2004).

This research was aimed to investigate the characteristics of lactic acid bacteria isolated from Dangke waste production, known as whey. So it can increase the repertoire of knowledge and information about the diversity of lactic acid bacteria that can have a positive impact in the health sector, food and beverages nor livestock.

## MATERIAL AND METHODS

Whey obtained from home industry of Dangke in Enrekang City, Enrekang District South Sulawesi. Isolation and characterization of lactic acid bacteria has been conducted in the Laboratory of Microbiology, Biology Department, Faculty of Science and Technology UIN Alauddin Makassar, take place in January until March 2014. Isolation and characterization of lactic acid bacteria has been conducted in the Laboratory of Microbiology, Department of Biology, Faculty of Science and Technology UIN Alauddin Makassar, take place in January until March 2014.

Whey sampled aseptically at home Dangke producer, shortly after their production process and immediately taken to the laboratory and soon be isolated using selective media MRS. Isolation of lactic acid bacteria conducted by inoculating suspension whey that has been previously sampled by 10 ml in 90 ml of medium de Man, Rogosa and Sharpe (MRS), incubated for 1 x 24 hours at 37 ° C using a shaker incubator. Culture from MRS medium and then inoculated on MRSA were added indicator Bromcresol purple (BCP) 0.1%, incubated for 2 x 24 hours. Growing colony forming clear zone surrounding purified with streak on MRSA medium and incubated for 2 x 24 hours. Cultivation is done repeatedly on the same medium and conditions to obtain single colonies. Single colonies obtained on agar slant was transferred as a pure isolate stock and stored in the refrigerator at 4° C

Characterization of lactic acid bacteria include observation of colony morphology and cell observation as well physiological properties through biochemical tests. Morphological observation includes observation of colony morphology and Gram's staining, whereas observations with biochemical test consists of motility test, indole test, citrate test, methyl red test, Voges Proskauer test, carbohydrate fermentation test, test lysine, and catalase test.

Hafsan, M.H. Mustami

## RESULT

Isolation conducted by firstly culturing the whey suspension on MRS medium for 1 x 24 hours at 37oC. MRS liquid culture from inoculated by pour plate method on MRSA medium BCP + 0 1% for 1 x 24 hours at 37oC. Medium purple color turns into yellow color indicates a growing isolates of lactic acid bacteria. Observation of colony morphology and biochemical testing of the bacterial cell as well as Table 1 is the physical properties and physiological lactic acid bacteria from waste of Dangke.

**Table 1. Characteristics of Lactic Acid Bacteria From Dangke Waste Production**

Sifat	LAB isolates code	
	A1	A2
Koloni	Round shape, flat edge, convex, white milk	Small round shape, flat edge, convex, white milk
Gram	Positive	Positive
Cell shape	Basil pendek	Basil panjang
Spore	-	-
KIA	A/A, -/-	A/A, -/-
MIO	-	-
Citrat	-	-
Urea	-	-
D-Glucosa	+	+
Galactosa	+	+
Lactosa	+	-
Sucrosa	+	+
Maltosa	+	-
L-Arabinosa	+	+
D-Silosa	-	-
Rhaminosa	+	-
Sorbitol	-	+
Salisin	+	+
Raffinosa	+	-
Trehalosa	+	+
Manitol	-	-
Malonat	-	-
LIA	-	-
MR	+	+
VP	-	-
Catalase	-	-
<b>Spesies</b>	<b><i>Lactobacillus acidophilus</i></b>	<b><i>Lactobacillus plantarum</i></b>

## DISCUSSION

Bromcresol purple is an color indicator of a decrease in the pH of the medium. MRSA medium discoloration added BCP, from purple to yellow as a result of acid produced by LAB isolates from the fermentation of glucose thus lowering the pH of the media (Lay, 1994). Fermentation can occur because of the activity of microbes that cause fermentation in suitable organic substrates. Fermentation as a gradual change in the enzyme from several bacteria, molds, and yeasts in the growth medium. Bacterial growth generally decreased, due to changes in fermentation environmental conditions be acidic, resulting in bacteria are not tolerant of the situation. According Pato *et al.* (2004) that, the bactericidal effect from lactic acid related to the decrease in the pH of the environment to be 3 to 4.5 so that the growth of other bacteria, including spoilage bacteria will be inhibited.



Hafsan, M.H. Mustami

Identification of lactic acid bacteria genera requires the main characters from ie bacterial colony morphology, cell morphology (cell shape and arrangement of cells) and biochemical tests. Observation of morphological and biochemical tests can be performed to identify the level of genera of lactic acid bacteria. Morphology test is performed to determine the morphology from the bacteria to be identified whereas biochemical test used to determine the ability of bacteria to use chemical substances for its growth.

The morphology of bacterial colonies that grow in the bacterial growth medium ie MRSA visible form a round (circular), flat edge (entire), convex surface and milky white. This same characteristic colony morphology between isolates A and B. According Mutmainnah isolates (2013) colony morphology that is round (circular), flat edge (entire), convex surface and a creamy white morphological features from lactic acid bacteria of the genus *Lactobacillus*. *Lactobacilli* are widespread in the environment, particularly in the dairy and food products vegetables, normally inhabit the intestinal tract of birds and mammals, and mammals vaginal as well non-pathogens.

On the results of Gram staining, both isolates ie isolates A and B isolates showed rod-shapes. This is consistent with the statement from Wibowo and Ristanto (1988), ie bacterial cells from the genus *Lactobacillus* are rod-shaped \_ varied from very short to long rod, are homofermentatif or heterofermentatif. Both isolates were also included in the Gram-positive. Gram staining performed on bacterial culture was 1 x 24 hour MRSA grown on medium. Lactic acid bacteria are Gram-positive bacteria. Gram-positive bacteria will show violet color when colored Gram. The violet color occurs because the cell walls of bacteria bind crystal violet (Gram A) were amplified by iodine (Gram B) and crystal violet will not be lost on a given laxative (Gram C), so it is not affected when the colored cover (Gram D ) are colored red. Gram's staining From the the results can also be used to see the shape and arrangement of cells of lactic acid bacteria. Observations were made at a magnification of 1000 x. The results of microscopic examination showed rod shape or bacillus whereas cell arrangement and clustered mostly chain. The results of this test in accordance with test results that had been conducted by Mutmainnah (2013), ie has the form of rods and Gram-positive trait is characteristic from genus *Lactobacillus*.

The principle of gram staining is ability of the cell wall binding basic dye (crystal violet) after washing with 95% alcohol. This situation is related to the structure of the cell wall constituent compounds. Gram-positive bacteria contain more peptidoglycan and less fat than gram-negative bacteria (Syulasmı *et al.*, 2005). *Lactobacilli* have long shaped cells, the cylinder rod (sometimes curved) while others are short rod-shaped, frequently Coryne-shaped or round rod. Cell also often form chains. Although belonging to the gram-positive bacteria, but there is also a bipolar (Gram positive and Gram negative) may occur due to granulation in the cells. Metabolism is fermentative metabolism by converting at least 50% of carbon into lactic acid.

Gram-negative bacterial cell wall consists from peptidoglycan 5-20%, the rest is a polysaccharide, whereas Gram-positive bacterial cell wall peptidoglycan contains the remaining 90% is teikoat acid. Gram-positive bacterial cell visible purple, because it can form a complex bond with crystal violet ie complex crystal violet-iodine. Giving 95% alcohol solution to Gram-negative bacteria cell can increase the porosity of the cell wall by dissolving the lipids in the outer membrane, so that the crystal violet iodine complex will be separated and the cell be colorless. Furthermore, the cell will be colored red because it is colored by dye comparator ie safranin (Madigan *et al.*, 2011).

Hafsan, M.H. Mustami

Motility test were conducted on both isolates by means cultivated on upright agar media by way of put a inoculation needle that has a bacterial inoculation in the media. It is observed in this test that is the propagations of the puncture marks on the media to know the ability of the bacteria to move. The test results showed that both isolates did not do the movement in the media so it is said to be non-motile bacteria. According Surono (2004), probiotic bacteria have a very limited ability for biosynthesis, so that is non motile. Acquisition energy solely rely on the fermentative metabolism were conducted in place. Negative results on motility test showed that the bacteria do not have flagella as test movements appliance. Reported by Mutmainnah (2013) and Suryani *et al.* (2010), to obtain a negative result on the test the motility to lactic acid bacteria is their isolation.

Indole test was conducted to determine ability of bacteria to form indole from tryptophan. In this test added with Erlich reagent or Kovach which can lead to medium red. The results showed negative for both isolates. This means that the bacteria are not able to produce an triptofanase enzyme that can catalyze reactions solving tryptophan be indole, pyruvic acid and ammonia. The same thing also happened on research conducted by Paramitasari (2013), ie lactic acid bacteria do not produce indole. It is also consistent with the statement Soeharsono (2010) that the lactic acid bacteria do not produce indole.

Methyl red test was used to determine the presence of a mixed acid products from fermentation of glucose by mixed acid fermentation pathways are generally in the form of lactic acid, acetic acid, formic acid, and succinic acid. Based on the research results, both isolates showed positive results on the MR test is characterized by the formation of red in the medium after adding a solution of methyl red as indicator. Brown (2001) explains that the red color formed on the media by the media due to decrease in the pH of media by acidic products in large quantities are produced from glucose fermentation. The results of this test in accordance with Surono statement (2004) that the fermentation process involving lactic acid bacteria has a characteristic ie the accumulation of organic acids is accompanied by a decrease in pH.

Voges Proskauer test was used to determine ability of isolates produce a final product that is neutral (asetilmetilkarbinol) from fermentation of glucose through butanediol pathway (Brown, 2001). In Voges Proskauer test, Barritt's A ( $\alpha$ -naphthol) and Barritt's B (potassium hydroxide) reagent is added. Tests showed negative results against both isolates. This can be seen with, there are no red ring formed at the media after the reagent drops.

Citrate test (using Simmons citrate medium) is intended to know the ability of of bacteria to use citrate as the sole carbon source. test showed negative results against both isolates. This can be seen with no discoloration of the medium from green to blue. In this test, the bacteria can grow in order to increase ie pH from acid to alkaline, whereas lactic acid bacteria are bacteria that produce lactic acid. Lactic acid bacteria use carbohydrates as its main source of nutrients in the fermentation.

Urease test against both bacterial isolates showed negative results, which is indicated by the formation of reddish-purple in the culture medium urease test. This shows that both the bacterial isolates could not produce urease enzyme that can break down urea to CO<sub>2</sub> and NH<sub>3</sub>, so phenol red indicator remains yellow, which means not formed alkaline compounds such as NH<sub>3</sub> in the culture medium.

Lactic acid bacteria have special properties that are able to ferment carbohydrates types of monosaccharides and disaccharides. Carbohydrate test done using some type of sugar in a liquid medium is glucose, lactose, sucrose, mannitol, maltose and malonate. A isolates showed positive results on glucose, sucrose, lactose and maltose, while a negative result on the test mannitol and malonate. At A2 isolates showed positive results on tests of glucose and sucrose,

Hafsan, M.H. Mustami

while a negative result on the test maltose, mannitol and malonate. Different results in various types of fermentation on sugar so it can be seen that both the lactic acid bacteria of different species. According Helferich and Westhoff (1980), carbohydrate metabolism of lactose is hydrolyzed by the enzyme lactic acid bacteria with  $\beta$ -galactosidase into glucose and galactose. Glucose can be directly used to produce lactic acid, while galactose will accumulate. Sucrose is converted into one molecule of glucose and one molecule of fructose while lactose is converted into 2 molecules glukosa. fermentation test of isolates B, shows not able to ferment the lactose medium, mannitol, maltose and malonate. Statement Soeharsono (2010), *Lactobacillus acidophilus* species are not capable of performing fermentation using sugars mannitol, lactose, maltose and malonate in fermentation.

Testing the ability of fermentation also showed that the tube does not form gas bubbles which means that the mixture acid not formed in the fermentation. Similar things was reported by Lay (1994), so it can be affirmed that both isolates of lactic acid bacteria isolates A1 and A2 are lactic acid bacteria that are homofermentatif.

Catalase test is a test to identify microbes that are capable of producing the catalase enzyme is used to break down the hydrogen peroxide formed from the process of aerobic respiration and are toxic to bacteria, into dihydrogen oxide (H<sub>2</sub>O) and oxygen (O<sub>2</sub>) that is not toxic anymore. Results showed these bacteria are not able to produce the enzyme catalase thus giving a negative reaction. These results are consistent with research conducted by Sri *et al.*, (2009) who also obtained a negative result on catalase test is performed.

Based on morphological observations and chemical tests that have been done then both allegedly isolates belonging to the genus *Lactobacillus* to satisfy the criteria have the nature of catalase negative, Gram-positive, nonmotil, and rod-shaped cell. Selection is based on Buchanan and Gibbons (1974) in Bergey's manual of determinative Bacteriology stating that the nature of catalase has *Lactobacillus* negative and Gram-positive and rod-shaped cell. more specifically, that A1 isolates have characteristics resembling *Lactobacillus acidophilus* and A2 isolates have characteristics resembling *Lactobacillus plantarum*, but characterization has been done still requires confirmation by molecular level, so as can be ascertained species have been isolated.

Bacteria belonging to the genus *Lactobacillus* has a rod-shaped cells varied from very short rods to long rods, are homofermentatif or heterofermentatif (Wibowo and Ristanto, 1988). This bacterial genus also are microaerophilic, catalase negative, Gram-positive and ferment sugars to lactic acid as the primary products. When is homofermentatif will ferment sugars into lactic acid, whereas when is heterofermentatif will produce volatile products including alcohol other than lactic acid. Homofermentatif *Lactobacillus* grown with optimum temperature 37° C or lower, for example, *L. bulgaricus*, *L. helveticus*, *L. lactis*, *L. acidophilus* and *L. Thermophilus*. *L. Delbrueckii* and *L. fermentum* is heterofermentatif *Lactobacillus* which is able to grow at high temperatures (Frazier, 1988). Bacteria of this genus are found in plants, vegetables, whole grains, fresh milk and dairy and meat and fermented meat products, fermented vegetables and some of the species found in the digestive tract of humans and animals (Ray, 2004).

## ACKNOWLEDGEMENT

The authors would like thank to the entire academic of Biological department Faculty of Science and Technology UIN Alauddin Makassar for the support and facilities that we use, So too the same thing to LP2M UINAM. This research is supported by a grant from UIN Alauddin Makassar DIPA BLU grant 2013.



Hafsan, M.H. Mustami

## REFERENCE

- Abun. 2008. *Hubungan Mikroflora dengan Metabolisme dalam Saluran Pencernaan Unggas dan Monogastrik*. Universitas Padjajaran Press. Jatinangor.
- Asriani. 2006. *Kajian Efek Sinergi Antimikroba Metabolit Bakteri Asam Laktat dan Monoasilgliserol Minyak Kelapa terhadap Bakteri Patogen [Tesis]*. Bogor: Pertanian Bogor Institute.
- Brown, A. 2001. *Microbiological Applications Lab Manual. 8th Ed.* The McGraw-Hill Companies. New York.
- Buchanan, R.E., and Gibbons, N.E. 1974. “*Bergey’s Manual of Determinative Bacteriology 8<sup>th</sup> edition*”. The William & Wilkin Company/Baltimore. USA.
- Frazier William C, Dennis C. Westhoff. 1988. *Food Microbiology*. McGraw-Hill. U.S.A.
- Helferich, William C, Dennis C. Westhoff. 1980. *All About Yogurt*. Prentice Hall. U.S.A.
- Hafsan. 2011. *Mikrobiologi Umum*. Alauddin Press. Makassar
- Holt. G., Kreig, N.R., Sneath, P.H.A., Stanley, J.T. & Williams, S.T. 2000. *Bergey’s Manual Determinative Bacteriology*. Baltimore: Williamn and Wilkins Baltimore.
- Jawetz, Melnick, & Adelberg’s. 2001. *Medical Microbiology*. Salemba Medica. Jakarta.
- Lay, B. W. 1994. *Analisis mikrobiologi di Laboratorium*. PT Raja Grafindo Persada. Jakarta.
- Madigan MT, Martinko JM, Stahl DA, and Clark D. 2011. *Biology of Microorganisms Thirteenth Edition. Pearson Education International. USA.*
- Mutmainnah, Heni. 2013. *Isolasi Dan Karakterisasi Bakteri Probiotik Dari Saluran Pencernaan Ayam Kampung Gallus domesticus [Skripsi]*. Makassar: Universitas Hasanuddin.
- Paramitasari, Dewi. 2013. *Isolasi dan Karakterisasi Bakteri Asam Laktat dari Dangke [Skripsi]*. Gowa: Alauddin Islamic State University Makassar.
- Pato U, Surono IS, Koesnandar A, Hosono A. 2004. *Hypocholesterolemic effect of indigenous dadih lactic acid bacteria by deconjugation of bile salts*. Asian-Australian Journal of Animal Sciences 17:1741-5.
- Purwoko T, 2009. *Fisiologi Mikroba*. Jakarta: Bumi Aksara.
- Ray B, 2004. *Fundamental Food Microbiology*. CRC Press. New York.
- Setyaningsih, I. 1992. *Pengaruh Jenis Kultur L. Casei, Penambahan Susu Skim dan Glukosa terhadap Mutu Yakult Kedelai [Skripsi]*. Bogor: Pertanian Bogor Institute.
- Soeharsono. 2010. *Probiotik. Basis Ilmiah, Aplikasi dan Aspek Praktis*. Bandung: Widya Padjajaran.

*Hafsan, M.H. Mustami*

- Sri Sumarsih, T. Yudiarti, C.S.Utamar, E. S. Rahayu, dan E. Harmayani. 2009. *Isolasi dan identifikasi bakteri asam laktat pada caecum ayam Pedaging*. Semarang: Universitas Diponegoro
- Surono, I.S, 2004. *Probiotik Susu Fermentasi dan Kesehatan*. Tri Cipta Karya. Jakarta.
- Suryani Y, Astuti, Bernadeta Oktavia, Siti Umniyati. 2010. *Isolasi Dan Karakterisasi Bakteri Asam Laktat Dari Limbah Kotoran Ayam Sebagai Agensi Probiotik Dan Enzim Kolesterol Reduktase [Skripsi]*. Yogyakarta: Yogyakarta State University.
- Trisna dan Wahud N. 2012. *Identifikasi Molekuler dan Pengaruh Pemberian Probiotik Bakteri Asam Laktat (BAL) Asal Dadih dari Kabupaten Sijunjung Terhadap Kadar Kolestrol Daging pada Itik Pitalah Sumber Daya Genetic Sumatra Barat*. Artikel. Padang: Andalas University.
- Watson A.K., H. Kaspar, M.J. Lategan & L. Gibson. 2008. *Probiotic in aquaculture: The need, principles and mechanisms of action and screening processes*. *Aquaculture* 274: 1-14.
- Wibowo dan Ristanto, D. 1988. *Petunjuk Khusus Deteksi Mikroba Pangan*. Yogyakarta: Pusat Antar Universitas Pangan dan Gizi Universitas Gadjah Mada.
- Widiasih T. 2008. *Aktivitas Substrat Antimikroba Bakteri Asam Laktat Yang Diisolasi Dari Daging Sapi Terhadap Bakteri Patogen dan Konsentrasi Minimum Penghambatannya [Skripsi]*. Bogor: Pertanian Bogor University.