

Phytate Activity of Thermophilic Bacteria from Sulili Hot Springs in Pinrang District South Sulawesi

by Hafsan .

Submission date: 15-Jul-2019 12:01PM (UTC+0700)

Submission ID: 1151965822

File name: scriptaBiologica.pdf (321.09K)

Word count: 2460

Character count: 12855

PHYTATE ACTIVITY OF THERMOPHILIC BACTERIA FROM SULILI HOT SPRINGS IN PINRANG DISTRICT SOUTH SULAWESI

HAFSAN¹, NURJANNAH¹, CUT MUTHIADIN¹, ISNA RASDIANA¹, AHYAR AHMAD², LAILY AGUSTINA, ASMUDDIN NASIR³

¹Fakultas Sains dan Teknologi, UIN Alauddin Makassar, Jalan H.M. Yasin Limpo No. 36 Gowa, Sulawesi Selatan

²Faculty of Mathematics and Natural Sciences, Universitas Hasanuddin, South Sulawesi

³Faculty of Animal Science, Universitas Hasanuddin, South Sulawesi

ABSTRACT

This study aimed to determine the optimum temperature and pH of the phytase activity produced by isolate of thermophilic bacteria from Sulili hot springs in the district Pinrang south Sulawesi. This is a descriptive research, that measure of phytase activity by treated the variation of temperature and pH. The various of temperature has eight levels ie 20; 30; 40; 50; 60; 70; 80; and 90°C, the pH treatments consists of eight levels ie 2; 3; 4; 5; 6; 7; 8 and 9. Phytase activity was determined by converting the Na-phytate absorbance values produced by phytase enzyme against Na-phytate standard curve. One unit of phytase enzyme activity is defined as the amount of enzyme capable of hydrolysed Na-phytate and produce a peptide which is equivalent to one mole of tyrosine per minute under the conditions of analysis. The results of studies have shown that there is a significant effect of temperature on the phytase activity of *Bacillus coagulans*, thermophilic bacteria from Sulili hot springs, whereas no significant effect of pH treatment. Optimum phytase activity of *B. coagulans* are at 60°C and pH 7 ie 6.11 units/ mL.

KEY WORDS: *Bacillus coagulans*, phytase, Sulili hot springs, temperatures and pH, thermophilic bacteria

Corresponding author: HAFSAN | email: hafsan.bio@uin-alauddin.ac.id

INTRODUCTION

Phytase is one of the most popular enzymes, especially in poultry farms field. This is related to the application of these enzymes as an effort to improve the quality of poultry feed (Kusumadaja *et al.*, 2009). Some studies reported that phytase supplementation in feed, can improve the digestibility of nutrients that leads to improve the performance of poultry (Hafsan *et al.*, 2017). Increased nutrient digestibility is the result of the release of protein and essential minerals such as phosphorus and calcium from the myo-inositol ring contained in the phytate compound feed (Blaabjerg *et al.*, 2015). Therefore, it can be used maximally by poultry for metabolic requirements (Bhavsar *et al.*, 2012).

As a feed supplementation for industrial scale production, the most appropriately used is a thermostable phytase. It can produce a much better and more efficient phytase, because the utilization process can reduce the possibility of microorganisms contamination as it is done at high temperatures, otherwise it can save cooling costs in large-scale fermentation so processing by using high temperature is an option in the industry.

Microorganisms are the most widely used as a source of enzymes. This gives more benefit because their cell are relatively easy to culture, the growth rate is relatively faster, easy to increase cell production as desired, production does not depend on season change, time required in shorter production process, and uniform quality (Pawiroharsono, 2003; Hasyimuddin *et al.*, 2016). Superior microorganisms are an

important factor in enzyme production efforts. Therefore, it is necessary to explore local microorganisms that produce phytase in Indonesia. High biodiversity provides a great opportunity to get potential microorganisms to produce enzymes.

The wealth of Indonesia's natural resources is abundant. One of them is the source of hot springs that can be used not only as a tourist destination, but also the sources of thermophilic bacteria. A country with a rich source of hot springs will have high potential as a ride for thermophilic bacteria growth, but until now the exploration of thermoplastic hot spring microorganisms from potential sites is still uneven, especially in South Sulawesi. It is still not widely applied. In this research, we have measured the phytase activity of indigenous bacteria from hot springs on temperature and pH variations so that the optimum activity and its stability in temperature and pH variations are known. The results of this study are expected to be a reference in the effort of optimum phytase production.

METHODS

Thermophilic bacteria isolates from Sulili hot springs (*B. coagulans*) has rejuvenated with inoculated the pure isolates in slant medium and incubated for 1 x 24 hours, then stored at 4°C. Preparation of standard curve of growth was determined by measured the value of Optical Density (OD) fermentation media. Bacteria were inoculated in 50 ml of medium as much as 2 ose isolates. Bacterial culture that has been suspended incubated in an incubator shaker at 60°C with agitation of 200 rpm. OD values were measured every 2 hours with spectronic 20 on $\lambda = 280$ nm to obtain a series of OD using turbidimetric method. Standard curve of growth is the relationship between the numbers of cells per millilitre

with OD.

Production and isolation of phytase enzyme; 27 ses bacterial culture inoculated from agar slant into 50 ml of liquid medium and then shake at 60°C with agitation of 200 rpm until the culture reached the logarithmic phase and stored as a starter at the time of phytase production. Then about 5 ml starter culture medium was inoculated into the new liquid medium and then shaken at 60°C for 24 hours with an incubator shaker at 200 rpm. Incubed bacterial cell cultures were separated from the medium by centrifugation at 6000 g for 20 min at 4°C. The supernatant obtained a crude extract of enzyme activity were then tested (El-Toukhy *et al.*, 2011).

Measurement of phytase activity; determined based on the ability of the crude extract phytase to hydrolyse the myo-inositol bond of the phytate as substrate. The phytase activity is known by measuring the absorbance value of the crude extract of phytase at $\lambda = 700$ nm which then analyzed the amount of phosphate content formed (Unit/mL) in solution of crude extract of enzyme by substitution of absorbance value using linear regression equation of standard phosphate curve (El-Toukhy *et al.*, 2013). As for the treatment of variations in temperature and pH measurement of phytase activity carried out at a temperature variation (20, 30, 40, 50, 60, 70, 80; and 90°C) and pH (2; 3; 4; 5; 6; 7; 8; and 9). Absorbance value obtained is substituted into the formula determining the phytate activity. One phytase unit (U/mL) is defined as the number of enzymes catalyzing the formation of 1 μ mol phosphate per minute.

RESULT AND DISCUSSION

Bacterial Growth Curve. Phytase production has begun by determining the bacterial growth curve. It is important to know the optimum phase of phytase production. The growth curve of *B. coagulans* as in Figure 1. The growth curve is the relationship between the values of optical density (OD) to the incubation time. The OD value was measured every 2 hours with a spectrophotometer at λ 280 nm and obtained a series of OD using a turbidimetric method.




Figure 1. Growth curve of *Bacillus coagulans*

Generally enzymes are produced during bacterial growth and will achieve the highest activity at the end of the exponential phase or logarithmic phase. The growth curve showed the *B. coagulans* log phase achieved at 12 hours. The time has the basis for the length of incubation required as a starter, so the length of incubation required for the production of phytase is up to the end of the log phase which can be extended to half the stationary phase. At the end of the log phase

there is a peak increase in the number of cells as each active cell multiplies, whereas in the stationary phase there is no longer a doubling of bacterial cells.

Based on the growth curve *B. coagulans*, phytase production was performed by inoculated a 5 mL starter isolate culture suspension into 250 mL of sterile PPM medium and incubated at 60°C with an incubation time of 18 h using an incubator shaker at 100 rpm. The culture was then centrifuged 5000 rpm for 35 min at 4°C. The supernatant obtained by separation from its precipitates is crude phytase, then its activity is measured.

The phytase activity on variations of temperature and pH. The results of measurement of *B. coagulans* phytase activity on temperature variation and pH are shown in Figures 2 and 3. As shown in figure 2, that increase of temperature causes an increase in activity to reach the optimum point at 60°C. Increasing the temperature further showed decreased activity of phytase, which began to look at 70°C.




Figure 2. Phytate activity of *Bacillus coagulans* on the temperatures variation.

At first with the increase in temperature there is an increase in the speed of the enzyme reaction because of the increased kinetic energy that accelerates the vibrational motion, translation and rotation of the enzyme and substrate thereby increasing the chances of both to react. At exposure to higher temperatures than optimum temperature, the protein will undergo conformational changes. Similarly, the substrate may undergo a conformational change so that the reactive group does not correspond to the active site, or experience resistance to entering the enzyme's active site so as to significantly affect its catalysis activity. At the optimum temperature, the enzyme activity is optimal and optimal product produced also (Vielle and Zeikus, 2001).

The activity of *B. coagulans* phytase was seen from the pH 2 treatment and showed an increase starting at pH 3 to reach the optimum at pH 4 and at pH 5 to pH 9 become decreased. The difference in activity at different pH results from the occurrence of intramolecular changes from enzymes caused by ionization binding and releasing protons (hydrogen ions) in amino, carboxyl and other functional groups. If the change is too large it can result in the denaturation of the enzyme, so that its activity will be lost (Madigan

et al., 2004).




Figure 3. Phytase activity of *Bacillus coagulans* on the pH variation

The results of this measurement are in line with the properties of phytase which is a heterologous group of enzymes, a subclass of phosphatases, having the ability to hydrolyse phosphate esters and optimum at low pH. At neutral pH and bases, catalytic activity decreases. This is due to the instability of structures in the enzyme protein molecules thus causing structural changes under these pH conditions (Vats and Banerjee, 2004; Tomschy et al., 2002)

The optimum temperature and pH of the enzyme depend on the type and source of the enzyme. Based on the measurement of phytase activity on temperature and pH variations that have been done, it is known that the temperature and pH of optimum phytase produced by *B. coagulans* is 60°C and pH 7. Variation of optimum pH and temperature of phytase from various microbes have been reported. Choi et al. (2012) obtained phytase from *Bacillus* sp. KHU-10 with optimum pH and temperature for phytase activity were pH 6.5-8.5 and 40°C without 10 mM CaCl₂ and pH 6.0-9.5 and 60°C with 10 mM CaCl₂. Phytase is also produced by *Lactobacillus sanfranciscensis* (pH 4.0 and 45 °C), *Mitsuokella jalaludinii* (pH 7.0 and 39 °C), *Pedobacter nyackensis* (pH 7.0 and 45 °C), and *Aspergillus niger* (pH 5.0 and 50 °C) (Mandviwala and Khire, 2000; Lan et al., 2002; Angelis et al., 2003; Huang et al., 2009).

CONCLUSION

The results of studies have shown that the activity of phytase from *Bacillus coagulans*, thermophilic bacteria from Sulili hot springs, optimum at 60°C and pH 7 ie 6.11 units/ml. The phytase activity of *B. coagulans* at the optimum temperature and pH of treatment indicates the enzyme probability to be produced on an industrial scale and highly applicable to be used as feed additives in poultry feed, as it has a high temperature resistance and relatively stable at pH variations.

REFERENCES

Angelis MD, Gallo G, Corbo MR, McSweeney PLH, Faccia M, Giovine M, Gobetti M. 2003. Phytase activity in sourdough lactic acid bacteria: purification and characterization of a phytase from *Lactobacillus sanfranciscensis* CB1. International Journal of Food

Microbiology 87(3):259-270. doi: 10.1016/S0168-1605(03)00072-2.

Bhavsar K, Kumarb VR, Khirea JM. 2012. Downstream Processing of Extracellular Phytase from *Aspergillus niger*: Chromatography Process vs. Aqueous Two Phase Extraction for its Simultaneous Partitioning and Purification. Process Biochem. 7(47):1066-1072

Blaabjerg K, Thomassen AM, Poulsen HD. 2015. Microbial phytase addition resulted in a greater increase in phosphorus digestibility in dry-fed compared with liquid-fed non-heat-treated wheat-barley-maize diets for pigs. Anim. 9(2):243-248.

Choi YM, Suh HJ, Kim MJ. 2001. Purification and Properties of Extracellular Phytase from *Bacillus* sp. KHU-10. Protein Chem 20: 287. doi: 10.1023/A:1010945416862.

El-Toukhy NMK, Youssef AS, Mikhail MGM. 2013. Isolation, purification and characterization of phytase from *Bacillus subtilis*. African J. Biotechnol. 12:2957-2967.

Hafsan, Irwan I, Agustina L, Ahmad A, Natsir A. 2017. Isolation and characterization of phytase-producing thermophilic bacteria from sulili hot springs in South Sulawesi. SCIRJ. 5(12):16-19.

Hasyimuddin, Djide MN, Samawi MF. 2016. Isolasi Bakteri Pendegradasi Minyak Solar Dari Perairan Teluk Pare-Pare. Biogenesis 4(1):41-46. doi: 10.24252/bio.v4i1.1119.

Huang H, Shao N, Wang Y, Luo H, Yang P, Zhou Z, Zhan Z, Yao B. 2009. A novel beta-propeller phytase from *Pedobacter nyackensis* MJ11 CGMCC 2503 with potential as an aquatic feed additive. Appl Microbiol Biotechnol 83: 249. doi: 10.1007/s00253-008-1835-1.

Kusumadjaja S, Piliang WG, Ketaren PP. 2009. Suplementasi enzim pemecah serat dan phytase terhadap performans. JITV. 1(14):40-46.

Lan GQ, Abdullah N, Jalaludin S, Ho YW. 2002. Culture conditions influencing phytase production of *Mitsuokella jalaludinii*, a new bacterial species from the rumen of cattle. Journal of Applied Microbiology 93(4):668-674. doi: 10.1046/j.1365-2672.2002.01727.x.

Madigan MT, Martinko J, Parker J. 2004. Brock Biology of Microorganisms 10th Edition. NJ: Pearson Prentice Hall.

Mandviwala TN and Khire JM. 2000. Production of high activity thermostable phytase from thermotolerant *Aspergillus niger* in solid state fermentation. J Ind Microbiol Biotech 24: 237. doi: 10.1038/sj.jim.2900811.

Pawiroharsono S. 2003. Microbial enzyme and their application in industry. Proceeding of Seminar Industri Enzim dan Bioteknologi. Malang: CIEB.

Tomschy A, Brugger R, Lehmann M, Svendsen A, Vogel K, Kostrewa D, Lassen SF, Burger D, Kronenberger A, van Loon APGM, Pasamontes L, Wyss M. 2002. Engineering of Phytase for Improved Activity at

- Low pH. *Appl. Environ. Microbiol* 68(4):1907-1913.
doi: 10.1128/AEM.68.4.1907-1913.2002.
- Vats P and Banerjee UC. 2004. Production studies and catalytic properties of phytases (myo-inositolhexakisphosphate phosphohydrolases): an overview. *Enzyme and Microbial Technology* 35(1):3-14. doi: 10.1016/j.enzmictec.2004.03.010
- Vielle C and Zeikus GJ. 2001. Hyperthermophilic enzymes: sources, uses, and molecular mechanism, for thermostability. *Microbiol Mol Biol Rev.* 7(65):14-18.

Phytate Activity of Thermophilic Bacteria from Sulili Hot Springs in Pinrang District South Sulawesi

ORIGINALITY REPORT

22%

SIMILARITY INDEX

19%

INTERNET SOURCES

9%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

iicbe.org

Internet Source

15%

2

link.springer.com

Internet Source

2%

3

academicjournals.org

Internet Source

1%

4

Hasan B. Coban, Ali Demirci. "Phytase as a Diet Ingredient: From Microbial Production to Its Applications in Food and Feed Industry", Elsevier BV, 2017

Publication

1%

5

"Recent Advancement in White Biotechnology Through Fungi", Springer Science and Business Media LLC, 2019

Publication

1%

6

proteomesci.biomedcentral.com

Internet Source

1%

7

Karl-Heinz Schleifer. "Phylum XIII. Firmicutes

Gibbons and Murray 1978, 5 (Firmacutes [sic]
Gibbons and Murray 1978, 5)", Systematic
Bacteriology, 2009

Publication

1%

8

A. BOYCE. "Comparison of Selected
Physicochemical Characteristics of Commercial
Phytases Relevant to their Application in
Phosphate Pollution Abatement", Journal of
Environmental Science and Health Part A
Toxic/Hazardous Substances & Environmental
Engineering, 6/1/2006

Publication

1%

9

repositori.uin-alauddin.ac.id

Internet Source

1%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On