

# Test of Hepatoprotector Effect of Merak Leaf Ethanol Extract (Caesalpinia Pulcherrima L) with SGPT Enzyme Parameter and SGOT of Induced Parasetamol Heart Rats (Rattus Norvegicus)

*by* Gemy Nastity Handayany

---

**Submission date:** 20-Jun-2020 10:18PM (UTC+0700)

**Submission ID:** 1347039398

**File name:** GEMY\_3.pdf\_JURNAL\_SCOPUS\_Q3.pdf (8.94M)

**Word count:** 2950

**Character count:** 15326

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

Volume 10 / Number 1 / January 2019



# Indian Journal of Public Health Research & Development

An International Journal

## SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development

Scopus coverage years: from 2010 to 2018 Publisher:

R.K. Sharma, Institute of Medico-Legal Publications

ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:

Public Health, Environmental and Occupational Health

Cite Score 2017- 0.03

SJR 2017- 0.108

SNIP 2017- 0.047



Website:

[www.ijphrd.com](http://www.ijphrd.com)

2

## Test of Hepatoprotector Effect of Merak Leaf Ethanol Extract (Caesalpinia Pulcherrima L) with SGPT Enzyme Parameter and SGOT of Induced Parasetamol Heart Rats (Rattus Norvegicus)

Gemy Nastity Handayani<sup>1</sup>, Trimaya Cahya M<sup>1</sup>

<sup>1</sup>Universitas Islam Negeri Alauddin Makassar

### ABSTRACT

Tests have been carried out research hepatoprotective Peacock Flower Leaves Ethanol Extract (*Caesalpinia pulcherrima L*) with Parameter enzymes SGOT and SGPT on Male Rat (*Rattus norvegicus*) induced Paracetamol. This study aims to assess the hepatoprotective effect of the ethanol extract of leaves of peacock flower (*Caesalpinia pulcherrima L*). In this study used 15 rats were divided randomly into 5 groups. Group I was given 1% CMC-Na as the control, group II was induced with paracetamol 1000 mg / kg for 3 days as the toxic dose to induce liver damage, group III, IV and V induced by paracetamol 1000 mg / kg for 3 days and then proceed with the extract ethanol leaf peacock flower (*Caesalpinia pulcherrima L*) for 7 days in a row with a dose of 270mg / 200 g BB rats, 540mg / 200 g BB rats, 1080mg / 200 gram BB rats. Before treatment, the mice blood drawn through the lateral vein then measured the levels of SGPT and SGOT early. After day 8, re-performed blood sampling to measure the levels of SGPT and SGOT end.

SGPT and SGOT The data were then analyzed using a Rancangan Acak Lengkap (RAL). The research proves that Paracetamol can cause hepatotoxicity with increased levels of SGOT and SGPT are high compared with the control group Na- CMC 1%. The results of this experiment indicate that the ethanol extract of leaves of peacock flower (*Caesalpinia pulcherrima L*) as a hepatoprotective effect in which the IV dose of extract (540 mg / 200 grBB mice) showed reduced levels of SGOT and SGPT largest, while the third dose of the extract (270 mg / 200 grBB mice) and V extract dose (1080 mg / 200 grBB mice) lacking both in effect hepatoprotective .

**Keywords:** Leaf Extract Peacock Flower (*Caesalpinia pulcherima*), hepatoprotective, Male Rat (*Rattus norvegicus*), SGPT & SGOT

### INTRODUCTION

The use of paracetamol as an analgesic and antipyretic has been known by the general public and many are sold freely on the market. This causes the community to consume it without having to use a doctor's prescription, in addition, public knowledge about the dangers of drug toxicity is still lacking, especially if used in excessive doses<sup>1,2</sup>.

The most common enzyme associated with liver cell damage is aminotransferase damage to liver parenchymal cells will increase levels of Serum Glutamic Pyruvate Transaminase (SGPT) and Serum Glutamic Oxaloacetic Transaminase (SGOT) in plasma. SGPT is more specific

than SGOT. SGOT is more in the myocardium than in liver cells, also SGOT is in the striated muscles, kidneys, and brain (Bastiansyah, 2012: 54). Liver damage can certainly affect liver function. But until now, there is no effective drug to stimulate liver function, protect liver cells from damage, and help regenerate liver cells even though the progress of modern medicine is growing rapidly. On the other hand, various attempts to treat liver function disorders are clinically expensive and often cause adverse side effects. Therefore, the community began to switch to traditional treatment in accordance with the motto "back to nature" which often had a significant effect. Until now there are also various studies to obtain active ingredient components that are

capable of acting as hepatoprotectors<sup>3-5</sup>.

Peacock flower is one of the popular ornamental plants of the tribe of legumes (Fabaceae), usually planted in the yard, sometimes clustered, with bright yellow-red flowers. Some chemicals contained in peacock flowers include tannin, gallic acid, resin, red matter, and benzoid acid. The leaves contain alkaloids, saponins, tannins, glucosides, calcium oxalates. While on bark contained plumbagin, lumbago, tannin, tanned substances, alkaloids, saponins, and calcium oxalate. Flavonoid content is found in all parts of the Peacock Flower plant, namely leaves, roots, wood, bark, pollen, nectar, fruit, and seeds<sup>6-8</sup>.

Peacock flower has various benefits, including blood circulation, menstruation and abortivum. Peacock flowers are used for the treatment of several diseases such as pyrexia, monoxenia, bronchitis and malaria infections. Peacock flower leaves are used as antipyretic, antimicrobial, antibacterial, and inflammation of the liver (hepatitis), canker sores, fever and skin diseases<sup>9</sup>. Empirically the community uses peacock flower leaves by boiling 30 g of peacock flower leaves are then drunk<sup>6</sup>. In accordance with the description of the efficacy of peacock flowers, a study on the effects of hepatoprotector on treatment for liver disease was selected based on the empirical experience of the existing community, with the aim of scientifically proving the experience of the community.

**MATERIALS AND METHOD**

**Sample extraction**

The 600 g peacock flower sample was put into a maceration container, moistened with 70% ethanol until all the simplicia was moistened, then ethanol was added until the simplicia was submerged. The maceration container is closed and stored for 24 hours in a place that is protected from direct sunlight while stirring

occasionally. Next filtered. Separated between pulp and filtrate. The pulp is re-extracted with the same amount of ethanol. This is done for 3x 24 hours. The ethanol extract obtained was then collected and evaporated by the liquid solution until a thick ethanol extract was obtained.

**Selection and preparation of experimental animals**

The experimental animals used were healthy male rats (*Rattus norvegicus*) and normal activities, body weight between 130-240 g. Male rats are adapted in cages for 7 days, adaptation is done to avoid the risk of stress that can affect blood serum content. During the adaptation period, male rats are only given standard feed and have not been given any treatment. At the end of the adaptation period a serum analysis was carried out in the five treatment groups. This is done to determine the normal state of SGPT and SGOT activities.

**Measurement of SGPT and SGOT of blood of test animals**

The principle of measuring SGPT and SGOT activity is measuring the rate of reduction in the amount of NADH to NAD + in the reaction that occurs between the enzyme and the substrate which can be measured at a wavelength of 340 nm. The rat blood sample was centrifuged at 3000 rpm for 15 minutes to get the serum. After that, ALT and AST levels were analyzed. A total of 100 µl of mouse blood serum were mixed with 1000 µl of reagent, then the absorption was measured by using a spectrophotometer at a wavelength of 340 nm, in the 1st, 2nd and 3rd minutes

**RESULTS**

From the results of research that has been done on Ethanol Extract of Kembang Merak Leaf (*Caesalpinia pulcherrima* L.) as a hepatoprotector tested in experimental animals, the following results were obtained:

**Table 1. Data on observations of SGPT and SGOT levels**

Treatment	Replication	Measurement			
		SGPT		SGOT	
		Early	End	Early	End
NA-CMC 1% (Control)	1	50	56	84	95
	2	73	80	60	76
	3	96	85	65	74

**Table 1. Data on observations of SGPT and SGOT levels**

Paracetamol 1000mg/ KgBB	1	65	118	62	174
	2	81	138	86	200
	3	76	143	69	198
Extract Dose I 270mg/200grBW rat	1	33	48	80	91
	2	57	77	67	89
	3	80	83	70	99
Extract Dose II 540mg/200grBW rat	1	67	73	78	85
	2	28	41	64	82
	3	46	59	68	96
Extract Dose III 1080mg/200grBW rat	1	64	86	88	122
	2	89	110	72	118
	3	80	97	65	106

## DISCUSSION

Serum Glutamic pyruvic transaminase (SGPT) / Alanine aminotransferase (ALT) is an enzyme that catalyzes the transfer of one amino group including alanine and alpha ketoglutarate acid. There are many in hepatocytes and their concentrations relatively low on other networks. Normal blood levels of 5-35 IU / liter and ALT are more sensitive than AST (Sacher and McPerson. 2011; 369). Serum glutamate oxaloacetic transaminase (SGOT) / Aspartate aminotransaminase (AST). AST is a mitochondrial enzyme that is also found in the liver, heart, kidneys and brain. If the tissue has acute damage, the level in serum increases. Allegedly this is due to the release of intracellular enzymes from damaged cells into the circulation. Very high levels are found in hepatocellular necrosis or myocardial infarction<sup>10</sup>.

In this study, we used a sample of peacock flower leaves (*Caesalpinia pulcherrima* (L) which contains many chemical compounds in the form of alkaloid flavonoids, saponins, tannins, glukoside, and calcium oxalate. , skin, pollen, nectar, fruit and seeds Flavonoids are phenol compounds that can be coagulators of protein and as antioxidants Traditionally, peacock flower leaves are used as a medicine for inflammation of the liver (hepatitis), canker sores, fever, skin disease<sup>9,10</sup>.

In this study using active ingredients of kembang merak leaves (*Caesalpinia pulcherrima* (L) extracted using 70% ethanol dancr. Ethanol is a universal dancr which can dissolve polar compounds and nonpolar compounds. Ethanol is the best for producing optimal active compounds is ethanol 70%<sup>11</sup>. The extraction

method used is maceration method. This method was chosen because it has several advantages, namely the way the process is easy, the tools that are used are simple, suitable for materials that are not heating resistant. The maceration method is done by soaking simplicia as much as 600 g in 70% ethanol. The dancr liquid will penetrate the cell wall and enter the cell cavity containing the active substance, the active substance will dissolve and because of the difference in concentration between the solution of the active substance in the cell and the outside of the cell, the concentrated solution is pushed out. The event continued so that there was a concentration equilibrium<sup>1-5</sup>. This method is done 1 X 24 hours which is done 3 times. Maserat results later it was relayed at 600C and then evaporated until a thick extract was obtained. Extract obtained is stored in the desiccator. Then the extract is weighed using an analytical balance.

Rat that will be used are first adapted for 1 week which aims to condition animals with a laboratory atmosphere. To induce liver damage, toxic doses of paracetamol were used (1000 mg / kg). Rat that will be used are first adapted for 1 week which aims to condition animals with a laboratory atmosphere. To induce liver damage, toxic doses of paracetamol were used (1000 mg / kg). According to Abraham's research, the administration of paracetamol at a dose of 1000 mg / kg body weight has been shown to show liver damage characterized by elevated levels of SGPT enzymes. This is also supported by the research of Roy and Das where administration of paracetamol at a dose of 1000 mg / kg BW orally in male white rats after 48 hours showed an increase in ALT, AST, ALP, and serum bilirubin levels.

From the data obtained then it is processed by

calculating the Completely Randomized Design (CRD) which can be seen in appendix 4 where F Calculate > F Table at the confidence level of 5% and 1%, this means that there is a very significant difference from each variety of doses of extract ethanol flower peacock (*Caesalpinia pulcherrima* L). The results of measurement of the average level of SGPT the dose groups of extracts I, II and III were 63 U / L, 52.3333 U / L and 87.66667 respectively U / L and SGOT mean levels are 81.33333 U / L, 80.16667 U / L and 95.16667 U / L, respectively. It can be seen from the three dosage groups that SGPT and SGOT enzyme levels decreased significantly with the Na-CMC and Paracetamol control groups. Although SGPT and SGOT enzyme levels decreased from all three doses, only the dose group extract II (540 mg / 200 gBB) showed a better reduction in the value of SGPT and SGOT enzymes. The results of the calculation of percent (%) decreased SGPT levels in the Na-CMC Control group, the Paracetamol group (1000mg / 200g BB Mice), group I (270mg / 200g BB Mice), group II (540 mg / 200g BB Mice) and Group III (1080 mg / 200 g BB Mice), respectively 10.7%, 44.3%, 20.2%, 12.5% and 20.6%. From the data generated shows that group IV (540 mg / 200 g BB Mice) shows a percent decrease which is almost close to the Na-CMC control group, which means that group IV can reduce SGPT levels well compared to other groups. While the results of the calculation of percent (%) decreased SGOT levels in the Na-CMC Control group, Paracetamol group (1000mg / 200g BB Mice), group I (270mg / 200g BB Mice), group II (540 mg / 200g BB Mice) and Group III (1080 mg / 200 g BB Mice), respectively 14.8%, 62.1%, 21.9%, 19.7% and 35.1%. From the data generated shows that group IV (540 mg / 200 g BB Mice) shows a decrease in percentage which is almost close to the Na-CMC control group compared to other groups, which means that group IV can reduce SGOT levels well compared to other groups.

The decrease in the levels of SGPT and SGOT enzymes is one indication of the healing of liver cells damaged by paracetamol after the administration of peacock leaf extract (*Caesalpinia pulcherrima* L). This is because the leaves of the peacock flower contain a group of compounds including flavonoids which are antioxidants. Antioxidant compounds can act as neutralizing free radicals produced by the metabolites of paracetamol. The antioxidant activity of flavonoids found in peacock leaves can be associated with the potential

of peacock hepatoprotector. According to research by Sulistiyani found secondary metabolites (flavonoids and phenolics) that have the potential as antioxidants can reduce lipid peroxide formation in the blood of rats damaged by their paracetamol. The mechanism of action of antioxidant compounds by giving electrons or stopping the reaction of free radicals, so as to prevent the continued chain reaction from fat peroxidation and also proteins due to the effects of free radicals, thus further cell damage can be prevented.

In this study paracetamol was used as a liver damage induction agent. This is because high doses of paracetamol will consume the conjugated capacity of glucuronic acid and sulfuric acid, so that the formation of NAPQI reactive metabolites increases over the GSH conjugate capacity. This NAPQI will then bind covalently with vital liver cell macromolecules (lipids and membrane proteins) so as to cause liver damage. Damage to liver cells results in an increase in SGPT, SGOT, alkaline phosphatase, etc. Besides chemical agents, stress factors such as lack of oxygen supply, excessive physical activity, trauma, unstable environment temperature, is also one of the causes of cell damage.

## CONCLUSION

From the results of observations that have been made on Ethanol Extract of the Leaf of Peacock Flower (*Caesalpinia pulcherrima* L) on test animals it can be concluded that: 1. Ethanol Extract of Leaves of Peacock Flower (*Caesalpinia pulcherrima* L) dose I (270mg / 200 gram BB rat), dose II (540mg / 200 gram BB rat) and dose III (1080mg / 200 gram BB rat) showed hepatoprotector effect. 2. The most effective dose as hepatoprotector is the dose of extract III (540 mg / 200 gBB)

**Conflict of Interest :** None

**Source of Funding:** Middle cluster dictis.

**Ethical Clearence:** Obtained from faculty committee

## REFERENCES

1. Director General of POM. Indonesian Pharmacopoeia. Ed. III. Jakarta: Indonesian Ministry of Health, 1979.
2. Fachruddin, H. Plant Phytochemical Analysis. Faculty of Pharmacy. Hasanuddin University. Makassar, 2001.

3. Ismeri, Ethanol-Water Extract Activity Curry leaves (*Murraya kuenigii*) as Hepatoprotector in sprague Dawley rats, Bogor: Faculty of Mathematics and Natural Sciences, Bogor Agricultural University. 2011
4. Corwin, E.Z. Pathophysiology pocket book. Happened. From the Handbook of pathophysiology by Brahm, U. Medical Book Publishers EGC, 2010; 3rd revised edition of Jakarta
5. RI Ministry of Health. How to Manage Good Simplicia. Directorate General of Drug and Food Control, Jakarta, 1999.
6. Hembing Wijaya, Tumpas Hepatitis with herbal ingredients. Agromedia library Jakarta, 2010
7. Pulubuhu, D.A.T., Evans, K., Arsyad, M., Mallongi, A. Understanding the perspectives of village leaders and institutions in transforming social conflict into peace and health. *Indian Journal of Public Health Research & Development*, March 2018, Vol. 9, No. 3
8. Amran, Stang, and Anwar Mallongi, AIP Conference Proceedings **1825**, 020002 2017; doi: 10.1063/1.4978971
9. Wijayakusuma, M.H., Dalimartha, S., Wirian, A. *Medicinal Plants in Indonesia*, 2008; Volume IV Kartini Library, Jakarta
10. Sacher and McPerson, *Clinical Review of Laboratory Examination Results*. Edition Medical Book Publisher EGC. Jakarta, 2011
11. Voight, Rudolf. *Pharmacy Technology Textbook*. Gadjah Mada University Press. Yogyakarta, 1995.
12. Arianti, Rini, Acute Hepatoprotective and Toxicity Activity of alang-alang root extract (*Imperata Cylindrica*), Thesis, Bogor, Faculty of Mathematics and Natural Sciences, Bogor Agricultural Institute, 2012
13. Bastiansyah, Eko, *Complete Guide to Reading Health Test Results*. Penebar Plus, Jakarta 2012
14. Mescher, A.L. *Juncuiera's Basic Histology Text & Atlas* 12<sup>th</sup> ed. New York : The McGraw-Hill Companies, Inc. 2010
15. Sriasih, N.G.K., Ariyani, N.W., Mauliku, J., Budiani, N.N., Mallongi, A. Frangipani aromatherapy oil in the massage of labor first stage reduced events perineum ruptur spontan at the labo. *Indian Journal of Public Health Research & Development*. 2018.

# Test of Hepatoprotector Effect of Merak Leaf Ethanol Extract (Caesalpinia Pulcherrima L) with SGPT Enzyme Parameter and SGOT of Induced Parasetamol Heart Rats (Rattus Norvegicus)

## ORIGINALITY REPORT

4%

SIMILARITY INDEX

4%

INTERNET SOURCES

0%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

2%

2

[directory.indianjournals.com](http://directory.indianjournals.com)

Internet Source

1%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On