

# A Review on Selected Tropical Tree Legume Leaves in Poultry diet

The screenshot shows a Yahoo Mail interface. The browser address bar displays a search URL for 'Transylvanian Review' with email addresses. The navigation bar includes categories like AWAL, MAIL, BERITA, KEUANGAN, OLAHRAGA, SELEB, LIFESTYLE, and LAINNYA... The user is logged in as 'Muhammad'.

The left sidebar shows the email folder structure: Email Masuk (29), Belum Dibaca, Berbingung, Draft (583), Terkirim, Arsip, Spam, Sampah, and a 'Folder Sembunyikan' section with '+ Folder Baru'.

The main inbox area shows a list of emails from 'transylvanian review' in 2020 and 2019. The selected email is dated 13/3/2020 with the subject 'Re: Acceptance letter for our p...'. The email content is as follows:

**transylvanian review** <t...> Jum, 13 Mar 2020 jam 03.55  
Kepada: Muhammad Rusdy

Dear Muhammad Rusdy  
Please send your corrected full papers.

Sembunyikan pesan asli

On Thu, Feb 27, 2020 at 5:40 PM Muhammad Rusdy <muhrusdy79@yahoo.co.id> wrote:  
Dear Sir,  
Please forgive me because I forget to send acceptance letter for our second paper (title A review of selected tree legume leaves in poultry diet). The first author of the paper was Sofyan Nurdin Kasim, but because he could not able to pay the the article processing fee, I proposed to you to replace him with Andi Suarda as the first author, me as the second author and Tanrigiling Rasyid as the third author. Because you agreed to my proposal, we paid the article processing fee. Andi Suarda told me that he is very need the paper published because he will use the pulished paper to be proposed as professor. Thank you for your help and appreciation.

Kind regards

On the right side, there is a profile card for 'Transylvanian Review' with the email 'transylvanianreviewjournals@gmail.com' and a 'Dapatkan App-nya' advertisement for 'yahoo!mail TERATUR SPESIAL KEREN'.

Tulis

- Email Masuk 29
- Belum Dibaca
- Berbintang
- Draft 583
- Terkirim
- Arsip
- Spam
- Sampah
- Lebih sedikit
- Tam... Sembunyikan
- Foto
- Dokumen
- Langganan
- Promo
- Folder Sembunyikan
- + Folder Baru

Kembali


Transylvanian review Ok, no problem. On 2 Jul 2020 T Kam, 2 Jul 2020 jam 06.04

Transylvanian review <transylvanianreview> Kepada: Yahoo Desk Sab, 4 Jul 2020 jam 03.40

Dear Author  
Please find your full paper as attached. Also, please send your papers for further processing of the next issue.

Tampilkan pesan asli

Kind Regards  
Vasile Salijan  
Editor in Chief



TR-012333-1....pdf  
363kB

TR-012333-1... Halaman 2 dari 9

Andi Suardi et al  
Transylvanian Review Vol XXVIII, No. 46, Dec 2019

### A Review on Selected Tropical Tree Legume Leaves in Poultry Diet

Andi Suardi<sup>1</sup>, Muhammad Rusdy<sup>2</sup> and Tamri Giling Bayu<sup>1,2</sup>

<sup>1</sup>Faculty of Science and Technology, Alauddin Islamic State University, Makassar, Indonesia  
<sup>2</sup>Faculty of Animal Science, Hasanudin University, Makassar, Indonesia

**ABSTRACT**

The high prices of protein sources from fish meal and soybean meal are the main constraints to increasing poultry production in developing countries. One way to alleviate the problems is through utilization of fodder tree legume leaves to replace a part of fish meal and soybean meal in poultry diet. Fodder tree legumes of *Albizia chinensis*, *Leucaena leucocephala*, *Gliricidia sepium*, *Calliandra calothyrsus*, *Acacia mangium* and *Sesbania grandiflora* leaves have been studied as protein sources in poultry diets. To generate information concerning chemical composition and effect of incorporating tree legume leaves in diet on poultry performance, the literature in Google Scholar, Scopus, CAB Abstract, Pubmed and CrossRef databases were reviewed. Results revealed that leaves of those tree legumes contain protein higher than required by poultry but also have higher fiber than can be tolerated by poultry. They also contain lower metabolizable energy than conventional concentrate or proteinaceous concentrate and have nutritional factors that may hinder poultry performance. Compared with concentrate feeds, all those legume leaves can only be incorporated in poultry diets at low levels without affecting poultry performance. Except for *Sesbania grandiflora* leaves that toxic to poultry, all other tree legume leaves can be incorporated in poultry diets. The best growth performance of birds fed on high levels of those legume leaves was found in the leaves of *Albizia chinensis*, followed by *Leucaena leucocephala*, *Gliricidia sepium*, *Acacia mangium* and *Calliandra calothyrsus* leaves. It can be concluded that those tree legume leaves can be included in poultry diets with varying optimum levels.

**Keywords:** Antinutrients, Incorporation, Nutrients, Poultry performance, Tree legume leaves.

**Introduction**

Feed cost constitutes about 70 - 80% of the total cost of poultry production. In many developing country like Indonesia, scarcity and high prices of quality feeds containing high protein are the main constraints to increasing poultry production. In the recent years, increasing price of conventional protein sources such as soybean meal and fish meal has increased feed cost enormously. Therefore, there is a great need to exploit alternative locally available cheap protein sources which can replace a proportion of those ingredients which do not compete with human needs. A possible cheap protein source that can be utilized for this purpose is fodder tree legume leaves. Because of its ability to fix atmospheric N, fodder tree legume leaf generally contains higher protein than other family of plants. Leaves of tropical legumes are good protein source that can be exploited due to their nutritive value, availability, relatively low cost and non-competition of human (Corunha et al., 2010). Fodder tree legume leaves also have other nutrients like mineral and vitamin needed by poultry (Agbede, 2006) and have excellent potential of pigment due to their high xanthophyll content (Dilakic and Jovanovic, 1999). Due to their high-palating potentials and less susceptible to adverse climatic fluctuations, the utilization of tree legume leaves can be expected to be a sustainable feed source for reducing feed cost in humid tropical country like Indonesia.

The main problems that may limit the use of fodder tree legume in poultry feeding are high fiber, low palatability, low energy and high moisture contents (Makarewicz et al 2018). Incomplete essential amino acid composition and presence of anti-nutritive factors like toxic amino acids, tannins, cyanogenic glycosides etc. also limit their potential as good feed for poultry (Dilakic, 1999). Indicators of high fiber ingredients in poultry diet may reduce nutrients digestibility and nitrogen excretion (Dijger et al., 2004). Despite their negative effect on poultry performance, low to moderate amounts of fiber might be acceptable for broilers and development of a palatable level, thereby can enhance nutrients digestibility and growth performance (De Vries, 2018). Poultry producers have been advised to feed forages to