

A Review on Selected Tropical Tree Legume Leaves in Poultry Diet

by Andi Suarda

Submission date: 15-Sep-2020 01:57PM (UTC+0700)

Submission ID: 1387523714

File name: A_review_of_selected_tropical_tree_legume_leaves_1_3.docx (50.48K)

Word count: 4767

Character count: 26326

5

TRANSYLVANIAN REVIEW

VOL XXVII, No. 45, 2019



Transylvanian Review

Centrul de Studii Transilvane | str. Mihail Kogalniceanu nr. 12-14, et.5, Cluj-Napoca

Email: transylvanianreview@gmail.com

Online Submission System: <http://transylvanianreviewjournal.org/>

A Review on Selected Tropical Tree Legume Leaves in Poultry Diet

Andi Suarda¹, Muhammad Rusdy², Tanri Giling Rasyid²

¹Faculty of Science and Technology, Alauddin Islamic State University, Makassar, Indonesia

²Faculty of Animal Science, Hasanuddin University, Makassar, Indonesia

Corresponding E-mail: andi.suarda@uin-alauddin.ac.id

Abstract

The high prices of protein sources from fish meal and soybean meal are the main constraints to increasing poultry production in developing countries. One way to alleviate the problems is through utilization fodder tree legume leaves to replace a part of fish meal and soybean meal in poultry diet. Fodder tree legumes of *Moringa oleifera*, *Leucaena leucocephala*, *Gliricidia sepium*, *Calliandra calothyrsus*, *Acacia angustissima* and *Sesbania grandiflora* leaves have been studied as protein sources in poultry diets. To generate information concerning nutritional composition and effect of incorporating tree legume leaves in diet on poultry performance, the literatures in Google Scholar, Scopus, CAB Abstract, PubMed and CrossRef databases were reviewed. Results revealed that leaves of those tree legumes contain protein higher than required by poultry but also have higher fiber that hardly be tolerated by poultry. They also contain lower metabolizable energy than carbonaceous concentrate or proteinaceous concentrates and have antinutritional factors that may hinder poultry performance. Compared with concentrate feeds, all those legume leaves can only incorporated in poultry diets at low levels without affecting poultry performance. Except for *Sesbania grandiflora* leaves that toxic to poultry, all other tree legume leaves can be incorporated in poultry diets. The best growth performance of birds fed on high levels of those legume leaves was found in the leaves of *Moringa oleifera*, followed by *Leucaena leucocephala*, *Gliricidia sepium*, *Acacia angustissima* and *Calliandra calothyrsus* leaves. It can be concluded that those tree legume leaves can be included in poultry diets with varying optimum levels.

Key-words: antinutrients, incorporation, nutrients, poultry performance, tree legume leaves

Introduction

Feed cost constitutes about 70 - 80% of the total cost of poultry production. In many developing country like Indonesia, scarcity and high prices of quality feeds containing high protein are the main constraints to increasing poultry production. In the recent years, increasing price of conventional protein sources such as soybean meal and fish meal has increased feed cost enormously. Therefore, there is a great need to exploit alternative locally available cheap protein sources which can replace a proportion of protein sources which do not compete with human needs.. A possible cheap protein sources that can be utilized for this purpose is fodder tree legume leaves. Because of its ability to fix atmospheric N, fodder tree legume leaf generally contains higher protein than other family of plants. Leaves of tree legumes are good feed source that can be exploited due to their high availability, nutritive value, relatively low cost and non existence of competition in their consumption between animals and human (Oloruntola *et al.*, 2016). Fodder tree legume leaves also have other nutrients like mineral and vitamin needed by poultry (Agbede, 2006) and have excellent potential of pigment due to their high xanthophyll content (D'Mello and Acamovic,1989). Due to their high-yielding potentials and less susceptible to adverse climatic fluctuations, the utilization of tree legume leaves can expected to be a sustainable feed sources for reducing feed cost in humid tropical country like Indonesia.

Methodology

Recently published literatures in Google Scholar, CAB Abstract, Scopus, PubMed and CrossRef databases using key-words of antinutrients, incorporation, nutrients, and tree legume leaves were searched through July to October 2019. The synonym and antonym of

The main problems that may limit the use of fodder tree legume in poultry feeding are high fiber, low palatability, low energy and high moisture contents (Ndelekwute *et al.* 2018). Incomplete essential amino acid composition and presence of anti-nutritive factors like toxic amino acids, tannins, cyanogenic glycosides etc. also limit their potential as good feed for poultry (D'Mello, 1991). Inclusion of high fiber ingredients in poultry diet may reduce nutrients digestibility and increased nitrogen excretion (Dilger *et al.*, 2004).

Despite their negative effect on poultry performance, low to moderate amounts of fiber might be beneficial for function and development of gastrointestinal tract, thereby can enhance nutrients digestibility and growth performance (De Vries, 2015). Poultry producers have been advised to feed forages to their animals to meet the fibre, protein and vitamins requirements and reducing feed cost (Ndelekwute *et al.*, 2018)

Of about 12 000 species of legume (NAS, 1979), there are about one hundred of legume species have been studied as poultry diet. Although considerable work on incorporation of tree legume leaves in poultry diets have been published, however they have been rarely reviewed. Therefore, the objective of this paper was to review the available literatures on chemical composition and the effects of incorporating of selected fodder tree legume leaves (*Leucaena leucocephala*, *Gliricidia sepium*, *Calliandra calothyrsus*, *Acacia angustissima* and *Sesbania grandiflora*) in diets on poultry performance.

those key-words was used for searching relevant literatures. Of about 100 literatures were read, 58 literatures were randomly selected for review. Chemical compositions of each legume species leaves from 2 – 5 literatures and effect of incorporating of each legume leaf meal on poultry performance from the literatures were presented and reviewed.

Results

Chemical composition and effects of inclusion of selected fodder tree legumes on poultry

performance are shown in Table 1 and Table 2, respectively.

Table 1. Nutritional composition and antinutritional substances in selected fodder tree legume leaves

| Plant species | Protein (%) | Fiber (%) | Ca (%) | ME Kcal/kg | Anti-nutrients | Author |
|-------------------------------|-------------|-----------|--------|------------|---|---|
| <i>Moringa oleifera</i> | 28.00 | 12.57 | | 2625 | Tannins, saponin, oxalate, phytate | Okiki <i>et al.</i> , 2015 |
| | 25.27 | 9.92 | 1.70 | 2852 | | Kumar <i>et al.</i> , 2018 |
| <i>Leucaena leucocephala</i> | 29.17 | 13.77 | | | | Adekojo <i>et al.</i> , 2014 |
| | 24.9 | 14.20 | 1.80 | 2573 | Cyanogenic glycoside, phytate, tannin, saponin, mimosine, Oxalate | Ayssiwede <i>et al.</i> , 2010 Agbo <i>et al.</i> , 2017 |
| <i>Gliricidia sepium</i> | 22.88 | 10.12 | | | Saponin, phytate, tannins, cyanide | Paranamana <i>et al.</i> , 2015 |
| | 24.37 | 12.47 | | 3162 | Cyanide, tannin | Oloruntola, 2018 |
| | 28.4 | | 1.93 | | | Yousuf <i>et al.</i> 2007 |
| <i>Calliandra calothyrsus</i> | 25.30 | 19.1 | | | Tannin, | Salawu <i>et al.</i> , 1997 |
| | 17-28 | | 1.60 | 2147 | | Machanja, 2007 Pamo <i>et al.</i> , 2005 |

| | | | | | | |
|-----------------------------|-------|-------|------|------|---|---|
| <i>Acacia angustissima</i> | 22.13 | 9.86 | 1.20 | | Tannins | Ncube <i>et al.</i> , 2012 |
| | 23.4 | 13.00 | 0.94 | | Tannin Tannin, non-protein amino acids | Ncube <i>et al.</i> , 2018 Odenyo <i>et al.</i> , 2003 |
| <i>Sesbania grandiflora</i> | 34.9 | 7.47 | 2.42 | | | Reddy and Elanchezhian, 2008 |
| | 32.3 | 6.70 | 3.26 | 1477 | Tannins, canavanine Saponin | Prasad <i>et al.</i> , 1970 Sarvade and Parmar, 2014 Anonymous, 2019 Mlay <i>et al.</i> , 2006 |

Table 2. The effects of incorporating of fodder tree legume leaf meals on performance of poultry

| Plant species | Type of chicken, leaf meal doses | Growth/laying performance | Authors |
|------------------------------|---|---|--------------------------------|
| <i>Moringa oleifera</i> | Broiler chicken, doses: 0, 25, 50, 75, 100% | Up to 25% inclusion level, growth rate, final weight and feed intake were similar | Gadzirayi <i>et al.</i> , 2012 |
| | Broiler chicken, study lasted for four weeks Doses : 0, 12.5, 25, 37.5 and 50% | Growth rates and feed conversion were similar up to 25% level | Imoru <i>et al.</i> , 2019 |
| <i>Leucaena leucocephala</i> | Broiler chicken, Doses: 5, 10, 15 and 20% | Up to 15% level, weight gain and feed efficiency was similar but at 20% level, weight gain and feed efficiency were adversely affected. | Hussain <i>et al.</i> , 1991 |
| | Broiler, doses: 0, 7, 14 and 21% | Up to 21% level, intake, weight gain, feed conversion rate and nutrients utilization were similar. | Ayssiwede <i>et al.</i> , 2010 |

| | | | |
|-------------------------------|--|---|--|
| <i>Gliricidia sepium</i> | Broiler, from day 1 to 56 Doses 0, 5 and 10%. | At 5% inclusion level, weight gain, final weight, feed intake and feed conversion rate were not significantly differ with control | Oloruntola, 2018 |
| | Laying hens, doses: 0, 5, 10 and 15%. | At 0 and 5% level, egg production, body weight gain and feed conversion efficiency were similar, but at 10 and 15% level, they worsened | Odunsi <i>et al.</i> , 2002 |
| <i>Calliandra calothyrsus</i> | Laying hens, study lasted for 67 days Doses: 0, 5, 7.5, 10% | Feed intake increased as increasing leaf meal levels, but egg produced, weight gain and feed efficiency were lower compared with control | Paterson <i>et al.</i> , 1996 |
| | Layers, doses: 0, 5, 10 and 15% | Up to 5% level, no decreased in egg production and average egg weight, but at 10%, egg production and feed intake decreased, egg weight increased. | Patterson <i>et al.</i> , 2000 |
| <i>Acacia angustissima</i> | Broiler age of 1 to 28 days, doses: 0, 5, 10 and 15% | Feed intake, weight gain and feed conversion ratio at 0 and 5% was similar, but weight gain at 10 and 15% level was lower | Ncube <i>et al.</i> , 2012 |
| | Broiler, age of 1 to 42 days Doses: 0, 5, 10% | Up to week 4, feed intake, weight gain and feed conversion were similar across treatments, but at weeks 4 and 6, control and 5% group outperformed the 10% group. | Ncube <i>et al.</i> 2018 |
| <i>Sesbania grandiflora</i> | Growing chicken, doses : 0, 5, 10, 15% | Feed intake increased but final weight decreased as increasing leaf meal level. | Prasad <i>et al.</i> , 1970 |
| | Day-old chicks, doses: 0, 5, 10% | Feeding leaf meal increased mortality, reduced feed intake and growth rates | Ash and Petaia, 1992; Shqueir <i>et al.</i> , 1989 |

Discussion

1. Chemical composition of selected fodder tree legume leaves

Protein is an important nutrient for poultry and plays a significant role in growth, egg production and other biological functions. Protein contents of tree legume leaves ranged from 22.13% to 34.90% (Table 1). These values are above the requirements of both broiler (18 – 23%) and layer (15 – 18%) (NRC, 1994). However, compared with animal protein, contents of certain amino acids of tree legume leaves might be lower and their essential amino acid compositions are not complete, because according to Gorrissen *et al.* (2018), lysine and methionine are typically lower in plant based protein than animal based protein and there are differences in essential amino acid and non-essential amino acid composition among various plant-based isolates.

Fiber is naturally found in plants and can interfere with digestibility of feed. Fiber contents of tree legume leaves are varied between plant species. The highest value was found in *Calliandra calothyrsus* (18.46 – 22.56%) and the lowest was found in *Sesbania grandiflora* (6.70%) (Table 1). These values are above of 3 – 4% that is considered as maximum value of fiber content in broiler diet (Sw₃nen *et al.*, 2010) or up to 5% that is beneficial for gastrointestinal development, function and health and enhancing nutrient digestibility (De Vries, 2015). Inclusion of high levels of fiber in poultry diet have negative impact on voluntary feed intake and nutrient digestibility (Mateos *et al.*, 2012).

Calcium (Ca) contents of selected legume tree leaves are differ widely, the values ranged from 1.60% in *Calliandra calothyrsus* to 3.26% in *Sesbania grandiflora* (Table 1). Calcium content of *Sesbania grandiflora* leaves are comparable with other high-protein non-leguminous plants like sweet potato (*Ipomoea batatas*) leaves (2.10%) (Tsega and Tamir, 2009) and *Morus alba* leaves (2.96%) (Herrera *et al.*, 2019), while

calcium content of *Leucaena leucocephala* leaves was comparable with *Moringa oleifera*, *Gliricidia sepium*, and *Calliandra calothyrsus* leaves (Table 1) but higher than paw-paw leaf extract (1.09%) (Ugo *et al.*, 2019). Supplementation with high-calcium leaf can be used to increase egg shell thickness. Inclusion of *Moringa oleifera* leaf meal at 15% level has been reported could improve egg shell strength (Lu *et al.*, 2016).

Metabolizable energy (ME) is portion of energy utilized by the animal. ME contents of selected fodder tree legume leaves are vary, ranged from 1477 kcal/kg (*Sesbania grandiflora*) to 3162 kcal/kg (*Gliricidia sepium*) leaves (Table 1). These values were lower than ME in corn (3849 kcal/kg DM and soybean 3496 kcal/kg DM (Tesfaye *et al.*, 2013). The low ME contents of tree legume leaves is due to the high fiber contents. Increasing fiber intake decreased nutrients digestibility and consequently, ME contents of fibrous feed decreased (Baer *et al.*, 1997). ME level affects digestibility of DM, energy and crude protein, and enzyme supplementation improved energy digestibility in diets with lower levels of ME (Zhou *et al.*, 2009). Birds fed on diets containing 11.51 MJ kg⁻¹ consumed more than diets containing 13.81 MJ kg⁻¹, whereas gain and feed conversion ratio improved when enzyme was added to 11.51MJ kg⁻¹ diet (Moftakharzadeh *et al.*, 2019)

Antinutritional factors are toxic compounds of natural or synthetic origin which interfere with absorption of nutrients, reduce nutrient intake and utilization and may produce other adverse effect (Popova and Mihaylova, 2019). All selected tree legume leaves contain antinutritional factors with tannin is the most widespread, followed by saponin, oxalate and phytate. Cyanogenic glycoside, mimosine and canavanine only found in certain plants. Most of antinutrients were found in *Leucaena leucocephala* and the least was found in *Calliandra calothyrsus* (Table 1). Tannin is an antinutrient that may impair digestion of various nutrients. Like in ruminants, in poultry tannin impairs feed utilization by forming complexes with protein and carbohydrate that reducing nutrients availability (Medugu *et al.*, 2012).

2. Effect of incorporating tree legume leaves in the diets on poultry performance

Effect of incorporating of various tree legume leaf meal in diets on poultry performance were different between species of legume and levels of leaf meal incorporation. The highest tolerance level of birds to leaf meal incorporation obtained when the birds fed *Moringa oleifera* leaf meal (25% level), followed by *Leucaena leucocephala* leaf meal (18% level), *Gliricidia sepium* leaf meal (5% level), *Acacia angustissima* (5% level), *Calliandra calothyrsus* (2.5% level) and the lowest was found in *Sesbania grandiflora* that may kill the birds even at low levels. The values above those levels had negative effects on poultry performance (Table 2).

The adverse effect of incorporation of high levels of tree legumes leaf meal might be in part, attributed to the high levels of fiber in diets because digestive system of poultry has limited capacity to digest fibrous feed efficiently as they lack of enzymes for digestion of high fibrous feed. A high proportion of leaf meal in diet may reduces nutrients digestibility and increased nitrogen excretion (Dilger *et al.*, 2004). However, in the present study, there is a lack of consistency with regard to dietary fiber content and growth performance of poultry. For example, *Sesbania grandiflora* leaves with low crude fiber content (averaged 7.80%), even at 5% level was harmful to birds performance, but *Moringa oleifera* leaves with higher fiber content (averaged 11.28%) yielded better growth performance, even though they were incorporated in poultry diets at high level (25%). (Table 1 and Table 2).

Other factors such as protein quality (composition and concentration of amino acids) and presence of antinutritional factors in the leaf meal may also affect growth performance of poultry. The good source of essential amino acids in *Moringa oleifera* leaves (Moyo *et al.*, 2011) might compensate the adverse effects of the high fiber contents in *Moringa oleifera* leaves. Conversely the lacks of sulphur containing amino acids in *Leucaena*

leucocephala (Tantung and Madamba, 1981), *Gliricidia sepium* (Alonso *et al.*, 2003) and *Calliandra calothyrsus* (Salawu *et al.*, 1997) may affect poultry performance fed on those legumes. *Acacia angustissima* and *Sesbania grandiflora* leaves might also lack of sulphur amino acids because legume plants are generally low in sulphur amino acids (Bressani and Elias, 1974).

Antinutritional factors as listed in Table 1 might also affect poultry performance. However, in the present study, there is no correlation between number of anti nutrient present in any legume species and poultry performance. Number of antinutrients was mostly found in *Leucaena* leaves and the least was found in *Calliandra* leaf (Table 1), however *Calliandra* was more harmful to bird performance than *Leucaena* (Table 2). It seems that besides kinds of antinutrient, concentration of antinutrient also is important factor that affect poultry performance. The higher condensed tannin contents in *Calliandra* (6.0 to 16.0%) (Rusdy, 2016) than *Leucaena* (3.79%) (Aye and Adegun, 2003) might be the reason for the low performance of poultry fed on *Calliandra* leaf meal. Although *Sesbania grandiflora* leaves contain high protein, calcium and low fiber, feeding of *Sesbania* leaves to poultry is not recommended because it contains saponin which toxic to young chick (Anonymous, 2019). Feeding of 20% *Sesbania* leaf meal killed 47% chick reared (Shqueir *et al.*, 1989). Saponin is antinutrients that may be beneficial because it can be used as feed additive for disease prevention and growth promoter, but at high levels can be toxic to poultry (Wina *et al.*, 2017).

To increase utilization of dietary inclusion of leaf meal in poultry diet, the use of exogenous enzyme and supplementation with sweetener ingredients (Ige *et al.*, 2006) and treatments that will lead to reduction or elimination of antinutrients should be used before leaf meal fed to poultry.

Despite their promises as supplement in poultry diets, however, there is a great variation in chemical composition of fodder tree legume leaves and their effects on poultry performance. Therefore, research on the evaluation of the effects of incorporating tree legume leaves on

poultry performance should be carried out wherever the plant occurs in abundance.

Currently, in Indonesia both promising plants, *Moringa oleifera* and *Leucaena leucocephala* easily found elsewhere, however *Moringa oleifera* leaf tends to be less competitive as poultry feed because it is commonly used as vegetable in human food. At present time, among the selected tree legume leaves, *Leucaena leucaena* leaf meal might be the best ingredient because besides they can be tolerated at high levels in poultry diets, it is also has higher leaf biomass potential and rarely used as human food. In the future, mass-planting of *Moringa oleifera* may be the best option to lower the high feed cost in poultry production.

References

Adekojo, S.A., Adama, T.Z., Aremu, A., Ijaiya, A. T., 2014. Effects of different method of processing *Leucaena leucocephala* leaf meal on growth performance and nutrient digestibility of rabbits. *Int. J. Agric. Forest.* 4 (5) : 380 – 385.

Agbede, J.O., 2006. Characterization of leaf meal, protein concentrates, and residues from some tropical leguminous plants. *J. Food Sci. Agric.*, 86 (9) : 1292 – 1297.

Agbo, A.N., Balogun, J.K., Oniye, S.J., Auta, J. 2017. Effect of differing processing methods on nutritional composition of *Leucaena leucocephala* (Lam De Wit) leaves as inclusion in fish feed. *J. Appl. Sci., Environ. Mngmt*, 21 (4) : 719 – 725.

Alonso, R., Pedraza, R.M., Apori, S.O., Orskov, E.R., 2003. Some chemical and biological measurements of two contrasting cultivars of *Gliricidia sepium* (Jack) Kunth ex Walp. *Livest. Res. Rural Dev.*, 15 (7).

Anonymous, 2019. *Sesbania grandiflora*. [https://uses.plantnetproject.org/en/Sesbania-grandiflora_\(PROSEA\)](https://uses.plantnetproject.org/en/Sesbania-grandiflora_(PROSEA)). Accessed on November 5, 2019.

Conclusion

1. Fodder tree legume leaves can be incorporated in poultry diets at low levels
2. The best tree legume leaves used as supplement in poultry diets is *Moringa oleifera* and the worst is *Sesbania grandiflora*
3. There is great variation in nutritional composition and efficiency of inclusion of fodder tree legume leaves in diets on poultry performance.

Ash, A.J., Petaia, L. 1992. Nutritional value of *Sesbania grandiflora* leaves for monogastrics and ruminants. *Trop. Agric.*, 69 (3) : 223 – 228.

Aye, P.A., Adegun, M.K., 2013. Chemical composition and some functional properties of *Moringa*, *Leucaena* and *Gliricidia* leaf meal. *Agric. Biol. J. North Am.*, 4 (1) : 71 – 77.

Ayssiwede, S.B., Dieng, A., Chrysostome, C., Ossebi, W., Hornick, J.L., Missohou, A. 2010. Digestibility and metabolic utilization and nutritional value of *Leucaena leucocephala* leaf meal incorporated in the diets of indigenous Senegal chicken. *Int. J. Poult. Sci.*, 9 (8) : 767 – 776.

Baer, D.J., Rumpler, W.V., Miles, C.W., Fahey, G.C., 1997. Dietary fiber decreases metabolizable energy content and nutrient digestibility of mixed diets fed to humans. *J. Nutr.*, 127 (4) : 579 – 586.

Bressani, R., Elias, L.G. 1974. Legume Foods. In: A.M. Altschul, ed. *New Protein Foods*. Vol I A, p. 231 – 297, New York, Academic Press Inc.

De Vries, S., 2015. Fiber in poultry nutrition: bonus or burden ?. Conference: European Symposium on Poultry Nutrition (24 – 27 August 2015, Prague, Czech Republic, Vol. 20.

Dilger, R.N., Sands, J.S., Ragland, D., Adeola, O. 2004. Digestibility of nitrogen and amino acids in soybean meal with added soyhulls. J. Anim. Sci., 82 (3) : 715 – 724.

D'Mello, J.P.F., Acamovic, T., 1989. *Leucaena leucocephala* in poultry nutrition – a review. Anim. Feed Sci. Technol., 26 (1 – 2) : 1 – 28.

D'Mello, J.P.F., 1991. Nutritional potentialities of fodder trees and fodder shrubs as protein sources in protein nutrition. In: *Legume trees and other fodder trees as protein sources for livestock*. Ed. By Speedy, A. and Pugliese, P.L. Proceedings of FAO Expert Consultants Held at the Malaysian Agricultural Research and Development Institute (MARDI) in Kuala Lumpur, Malaysia, 14 – 18 October 1991.

Gadzirayi, C.T., Masamha, B., Masamha, B., Mupangwa, J.F., Washaya, S., 2012. Performance of broiler chicken fed on mature *Moringa oleifera* leaf meal as a protein supplement to soybean meal. Int. J. Poult. Sci., 11 (1) : 5 – 10.

Gorissen, S.H.M., Crombag, J.J.R., Senden, J.M.G., Waterval, W.A.H., Bierau, J., Verdijk, L.B., Loon, L.J.C.V. 2018. Protein content and amino acid composition of commercially available plant-based protein isolates. Amino Acids, 50 (12) : 1685 – 1695.

Herrera, S.M., Savon, L., Lon-Wo, E., Guiteres, O., Herrera, M., 2019. Inclusion of *Moringa oleifera* leaf meal : its effect on apparent retention of nutrient, productive performance and quality of the carcass of the naked neck fowl. Cuban J. Agric. Sci., 48 (33) : 259 – 264.

Hussain, J., Reddy, P.V.V.S., Reddy, V.R. 1991. Utilisation of leucaena leaf meal by broilers. British Poult. Sci., 32 (1) : 131 – 137.

Ige, A.O., Odunsi, A.A., Akinlade, J.A., Ojedapo, L.O., Ameen, S.A., Adernola, O.A., Rafiu, T.A., 2006. *Gliricidia* leaf meal in layer's diet : effect on performance, nutrient digestibility and economy of production. J. Anim. Vet. Adv., 5 (6) : 483 – 486.

Imoru, A., Onibi, G.E., Osho, I.B., 2019. Performance characteristics of broiler chicken fed graded level of moringa (*Moringa oleifera*) leaf meal in partial replacement of soybean meal. Livest. Res. Rural Dev., 31 (8).

Kumar, A., Kumar, K., Singh, P.K., Paswan, J.K., Chandramoni, 2018. Effect of feeding *Moringa oleifera* leaf meal on production efficiency and carcass characteristics of Vanaraja chicken in tropics. Int. J. Current Microbiol. Appl. Sci., 7 : 1213 – 1220.

Lu, W., Wang, J., Zhang, H.J., Wu, S.G. Qi, G.H., 2016. Evaluation of *Moringa oleifera* leaf in laying hens: effect on laying performance, egg quality, plasma biochemistry and organ histopathological indices. Italian J. Anim. Sci., 15 (4) : 658 – 682.

Machanja, J.U., 2007. Optimising the use of *Gliricidia*, *Calliandra* Leaf and Kenaf Seed Protein in Ruminant Feeding. Masters (Research) Thesis, James Cook University.

Mateos, G.G., Jimenez-Moreno, E., Serrano, M.P., Lazaro, R., 2012. Poultry response to high levels of dietary fiber resource varying in physical and chemical characteristics. J. Appl. Poult. Res., 21 : 156 – 174.

Medugu, C.I., Saleh, B., Igwebuike, J.U., Ndirmbita, R.L., 2012. Strategies to improve the utilization of tannins-rich feed materials by poultry. Int. J. Poult. Sci., 11 (6) : 417 -423.


Mlay, P.S., Pereka, A., Phiri, E.C., Balthazary, S., Igusti, J., Hvelplund, T., Weisbjerg, M.R., Madsen J. 2006. Feed value of selected grasses, legumes and concentrates. Veterinarski Arhiv. 76 (1) : 53 – 63.

Moftakharzadeh, S.A., Janmohammadi, H., Taghizadeh, A., Kianfar, R., Olyayee, M.G., 2019. Effect of enzyme addition on energy utilization and performance of broiler chicken fed wheat-based diet with different metabolizable energy level. Acta Scientiarum Anim. Sci., 41 : 2-10.

Moyo, B., Masika, P.J., Hugo, A., Muchenje, V., 2011. Nutritional characterization of *Moringa*

- (*Moringa oleifera*) leaves. Afr. J. Biotechnol., 10 (60) : 12925 – 12933.
- NAS., 1979. Tropical Legumes : Resources for the Future, National Academy of Sciences, Washington DC., USA., p. 345.
- Ncube, S., Hamudikuwanda, H. Banda, P. 2012. The potential of *Acacia angustissima* leaf meal as a supplementary feed source in broiler finisher diets. Int. J. Poult. Sci., 11 (1) : 55 – 60.
- Ncube, S., Halimani, T.E., Chikosi, E.V.I., Saidi P.T. 2018. Effect of *Acacia angustissima* leaf meal on performance, yield of carcass components and meat quality of broilers. South Afr. J. Anim. Sci., 48 (2) : 271 – 283.
- Ndelekute, E.K., Enyenni, G.E., Akpan, I.P. 2018. Potentials and challenges of utilizing forage sources for chicken production. J. Anim. Sci. Livest. Prod., 12 (1)
- NRC, Nutrient Requirement of Poultry, 1994 Ninth Revised Edition, National Research Council, National Academic Press Washington, D.C.
- Odenyo, A.A., Osuji, P.O., Reed, J.D., Smith, A.H., Mackie, R., McSweeney, C.S., Hanson, J., 2003. *Acacia angustissima* : its anti-nutrients constituents, toxicity and possible mechanism to alleviate the toxicity – a short review. Agroforest. Syst., 59 (2) : 141 – 147.
- Odunsi, A.A., Ogunleke, M.E., Alagbe, O.S., Ajani, T.O., 2002. Effect of feeding *Gliricidia sepium* on the performance and egg quality of layers. Int. J. Poult. Sci., 1 (1,2,3) : 26 – 28.
- Okiki, P.A., Osibote, I.A., Balogun, O., Oyinloye, B.E., Idris, O., Olufunke, A., Asoso, S.O., Olagbemide, P.T. 2015. Evaluation of proximate, minerals, vitamins, and phytochemical composition of *Moringa oleifera* Lam. cultivated in Ado, Ekiti, Nigeria. Adv. Biol. Res., 9 (6) : 436 – 443.
- Oloruntola, O.D., Ayodele, S.O., Agbede, J.O., Asanyan, E.K., 2016. Performance and apparent digestibility of broiler starter fed diets containing *Gliricidia sepium* leaf meal. Asian J. Biol. Life Sci., 5 (1) : 97 – 102.
- Oloruntola, O.D., 2018. *Gliricidia* leaf meal in broiler chicken diet : Effects on performance, carcass, and hemato-biochemical parameters, J. Appl. Sci. Int., 18 (3) : 1 – 9.
- Pamo, E.T., Kana, J.R., Tendonkeng, F., Boukila, B.M., Belfiang, M.E., 2005. Nutritive value and effects of different sources energy on *in vivo* dry matter digestibility of *Leucaena leucocephala* or *Calliandra calothyrsus* in Cameroon. Bull. Anim. Health Prod in Afr. 53 (3) : 149 – 159.
- Paranamana, N., Radampola, K., Bulugahapittya, V.P. 2015. Nutritional and antinutritional contents of alternative feed ingredients for fish feed formulation. J. Anim. Sci., 85 (2) : 212 – 215., 2000. The feeding leaf meal of *Calliandra calothyrsus* to laying hens. Trop. Anim. Health
- Paterson, R.T., Roothaet, R.L., Nyaata, O.Z., Akyeampong, E., Hove L 1996. Experience with *Calliandra calothyrsus* as feed for livestock in Africa. In : Evas. D.O. (Ed). Proceedings of the International Workshop on the genus *Calliandra*, Bogor, Indonesia, 23 – 27 January 1996..
- Paterson, R.T., Roothaert, R.L., Kiruiro, E 2000. The feeding of leaf meal of *Calliandra calothyrsus* to laying hens. Trop. Anim. Health Prod., 32 (1) : 51 - 61.
- Popova, A., Mihaylova, D., 2019. Antinutrients in plant based foods : a review. The Open Biotechnol. J., 13 (1) : 68 – 76.
- Prasad, D.A., Reddy, K.I., Reddy, C.S. 1970. Agathi leaf meal (*Sesbania grandiflora*) in the rations of growing chickens. Indian Vet. J., 47 : 164 -169.
- Reddy, D.V. and Elanchezian, N. 2008. Evaluation of tropical tree leaves as ruminant feedstuffs based on cell contents, cell wall fractions and polyphenolic compounds. Livest. Res. Rural Dev., 20 (5).
- Rusdy, M., 2016. *Calliandra calothyrsus* leaves as tanniferous fodder for ruminant animals. Int. J. Sci. Res. Publ., 6 (12) : 286 – 290.

- Salawu, M.B., Acamovic, T., Stewart, C.S. Maasdorp, B. 1997. Assessment of nutritive value of *Calliandra calothyrsus*: its chemical composition, and the influence of tannins, piperolic acid and polyethylene glycol on *in vitro* organic matter digestibility. Anim. Feed Sci. Technol., 69 (1 – 3) : 207 – 217.
- Sarvade, S. Parmar, Y.S., 2014. *Sesbania grandiflora* (L.) Poiret: A potential agroforestry tree species. Popular Kheti, 2 (3) : 204 – 207.
- Shqueir, A.A., Brown, D.L., Klasing, K.C., 1989. Canavanine content and toxicity of *Sesbania* leaf meal for growing chicks. Anim. Feed Sci. Technol., 1 – 2 : 137 – 147.
- Swennen, Q., Everaert, N., Debonne, M., Verbaeys, I., Careghi, C., Tona, K., Janssen, G.P.J., Decuypere, E., Bruggeman, V., Buyse, J. 2010. Effect of macronutrient ratio of the prestarter diet on broiler performance and intermediary metabolism. J. Anim. Physiol. Anim. Nutr., 94 :375 – 384.
- Tantung, R.L., Madamba, L.S.P. 1981. Fractination and characterization of Ipil-ipil (*Leucaena leucocephala* (Lam) de Wit seed proteins. The Bull. Philippine Biochem. Soc., 4 (1 & 2) :15 – 32.
- Tesfaye, E., Anmut, G., Urge, M., Dessie, T., 2013. *Moringa oleifera* leaf meal as an alternative feed ingredients in broiler ration. Int. J. Poult. Sci., 12 (5) : 289 – 297.
- Tsega, W., Tamir, B. 2009. The effect of increasing levels of dried leaves of sweet potato (*Ipomoea batatas*) on dry matter intake and body weight gain performance of broiler finisher chicken. Livest. Res. Rural Dev., 21 (12).
- Ugo, N.J., Ade, A.R., Joy, A.T., 2019. Nutrient composition of *Carica papaya* leaf extract. J. Food Sci. Nutr. Res., 2 (3) : 274 – 282.
- Vries, S.de., 2015. Fiber in poultry nutrition, bonus or burden ?. 20th European Symposium on Poultry Nutrition (24 – 27 August, 2015), Prague, Czech Republic.
- Wina, E., Pasaribu, T., Rakhmani, S.I.W., Tangendjaja, B., 2017. The role of saponin as feed additive for sustainable poultry production. Wartazoa, 27 (3) : 117 – 124).
- Yousuf, M.B., Belewu, M.A, Daramola, J.O., O Gundun, N.I., 2007. Protein supplementation value of cassava, leucaena and gliricidia leaf meal in goats fed low quality *Panicum maximum* hay. Livest. Res. Rural Devel., 19 (2)..
- Zhou, Y., Jiang, Z., Wang, T. 2009. Improved energy-utilizing enzyme by enzyme preparation supplement in broiler with different metabolizable energy levels. Poult. Sci., 88 (2) : 316 – 322.



A Review on Selected Tropical Tree Legume Leaves in Poultry Diet

ORIGINALITY REPORT

6%

SIMILARITY INDEX

%

INTERNET SOURCES

%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|----|
| 1 | Ben Salem, H.. "Spineless cactus (<i>Opuntia ficus indica</i> f. <i>inermis</i>) and oldman saltbush (<i>Atriplex nummularia</i> L.) as alternative supplements for growing Barbarine lambs given straw-based diets", <i>Small Ruminant Research</i> , 200401 Publication | 1% |
| 2 | www.abah.bioflux.com.ro Internet Source | 1% |
| 3 | hal.archives-ouvertes.fr Internet Source | 1% |
| 4 | periodicos.uem.br Internet Source | 1% |
| 5 | www.scribd.com Internet Source | 1% |

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On