

# Health workers

*by* Hasnah Hasnah

---

**Submission date:** 03-Sep-2021 10:06AM (UTC+0700)

**Submission ID:** 1640490529

**File name:** Manskrip\_OK\_Overview.....\_Hasnah.doc (121.5K)

**Word count:** 4679

**Character count:** 25886

# Overview of Health Workers' Knowledge of Pre-Eclampsia in Gowa Regency, Sulawesi Selatan, Indonesia: Cross-sectional Study

Hasnah Nur<sup>1, 2</sup>, Muh Syafar<sup>3</sup>, Ansariadi<sup>4</sup>, Ummuh Salmah<sup>5</sup>, Ratnasari<sup>2</sup> and Astuti<sup>2</sup>

<sup>1</sup>Doctoral Program at Hasanuddin University, Makassar, Indonesia

<sup>2</sup>Department of Maternity, Nursing Prody, Faculty of Medicine and Health Sciences, Universitas Islam Negeri Alauddin, Makassar, Indonesia

<sup>3</sup> Department of Health Promotion, Hasanuddin University, Makassar, Indonesia

<sup>4</sup> Department of Epidemiology Hasanuddin University, Makassar, Indonesia

<sup>5</sup> Department of Biostatistics Hasanuddin University, Makassar, Indonesia

<sup>1</sup>Correspondence Email: hasnah.nur@uin-alauddin.ac.id

**Abstract.** Saharan has higher mortality rates than in Switzerland. Likewise in Indonesia, maternal mortality rates vary from one source to another. The maternal mortality rate has decreased every decade but is still far from the 2015 MDGs-3 target. The cause of maternal death has also changed from bleeding to hypertension in pregnancy, followed by infection. The highest maternal mortality in south Sulawesi occurred in Gowa Regency, then Bone and Jeneponto Regencies.

Objective: describe the level of knowledge of health workers about the prevention of pre-eclampsia

Method: Cross sectional study design conducted at 3 Primary health care Sombaopu, Samata, and Bajeng, from April to June 2019. A sample size of 29 people were nurses, midwives, and doctors in the antenatal care room. The research instrument used a questionnaire of 25 questions about understanding, causes, symptoms, and prevention of pre-eclampsia. Reliability test with Cronbach Alpha value of 0.756 > 0.600 so that the instrument was declared reliable. Data analysis uses frequency distribution. Research ethic from the ethics committee of the Faculty of Medicine and Health Sciences Universitas Islam Negeri Alauddin Makassar Number A.110 / KEPK / FKIK / V / 2019 with No. Registrar UINAM19050110

Conclusion: Knowledge of health workers both in terms of understanding, signs, and symptoms, and prevention in the good category unless the cause is in the sufficient category.

## 1. Introduction:

Maternal mortality rates in various countries experienced differences between countries such as Switzerland, maternal deaths by 73 percent (Say et al., 2014), compared to countries in Sub-Saharan Africa and Asia, maternal deaths were higher by 90 percent (Barnes Josiah, Myntti, and Augustin, 1998). The cause of maternal death occurred due to direct obstetric causes, namely bleeding by 27.1 percent. (Gil González, Carrasco Portiño, and Ruiz, 2006) in African countries say that the cause of maternal death is postpartum hemorrhage (PP), followed by eclampsia and pre-eclampsia, delayed labor, infection, and indirect causes. Two-thirds of deaths due to bleeding occur in the postpartum period.

<sup>2</sup> Maternal death or maternal death is the death of a mother during pregnancy or within 42 days after the end of pregnancy, regardless of the place or age of the pregnancy. In 2008 the WHO (World Health Organization) <sup>6</sup> concluded that each year more than 500,000 women who die have given birth, one of the causes of maternal and fetal mortality and morbidity is preeclampsia (PE), the incidence rate ranges from 0.51% to 38.4 %. In developing countries, the maternal mortality rate due to preeclampsia and eclampsia is still high, while in developed countries the incidence of preeclampsia is around 6 to 7% and eclampsia 0.1 to 0.7% (Situmorang et al, 2016). This is because in developed countries better prenatal care. The incidence of preeclampsia is influenced <sup>14</sup> several factors such as race, parity, genetic and environmental factors. Maternal Mortality Rate (MMR) in Indonesia is very high compared to MMR in various countries in Southeast Asia (Purwoastuti, 2015). AKI is included in the 2015 Sustainable Development Goals (SDG's) achievement target of reducing AKI in 2019 to 306 / 100,000 live births (DIY Health Office, 2016).

<sup>2</sup> Maternal death or maternal death is the death of a mother during pregnancy or within 42 days after the end of pregnancy, regardless of the place or age of the pregnancy. In 2008 Indonesia was a developing country with a high maternal mortality rate (MMR) and perinatal mortality, the third-highest in ASEAN and the second highest in the South East Asian Nation Regional Organization (WHO, 2013). The Indonesia Demographic and Health Survey (IDHS) in 2018 showed that pregnant women with anemia increased from 2013 wherein 2013 with a presentation of 37.1 and in 2018 a presentation of 48.9 was obtained. The proportion of antenatal care (ANC access) for women aged 10 to 54 years according to the province of South Sulawesi is an increase in the presentation of visits of pregnant women in 2013 with 96.1 presentations and in 2018 96.3 (IDHS, 2018).

<sup>11</sup> The maternal mortality rate (MMR) in Indonesia is a benchmark of the quality of midwifery services and an important indicator of the degree of public health. Factors causing maternal mortality include puerperal hemorrhage of about 26.9%, eclampsia during childbirth 24%, bleeding 28%, infection 11%, puerperium complications 8%, obstetric trauma 5%, obstetric embolism 3%, abortion 5% and others 11% (Health Departement RI, 2015).

<sup>1</sup> Mortality and Morbidity in pregnant and childbirth women are a big problem. "Based on data from the Health Office 2016-2017, there are 11 regencies with high maternal and infant mortality rates, namely Gowa ranked first, Bone second, Jeneponto third and in fourth place are Bulukumba, Pangkep, Luwu. Then in fifth, there is Sinjai, Maros, Toraja, while North Luwu and Makassar were ranked sixth. "The cause of maternal death in South Sulawesi was caused by bleeding in the majority of 40 cases

and hypertension in pregnancy 35 cases. Over these conditions the importance of early detection at the time of Antenatal Care and appropriate and appropriate handling of standards can prevent both conditions (Health Profil Sulawesi selatan, 2016).

<sup>18</sup> Preeclampsia is one of the causes of maternal death worldwide, so the need for early detection and treatment that plays a role in reducing maternal mortality. Although there is no proven effective method to prevent the occurrence of preeclampsia, detection is needed to anticipate the event, so providing care antenatal care, management and appropriate treatment can be done as early as possible (O'Gorman et al., 2016).

<sup>1</sup> Based on the Regulation of the Minister of Health of the Republic of Indonesia Number 97 regarding Health Services, integrated antenatal which is a comprehensive and quality health service, things that must be done are among others by checking early detection of problems, illnesses, and complications / complications of pregnancy and carrying out case management in a timely manner and quick referrals if needed. So that the role of health workers, especially midwives is needed in the early detection of complications in pregnancy (Minister of Health Regulation RI No. 97, 2014) because one of the causes of AKI and IMR indirectly is preeclampsia. Preeclampsia is a hypertensive syndrome in certain pregnancies with multi-system conditions and multifactorial causes for humans, which is significantly related to maternal and perinatal mortality and morbidity.

<sup>1</sup> In reducing the morbidity and mortality rate of mothers and infants, efforts can be made, namely early detection of signs and hazards in pregnancy. It is important to pay attention to clinical signs during antenatal care by monitoring the increase in blood pressure, proteinuria, weight gain during pregnancy and nutritional status and avoid factors that are at risk for the occurrence of preeclampsia in pregnancy. The latest guidelines from the National Institute for Health and Clinical Excellence (NICE) recommend routine screening of specific risk factors for preeclampsia (nulliparity, older age, body mass index, family history of preeclampsia, pregnancy intervals of more than 10 years, history of kidney disease or hypertension chronic, multiple pregnancy, and a history of preeclampsia in a previous pregnancy). Difficulties that are often faced by pregnant women, namely being late in making decisions, resulting in late getting treatment. These complications can be immediately handled appropriately by health workers. So expect no maternal deaths due to preeclampsia (Situmorang et al, 2016).

Efforts to prevent perinatal mortality can be reduced if factors that have a predictive value of the incidence of preeclampsia can be identified. According to the results of the 2014 Artathi research on the determinant analysis affecting village midwives in the accuracy of referrals in cases of preeclampsia / eclampsia in Banyumas District, the study found no significant relationship between knowledge and accuracy in referral of village midwives to cases of preeclampsia / eclampsia.

<sup>2</sup> Based on the Health Profile of Gowa Regency in 2016, the highest number of maternal deaths was the Sombaopu Community Health Center in 2016 there were 4 mothers who died including 2 mothers who died during childbirth and 2 people during the postpartum period. And Samata health center found 2 mothers who died during

childbirth (Gowa Health Office, 2016). So that <sup>14</sup> research is needed to determine the knowledge of health workers for pre-eclampsia.

## 2. Methode

The research method used is quantitative. The design used is descriptive research design. Descriptive research is a study in which there is no analysis of the relationship between variables, there are no independent and dependent variables. Where this study aims to obtain information about the level of knowledge of health workers to early detection of preeclampsia. This research was conducted in the working area of Sombaopu Community Health Center, Samata Community Health Center, and Bajeng Health Center in April and June 2019.

<sup>4</sup> The population in this study were all health workers in the working area of primary health care Samata, Sombaopu and Bajeng. Samples are health workers who work in the ANC room and are willing to be respondents using the sensus sampling method.

Data collection tools used were questionnaires. The questionnaire used was a questionnaire consisting of 25 questions. Questions about understanding 4 questions, questions about causes 9, questions about signs and symptoms 5, questions about prevention 7 that have been tested for reliability

Data processing is done manually by filling out the observation sheet provided. The data processing is then processed using the SPSS program with Editing, Coding, Data tabulation and data analysis stages. Data analysis contains an explanation of the data on each variable studied which is then described. This study uses a way to analyze data, namely univariate data analysis. Univariate analysis is the process of analyzing each research variable that exists descriptively by calculating the attributional frequency and presentation of each variable (Notoatmodjo, 2005).

## 3. Result

This study of pre-eclampsia health professionals knowledge in the primary health care Sombaopu, Bajeng and Samata in Gowa District has been conducted since May and June 2019. Respondents in this study were respondents aged 20 and 50 years in the MCH space respondents 51 people with 37 midwife professions, 11 profession nurses, and 3 doctors.

### 1.1 Characteristics of Respondents

**Table 1.** A Distribution of Frequency of Respondents by Age, Profession, Education, and years of worker in Health Workers

Characteristics	Frequency	Presentation
Age		
21-30 year	34	66.6
31-40 year	5	9.8
41-50 year	12	23.5
Profesi		
Nurse	11	21.5
Midwifery	37	72.5
Doctor	3	5.8
Education		

D3	30	58.8
D4	14	27.4
Bachelor	7	13.7
<b>Years of worker</b>		
1 moon– 1.9 year	8	15.6
2 - 10 year	33	33
10.1 - 20 year	7	13.7
20.1 - 30 year	3	5.8
<b>Total</b>	<b>51</b>	<b>100</b>

10  
1.2 *Level of knowledge of health workers about preeclampsia in primery health care at Gowa District.*

10  
**Table 2.** A Distribution of Frequency *Level of knowledge of health workers* about preeclampsia in primery health care at Gowa District.

<b>Characteristics</b>	<b>Frequency (person)</b>	<b>Presentation (%)</b>
Knowledge about preeclampsia(all aspects)		
Well	38	74.5
Enough	13	25.4
Less	0	0
<b>Total</b>	<b>51</b>	<b>100</b>

1  
Table 2. shows that the majority of health workers have a level of knowledge about preeclampsia (all aspects) that fall into the category of quite as many as 13 person (24.4%) and health workers have a level of knowledge about preeclampsia (all aspects) that falls into the good category of 38 person (74.5%)

4  
1.3 *The level of knowledge of health workers about the meaning of preeclampsia in primery health care at the Gowa District*

**Table 3.** A Distribution of Frequency Level of the meaning of preeclampsia in primery health care at Gowa District.

<b>Characteristics</b>	<b>Frequency (person)</b>	<b>Presentation (%)</b>
Knowledge about preeclampsia(all aspects)		
Well	39	74.4
Enough	12	23.5
Less	0	0
<b>Total</b>	<b>51</b>	<b>100</b>

Table 3. Shows that based on the aspect of understanding of preeclampsia, most health workers have a sufficient level of knowledge of 12 person (23.5%) and health workers with a good level of knowledge found in 39 person (74.4%).

4  
1.4 *The level of knowledge of health workers about the causes of preeclampsia in Primary Health Care at Gowa District*

**Table 4.** A Distribution of Frequency Level cause of preeclampsia in primery health care at Gowa District.

Characteristics	Frequency (person)	Presentation (%)
Knowlodge about preeclampsia(all aspects)		
Well	19	37.274.5
Enough	21	41.1
Less	11	21.5
<b>Total</b>	<b>51</b>	<b>100</b>

Table 4 shows that based on the causes of preeclampsia, most health professionals have a sufficient level of knowledge of 21 people (41.1%) and health workers who have less knowledge level of 11 people (21.5%), and health workers who have a good level of knowledge with the number the smallest were 19 person (37.2%).

*1.5 The level of knowledge of health workers about the signs and symptoms of preeclampsia in Primary Health Care at Gowa District*

**Table 5.** A Distribution of Frequency Level of knowledge of health workers about preeclampsia in primery health care at Gowa District.

Characteristics	Frequency (person)	Presentation (%)
Sign and symptom preeclampsia(all aspects)		
Well	43	84.3
Enough	6	11.7
Less	2	3.9
<b>Total</b>	<b>51</b>	<b>100</b>

Table 4 shows that based on the signs and symptoms of preeclampsia some health workers have a good level of knowledge as many as 43 people (84.3%) and vice versa health workers who have less knowledge level are found in the smallest number of 2 people (3.9%) and health workers who have levels sufficient knowledge of 6 people (11.7%).

*1.6 The level of knowledge of health workers about the prevention of preeclampsia in the Primay Health Care Gowa District*

**Table 6.** Distribution Frequency The level of knowledge of health workers about the prevention of preeclampsia in the Primay Health Care Gowa District



Characteristics	Frequency (person)	Presentation (%)
Sign and symptom preeclampsia(all aspects)		
Well	49	96.01
Enough	2	3.9
<b>Total</b>	<b>51</b>	<b>100</b>

Table 6 shows that based on the prevention of preeclampsia, most health workers have a good level of knowledge about prevention as many as 49 person (96.01%) and vice versa health workers with a sufficient level of knowledge of 2 person (3.9%).

#### 4. Discussion

##### 4.1 Knowledge of health workers about preeclampsia

##### 4.1.1 Knowledge of health workers about definition of preeclampsia

Based on the aspect of understanding of preeclampsia, most health workers have a sufficient level of knowledge of 12 people (23.5%) and health workers with a good level of knowledge are found in 39 people (74.4%). This condition indicates that most health workers have good knowledge about preeclampsia.

Knowledge is a very important domain for the formation of open behavior (over behavior) which is the result of tofu that occurs through sensory processes, especially the eyes and ears of certain objects. Knowledge (knowledge) is also interpreted as a result of human sensing or the result of knowing someone about an object through their senses (eyes, nose and so on), by itself at the time of sensing so as to produce knowledge. Knowledge is an introduction to an object or thing objectively. Knowledge is an activity developed through the learning process and stored in memory will be dug up when it will be needed through the form of memory. Knowledge is an important domain factor for the formation of one's actions (Perdania, 2013).

From this, it is expected that health workers show a high curiosity about everything. But curiosity is not the main factor that influences the level of one's knowledge, but there are still other factors namely: the level of education, experience, information, culture.

The majority of health workers have a good level of knowledge found in 39 people (74.4%). so it can be concluded that the behavior is influenced by a person's experience, factors outside the person (environment), both physical and non-physical and socio-cultural, then the experience is known, perceived, believed, giving rise to motivation, intention to act and eventually becomes behavior. And look at the length of service with a span of 2-10 years at most worth 34 people or 66.6%. to be able to make an appropriate referral to preeclampsia / eclampsia cases depends on the experience and results of the analysis of health workers in handling these cases. Although in theory the midwife is able to understand, but if in terms



of experience and external factors are less supportive, it can have an impact on the inaccuracy in referencing pre-eclamptic / eclamptic cases

In making decisions for the follow-up to preeclampsia, an understanding is needed, especially regarding the understanding of preeclampsia, while in seeing a picture of the level of knowledge of health workers about preeclampsia is still obtained with a sufficient level of knowledge of 12 people (23.5%).

4

#### 4.1.2 Knowledge of health workers about the causes of preeclampsia

Knowledge of health workers about the causes of preeclampsia is still lacking. Factors that can increase the incidence of preeclampsia are first-time pregnancy (primigravida), pregnancy with a higher uterine strain such as pregnancy with most amniotic fluid, the incidence of preeclampsia will be higher in multiple pregnancy and pregnancy with a large fetus, the presence of maternal disease that accompanies pregnancy (kidney disease , high blood pressure) (Manuaba, 2016).

Preeclampsia is due to a complex immunologic mechanism and blood flow to the placenta is reduced. As a result, the supply of nutrients needed by the fetus is reduced. So that, if it lasts longer the incidence of preeclampsia will get worse and can inhibit fetal growth. Preeclampsia can cause harm to the mother and fetus. The symptoms are swelling in some parts of the body, especially the face and hands. And it will be more dangerous if accompanied by a sudden increase in blood pressure and high protein levels in the urine (Indiarti, 2010).

Preeclampsia must be treated immediately, because it can cause a substance that can destroy endothelial cells lining the blood vessels. If not treated quickly and correctly it will continue to become eclampsia characterized by seizures, even to the point of coma. This condition is very dangerous for pregnant women and the fetus, because if not treated immediately there will be permanent damage to the nerves, blood vessels or the mother's kidney. Meanwhile, babies will experience mental retardation due to lack of blood flow through the placenta and oxygen in the brain (Indiarti, 2015).

So seeing the theory of knowledge of health workers about preeclampsia is very important especially the causes of preeclampsia so that after knowing the cause of preeclampsia health workers are able to conduct early detection so as to reduce the incidence of preeclampsia in pregnant women, because according to manuaba almost 50% of maternal and fetal deaths are caused by preeclampsia therefore it is important to know the cause of preeclampsia as early as possible so that treatment can be carried out immediately. Based on this explanation, it is concluded that knowledge about early detection of causes of preeclampsia is very important for the lives of mothers and fetuses so that information and socialization regarding harmful causes are needed especially pregnant women in order to detect them as early as possible.

#### *4.1.3 Knowledge of health workers about the signs and symptoms of preeclampsia*

The results of the study showed that the majority of health workers had a high level of knowledge about the signs and symptoms of preeclampsia. Knowledge of health workers about preeclampsia is very important especially knowing as early as possible the signs and symptoms of preeclampsia so that after knowing or detecting as early as possible from preeclampsia health workers are able to make appropriate further handling in helping pregnant women, because according to manuaba maternal and fetal deaths are almost 50% caused by preeclampsia

Preeclampsia must be treated immediately, because it can cause a substance that can destroy endothelial cells lining the blood vessels. If not treated quickly and correctly, it will continue to become eclampsia, which is characterized by seizures, even to coma. This condition is very dangerous for pregnant women and the fetus, because if not treated immediately there will be permanent damage to the nerves, blood vessels or the mother's kidney. Meanwhile, babies will experience mental retardation due to lack of blood flow through the placenta and oxygen in the brain (Indiarti, 2015).

#### *4.1.4 Knowledge of health workers about the prevention of preeclampsia*

The results showed that most health workers had good knowledge about the prevention of preeclampsia. Various groups of people in Indonesia, especially mothers who consider pregnancy as kodart, natural, and ordinary. They felt they did not need to check themselves regularly to the health service. There are still many mothers who are less aware of the importance of antenatal care, so that no risk factors can be detected, including preeclampsia. This risk is only known at the time of delivery which is often difficult to handle so that the fatal result is death. This is caused by the low level of knowledge and lack of information about preeclampsia, including regarding the prevention aspects of preeclampsia (Rosliana, 2015).

In this case after knowing the level of knowledge of health workers about the prevention of preeclampsia, it is necessary to make efforts to provide counseling and education to pregnant women to reduce the incidence of preeclampsia, including ANC examination, classes of pregnant women, posyandu, and pregnancy exercises is an effort that can be done pregnant women is a prevention to reduce the incidence of preeclampsia (Indrawati, 2015).

Antenatal examinations are carried out regularly, thoroughly and quality, can recognize the signs of preeclampsia as early as possible and provide adequate treatment so that the disease does not become more severe, need to be aware of the possibility of preeclampsia, provide an explanation of the prevention of preeclampsia such as rest and the importance of

regulating diabetes low salt and maintain excessive weight gain (Mochtar, 2011).

#### 4.1.5 *Knowledge health works with early detection preeclampsia*

Preeclampsia is one of the main causes of perinatal morbidity and mortality in Indonesia. Even today, preeclampsia is still a health problem that cannot be completely resolved.

Risk factors for preeclampsia include knowledge, occupation, antenatal examination, and in reducing the Perinatal Death Rate (PPA) caused by preeclampsia. One of the efforts that can be done is to reduce the incidence of preeclampsia. Incidence rates can be reduced by efforts including prevention, early observation, and therapy. If the identified factors can be predicted then efforts to prevent perinatal mortality can be reduced.

Based on these data shows that the knowledge of health workers is mostly included in good knowledge. Knowledge is influenced by several factors including age, education, mass media / information, socio-cultural and economic, environment and experience (Notoatmojo, 2008).

According to the results of the 2014 Artathi research on the determinants of analysis affecting village midwives in the accuracy of referrals in cases of preeclampsia / eclampsia in Banyumas district showed no significant relationship between knowledge and accuracy in referring village midwives to cases of pre-eclampsia / eclampsia. Behavior is influenced by one's experience, factors outside the person (the environment), both physical and non-physical and socio-cultural which then experience is known, perceived, believed to cause motivation, intention to act and eventually become behavior.

So that researchers assume that knowledge is very important for our lives, and knowledge about health and health issues, especially the problem of preeclampsia because preeclampsia can affect the mother and fetus, but in correctly detecting the incidence of preeclampsia not only with good knowledge but also depends on the results of analysis and experience health workers in handling the case. Although in theory health workers are able to understand, but in terms of experience and external factors that are less supportive, it will have an impact on the inaccuracy in referring preeclampsia / eclampsia cases.

Based on four items of questions about the knowledge of health workers about preeclampsia, it can be seen that the level of knowledge of health workers regarding the causes of preeclampsia is still lacking. While early detection for the incidence of preeclampsia is very important in order to reduce the incidence of preeclampsia.

According to Manuaba in 2016 which can increase the incidence of preeclampsia there are several factors including first pregnancy (primigravida), pregnancy with a higher uterine strain such as pregnancy with mostly amniotic water, multiple pregnancy and pregnancy with a

large fetus, the incidence will be higher in the presence of maternal disease accompanying pregnancy (kidney disease, high blood pressure).

## 5. Research Ethics

In conducting this research, the researcher obtained recommendations from the institution for other parties by submitting an application for permission to the institution, emphasizing ethical issues that included informed consent, confidentiality, and anonymity. This study has passed the code of ethics from the health research ethics committee of the Faculty of Medicine and Health Sciences of UIN Alauddin Makassar Number A.110 / KEPK / FKIK / V / 2019 with Registration Number UINAM19050110.

## 6. Conclusion

Knowledge of health workers based on aspects of the understanding of preeclampsia, most have a picture of a good level of knowledge; Based on the cause of preeclampsia, most health workers have a sufficient level of knowledge; Knowledge of health workers based on signs and symptoms of preeclampsia some health workers who have a good level of knowledge Knowledge of health workers based on the prevention of preeclampsia, most health workers who have a good level of knowledge about prevention; The level of knowledge of health workers about preeclampsia (all aspects) falls into either category.

## REFERENCE

- [1] Artathi, Eka. 2014. Determinant Analysis that Affects Village Midwives in Referral Accuracy in Preeclampsia / Eclampsia Cases in Banumas District. YLPP Midwifery Academy Purwokerto. Imiah Journal Vol.5 NO.2
- [2] Barnes-josiah, D., Myntti, C., & Augustin, A. 1998. the " Three Delays " As a Framework for Examining Maternal Mortality in Haiti. Soc, Sci Med, 46 (8).
- [3] Central Statistics Agency, 2018. Indonesian Demographic and Health Survey (IDHS) 2018. Jakarta: Central Statistics Agency.
- [4] DIY Health Service. 2016. Annual Report of Yogyakarta Special Region Health Office.
- [5] Gil-González, D., Carrasco-Portiño, M., & Ruiz, M. T. 2006. Knowledge gaps in scientific literature on maternal mortality: a systematic review. Public Health Reviews, 84 (11), 903–909. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2627556/pdf/17143465.pdf>
- [6] Gowa district health office. 2016. Health Profile of Gowa Regency. Sungguminasa, 2016
- [7] Indiarti, 2015. Relationship between Age and Preeclampsia with Postpartum Hemorrhage Occurrence in Bantul Senopati Hospital General Hospital, Doctoral Dissertation. Yogyakarta: Aisha's Stikes.

- [8]Indrawati, 2015. Midwife level of knowledge about preeclampsia at the Husin Family Clinic in Medan in 2014. Volume 16 No. June 1, 2015.
- [9]Manuaba, Ida Bagus Gede. 2015. Introduction to the Obstetrics Lecture. Jakarta: ECG
- [10]Mochtar, 2011. Reference Book for Maternal and Neonatal Health Services. Yogyakarta: Bina Pustaka Sarwono Prawirohardjo Foundation.
- [11]Notoatmodjo, S. 2012. Health Promotion and Health Behavior. (Jakarta: Rineka Cipta)
- [12]Notoatmodjo, 2005. Health Research Methodology. Jakarta: Rineka Cipta.
- [13]Notoatmodjo, 2008. Health Education and Behavior. (Jakarta: PT Rineka Create)
- [14]Manuaba 2010. Obstetrics, Gynecology and Family Planning. Jakarta.
- [15]O'Gorman, Neil., Et al. 2016. Competing risks model in screening for preeclampsia by maternal factors and biomarkers at 11-13 weeks gestation. American Journal of Obstetrics & Gynecology, 214: 103.e1-12., [Http://dx.doi.org/10.1016/j.ajog.2015.08.034](http://dx.doi.org/10.1016/j.ajog.2015.08.034).
- [16]Permenkes, 2014. Pre-Pregnancy, Pregnancy, Childbirth, and Childbirth Services, Contraception Services, and Sexual Health Services. Jakarta: Regulation of the Minister of Health RI No. 97, 2014.
- [17]Perdania, Nurul. 2013. Level of Knowledge of Pregnant Women about Preeclampsia in Brujul Village, Central Java, Karanganyar in 2013. Stikeskusumahusada.ac.id. Accessed 31 July 2013.
- [18]Rosliana. 2015. Description of Pregnant Women Knowledge about Preeclampsia in Banguntapan II Health Center, Bantul. KTI D3 midwifery Stikes General Achmad Yani Yogyakarta.
- [19]Say L, Chou D, Gemmill A, et al. 2014. Global causes of maternal death: A WHO systematic analysis. Lancet Glob Heal. 2014; 2 (6): 323-333. doi: 10.1016 / S2214-109X (14) 70227-X
- [20]Situmorang, T.H, et al. 2016. Factors Associated with the Incidence of Preeclampsia in Pregnant Women at the KiaRsu Poly Anutapura Palu. Tadulako Health Journal 2 (1): 1- 75
- [21]The Indonesian Ministry of Health, 2015. Indonesian Health Profile. Jakarta: Indonesian Ministry of Health in 2015.
- [22]World Health Organization (WHO). 2013. A Global Brief Hypertension: Silent Killer, Global Public Health Crises.
- [23]World Health Organization (WHO). 2008. Maternal Mortality.

# Health workers

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

8%

INTERNET SOURCES

12%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1

"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020

Publication

4%

2

journal.unpak.ac.id

Internet Source

2%

3

Hasnah Nur, Amiruddin Ridwan, Triratnawati Atik, Muhammad Lalu S, Ar. Megawahyuni, Wahyuni, Wahdaniyah, Eny Sutria. "Makassar ethnic community perception about pre-eclampsia: A study phenomenology", Enfermería Clínica, 2020

Publication

1%

4

Siti Zahara Nasution, Bella Sastia, Cholina Trisa Siregar, Reni Asmara Ariga, Lufthiani, Muhammad Ridha Haykal Amal. "Knowledge, attitudes, and efforts of factory workers against the dangers of ISPA at Pt Asia karet medan", AIP Publishing, 2021

Publication

1%

5

jnk.phb.ac.id

Internet Source

1%



6	<a href="http://proceeding.unived.ac.id">proceeding.unived.ac.id</a> Internet Source	1 %
7	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	1 %
8	<a href="http://archive.org">archive.org</a> Internet Source	1 %
9	Submitted to iGroup Student Paper	1 %
10	Sunil Sapkota, Deb P. Pandey, Guru P. Dhakal, Dhan B. Gurung. "Knowledge of health workers on snakes and snakebite management and treatment seeking behavior of snakebite victims in Bhutan", PLOS Neglected Tropical Diseases, 2020 Publication	1 %
11	Tetty Ripursari. "Competence (Knowledge, Attitudes and Skills) Midwifeve to Handling of Obstetric Emergency According to Standard Operational Procedures", Journal for Quality in Public Health, 2019 Publication	1 %
12	Yuli Mulyanti Soenar, Deswani Kasim. "Reproductive Health Education in the Elderly", Asian Journal of Applied Sciences, 2019 Publication	1 %

13	Submitted to Pennsylvania State System of Higher Education Student Paper	1 %
14	elibrary.almaata.ac.id Internet Source	1 %
15	www.karger.com Internet Source	1 %
16	www.bmj.com Internet Source	1 %
17	digitalcommons.ric.edu Internet Source	1 %
18	www.neliti.com Internet Source	1 %

Exclude quotes    On  
Exclude bibliography    Off

Exclude matches    < 1%