Biodiversitas Journal of Biological Diversit	ty Tasks	0	Ə English 💿 Vie	w Site 🔒 mas	huribiodiversit
OPEN JOURNAL SYSTEMS	) / Masri et a	al. $/$ Novel chitinolytic bacteria from the shrimp shell processing waste			Library
Submissions	Workflow	Publication			
	Submission	Review Copyediting Production			
	Submissio	on Files		Q Sea	rch
	▶ 🕅 382	33-1 mashuribiodiversitas, MASHURI MASRI BIODIVERSITAS.doc	February 19, 2021	Article Text	
	► 🕅 384	35-1 aseptiasari, Novel chitinolytic bacteria from shrimp.doc	February 23, 2021	Article Text	
				Download All F	iles
	Pre-Revie	w Discussions		Add discuss	ion
	Name	From	Last Reply	Replies (	Closed
	<u>Submission</u>	aseptiasari 2021-02-20 05:20 AM	mashuribiodiversitas 2021-02-22 08:31 AM	; 1	

Platform & workflow by OJS / PKP

Submission	×	
Participants		Libra
- Anisa Septiasari (aseptiasari)		
Mashuri Masri (mashuribiodiversitas)		
Messages		
Note	From	
Dear Sir,	aseptiasari	Q Search
Thank you very much for your manuscript submission.	2021-02-20 05:20 AM	Text
Unfortunately, this mss has outdated references to be published in Biodiversitas Journal. At		
least, you need to compose 20 citations which:		Text
<ol> <li>80% of international scientific journals published in the last 10 years, and also 10% of national scientific journals published</li> </ol>		
2. Please write references based on author's guidelines.		oad All Files
Kindly check and correct accordingly.		
<ul> <li>Dear Mrs.Anisa and Mr Ahmad setyawan</li> </ul>	mashuribiodiversitas	
Thank you for your fast reply	AM	discussion
		lies Closed
I have checkhed and correct based on your instruction.		
Right now I have 59 reference :		
$80\% \times 59 = 47,2$ . I have 48 international scientific journals published in the last 10 years		
$10\% \times 59 = 5.9.$ I have 7 national scientific journals published.		
I have only 4 international scientific journals published in the more 10 last years		
And we have written references based on author's guidelines		
Thank you very much for your collaboration.		
Best Regards		
Mashuri Masri		
Note	From	

umal of Biological Dive	rsity Tasks 🕕		🛛 English 🛛 👁 Vie	w Site 🔺 n	nashuribio
L SYSTEMS	019 / <b>Masri et al.</b> / Novel chitinolytic bacteria from t	the shrimp shell processing waste	2		Lib
	Workflow Publication				
	Submission Review Copyediting P	roduction			
_	Round 1				
	<b>Round 1 Status</b> Submission accepted.				
	Notifications				
	[biodiv] Editor Decision			2021-03-18 02	2:35 AM
	Reviewer's Attachments			Q s	earch
	40806-1 , REVIEW TRI DEWI(1).pdf		April 15, 2021		
	40807-1 , REVIEW BU EVIKA-V-Novel chiting	olytic bacteria from shrimp (1) (2)	.doc April 15, 2021		
	Revisions		Q Sea	rch Uploa	d File
	40808-1 Article Text, FINAL ARTICLE.doc		April 15, 2021	Article Text	
	40809-1 Other, PROFESSIONAL LANGUAG	GE EDITOR.pdf	April 15, 2021	Other	
	Review Discussions			Add discu	ission
	Name	From	Last Reply	Replies	Closed
	Uncorrected Proof	dewinurpratiwi 2021-04-13 09:28 PM	mashuribiodiversitas 2021-04-15 08:22 AM	1	
	BILLING	dewinurpratiwi 2021-04-13 09:30 PM	dewinurpratiwi 2021-04-19 09:28 AM	2	

Biodiversitas Journal of Biological Diversit	y Tasks 0		😢 English 🛛 👁 View Site	a mashuribiodiversitas
	Uncorrected proof [02]	editors 2021-04-20 01:28 PM	mashuribiodiversitas 2021-04-20 03:54 PM	1
		Platform &		
		OJS / PKP		

# REVIEW

Manuscript Title: **Novel chitinolytic bacteria from shrimp shell waste** Reviewer: Dr.rer.nat. Tri Dewi K. Pribadi (Universitas Padjadjaran)

# Abstract

- Please be specific in introduction sentences with the former research.
- Please separate the purpose sentence from the methods. Purpose is NOT the methods.
- What is the important of mention place (Makassar Indonesia) in the shrimp shell waste? Does it have specification on characteristic of the shell/waste/species? If not, please omit.
- How can you be sure that this is the novelty? No former research being mentioned in this topics.

Introduction

- The introduction to the chitin and chitinase is more than enough, you can probably choose the point directly related to the topics.
- Need to add a bit further about former research/update research related to the shell shrimp waste, and be specific, so you can emphasize the novelty of this study.
- Please include 'the state of the art' or at least the research urgency in relation to the local/regional/global needs of the chitinolytic bacteria.

Material and Methods

- No clear reference in the methods (Isolation and and Selection of Chitinase Enzyme Producing Bacteria). Is Soeka and Sulistiani 2012 the only reference you have in this topics? Or they only for the index formula?
- No clear references in the step of Molecular Identification of Chitinase-Producing Bacteria Isolates. Please add the references.
- No specification of the tools and equipment being used in this study. Please add the references (e.g. UV-transilluminator: brand, type, number, etc.).

Results and discussion

- Please not to mention the procedure in the results and discussion part, but mention in the methods.
- Please connect the results with the discussion, answer the 'why' of your research questions, not just performing the results.
- Please connect other research with yours, compare theirs with your results, and discuss it with all the reason.
- Please reduce describing and explaining other people's results unless it is used to explain yours.
- Please use other people's results to explain yours, either they have the same results with yours or different.

Overall comment

- Need English improvement.
- Need deeper discussion, further explanation on results.
- Accepted with revision.

Bandung, 29 March 2021

Dr.rer.nat. Tri Dewi K. Pribadi

Department of Biology Faculty of Mathematics and Natural Sciences Universitas Padjadjaran Email: tridewi.pribadi@unpad.ac.id

# Novel chitinolytic bacteria from shrimp shell waste

Abstrack. Chitinase is a capable enzyme hydrolyze chitin (a polymer of  $\beta$ -1,4 Nsetyl-D-glucosamine), it is produced by several types of microorganisms. Chitinase in nature contributes for basic science and applied. Research on the role of enzymes Chitinase is widely carried in a variety fields, in the world health because it has anti-tumor activity. In agriculture, chitinase and the producing microorganisms act as plant disease biological control agents caused by fungus. The purpose of this study was to isolate novel chitinolytic bacteria from shrimp shell waste in Makassar Indonesia, using the scatter plate method on chitin agar selective media and incubated at 37°C for 3 days, a clear zone formed around the colony as an indicator that the isolate has chitinase activity. The selection was carried out by recycling all isolates on chitin agar media and incubating them at 37°C for 3 days, then measuring the chitinolytic index. 10 isolates had chitinase activity, two isolates with the highest chitinolytic index, isolate KLA-2 (4,5) and KLA-4 (15). The molecular identification results showed that the KLA-2 and KLA-4 isolates were *Lysinibacillus fusiformis* strain WH22 and *Brevibacillus reuszeri* strain CMB-15, respectively. A novel of This research is the *B. reuszeri* can produce chitinase enzymes from shrimp shell waste.

Key words: Brevibacillus reuszeri, Chitinase enzyme, Lysinibacillus fusiformis, Shrimp Shell Waste.

Running Title: Novel Chitinolytic Bacteria.

# INTRODUCTION

Chitin is a polysaccharide, a linear polymer composed of the monomer  $\beta$ -1,4-N-acetylglucosamine (Halder *et al.*, 2013; Junianto, Wahyuntari And Setyahadi, 2013). Chitin has a rigid structure in the animal exoskeleton, because in the N-acetylglucosamine polymer chain, there are hydrogen bonds between molecules to form microfibrils to produce a stable, rigid, water-insoluble structure so that it can crystallize (Dliyauddin et al., 2020; Elieh-Ali-Komi & Hamblin, 2016)

The abundance of chitin in nature ranks second after cellulose and is widely distributed in the biosphere, such as the skin of crustaceans (crab, shrimp, and lobster), jellyfish, structural components of insect exoskeletons, fungal cell walls (22-40%), algae, nematodes, animals, plants and microorganisms contribute to the availability of chitin in the soil (Yurnaliza et al., 2012; Asif *et al.*, 2020). The source of chitin is also obtained from shrimp processing waste which consists of a head and shell (Benhabiles et al., 2012; Ghorbel-Bellaaj et al., 2012). Both of these wastes contain large chitin, the waste is thought to be a habitat for chitinase - producing bacteria (Benhabiles et al., 2012; Das et al., 2016; Ghorbel-Bellaaj et al., 2012).

Chitin can be degraded in two ways; the first is degradation by chitinolytic mechanisms which hydrolyze  $\beta$ -1,4-glycoside bonds, the second is that the polymer undergoes the first deacetylation, which is then hydrolyzed by chitosanase (Yurnaliza et al., 2012).

Exokitinase, endokitinase, chitosanase, and chitin deacetylase are chitinase enzymes (Yan & Fong, 2015), The chitinase enzyme is produced by various organisms, including bacteria, yeasts, fungi, insects, plants, vertebrates, etc (Das et al., 2016). The chitinase enzyme is included in the hydrolase enzyme group that can degrade chitin directly into small molecular weight products produced by microorganisms (Yan & Fong, 2015), both intracellular (Y. Li et al., 2016) and extracellular (Kim et al., 2017).

Chitinase enzyme has excellent potential for application in various fields (Das et al., 2016). The chitin hydrolysis results in chitooligosaccharide compounds (chitohexose and chitoheptose) are beneficial in the world of health because they have anti-tumor activity (Soeka & Sulistiani, 2012). Chitooligosaccharide compounds can also increase immunity, which is very helpful for the treatment of AIDS, cancer, heart, and blood diseases can be applied to food, medicine, and cosmetic products (Haliza, 2016).

In agriculture, chitinase and the microorganisms that produce it act as biocontrol for plant diseases caused by fungi (Berini et al., 2018; Parnell et al., 2016). Chitinase causes lysis of fungal cell walls, which generally contain chitin (A. Veliz et al., 2017; Asif et al., 2020).

Chitinase is very promising in biotechnology to be used as a safe and environmentally friendly biopesticide because chitinase can degrade chitin, a component in insect exoskeletons, nematodes, and fungal cell walls (Berini et al., 2018; Gupta et al., 2019; Le & Yang, 2019; R. Singh et al., 2020). Chitinase also plays an environmental role as a bioconversion agent for chitin waste into single-cell proteins (Om, 2020; Suryadi et al., 2016).

Comment [evika1]: Sebaiknya introduction memuat 4 kata kunci ini

The chitinase enzyme's role, which is very prospective in people's lives, encourages scientists and researchers to explore chitinolytic microorganisms. including this research entitled isolation of novel chitinolitic bacteria from shrimp shell waste.

## MATERIALS AND METHODS

#### **Chitinolytic Bacterial Isolation**

Samples in the form of solid waste processing shrimp consisting of skin, head, and tail were taken from the shrimp processing factory in Makassar Industrial Area (KIMA) Indonesia, then weighed as much as 100 grams and put into an Erlenmeyer containing 400 mL of lactose broth (LB) media as a medium. Fermentation. The fermentation medium is kept closed for  $\pm$  one month. This fermented liquid will be used as a sample in the process of isolating chitinase enzyme-producing bacteria from shrimp processing waste.

# **Colloidal Chitin Preparation**

It weighed as much as 20 grams of chitin from shrimp shells obtained from PT. Biotech Surindo-Cirebon Indonesia, added 400 mL of concentrated HCl, stirred for 2 hours then incubated in the refrigerator for 24 hours. The solution is filtered with glass wool, the filtrate is added with 200 mL cold distilled water and neutralized with 10 N NaOH ( $\pm$  250 mL) to pH 7. The solution is centrifuged at 8,000 rpm for 20 minutes at 4°C. The supernatant was removed, the sediment was added with distilled water and stirred to dissolve the remaining salt, then it was again centrifuged at a speed of 8,000 rpm for 20 minutes at 4°C. The precipitate in the form of colloidal chitin is stored in the refrigerator (Widhyastuti, 2007).

#### Isolation and Selection of Chitinase Enzyme Producing Bacteria

Media Isolation and Selection: Chitin agar (1% colloidal chitin, 0.1% peptone, 0.1% yeast extract, 0.1%  $KH_2PO_4$ , 0.05%  $MgSO_4.7H_2O$ , and 2% agar) in 100 mL of distilled water. The solution was then homogenized with a magnetic stirrer and heated until it dissolved and then adjusted the pH of the media 7. The solution was sterilized for 15 minutes at a temperature of 121°C. The agar chitin medium was used to isolate and select pure bacteria that could break down chitin and poured it into a petri dish. Isolation was carried out using the spread plate method. In the sample (liquid from fermented shrimp processing waste), dilution was carried out in stages up to 10-6, for all dilutions, it was grown by spreading it on chitin agar media, then incubation was carried out at 37°C for 3 x 24 hours.

The qualitative test results showed that the chitinolytic isolate was characterized by a halo area (clear zone) around the colony. The obtained isolates were grown simultaneously by being recycled on chitin media to use a straight loop. The selection was carried out by measuring the chitinolytic index of each isolate. The chitinolytic index is the ratio value between the diameter of the clear zone and the diameter of the bacterial colony as in the following formula: CHITINOLITIC INDEX =  $\frac{\theta \text{ HALO ZONE DIAMETER}}{\theta \text{ BACTERIAL COLONY DIAMETER}}$  (Soeka & Sulistiani, 2012)

All isolates of chitinase enzyme-producing bacteria were observed, including colony macroscopic observations (size, shape, color, elevation, edges, and the surface of the colony) and microscopic cell observations (cell shape, Gram properties, and presence or absence of endospores). All isolated chitinolytic isolates and isolates with the highest chitinolytic index value will be purified and cultured again by inoculating into chitin agar and nutrient agar (NA) slants as

#### Molecular Identification of Chitinase-Producing Bacteria Isolates

stock for further treatment and stored in the refrigerator at 4°C (Soeka & Sulistiani, 2012).

Molecular identification was only carried out on two bacterial isolates that had the highest chitinolytic index. These bacterial isolates were determined using a 16S-rRNA sequence. The steps for molecular identification and analysis of the 16S-rRNA gene were carried out as follows:

#### a. DNA extraction

DNA extraction is a series of processes for separating DNA from other cell components. DNA extraction in prokaryotic organisms is carried out through sample preparation, cell lysis, DNA binding, wash, and elution. The DNA extraction steps are as follows:

1.Sample preparation

A total of 1 ose of bacterial samples were inserted into a sterile 1.5 mL microcentrifuge tube containing 200  $\mu$ L grams (+) of a buffer that had been added with lysozyme. Homogenize by pipetting, then incubated at 37°C for 30 minutes. The tube was vortexed after incubating, then added 20  $\mu$ L Proteinase K and 200  $\mu$ L gram (-) buffer, then vortexed and reincubated at 60°C for 10 minutes, every 3 minutes, the tube was reversed to make it homogeneous.

# 2. Cell lysis

Add 200  $\mu$ L GB Buffer (Geneaid), then vortexed and incubated again at 50°C for 10 minutes, turning the tube every 3 minutes.

3. DNA Binding

Add 200 µL of absolute ethanol 96% and vortex for 10 seconds. Transfer all the mixture into a GD column (spin

column) in a 2 mL collection tube, centrifuged at a speed of 13,100 rpm for 2 minutes. Remove the collection tube that is under the spin column and replace it with a new collection tube. 4.Wash

Adding 400  $\mu$ L of W1 buffer (Geneaid) then centrifuged at 13,100 rpm for 30 seconds, then discarding the liquid in the collection tube. Add 600  $\mu$ L of wash buffer and centrifuge again, then drain the liquid into the collection tube and centrifuge again for 3 minutes. Discard the collection tube and place the sterile microcentrifuge tube at the bottom of the spin column.

5. Elution

Add 100  $\mu$ L of Elution buffer (Geneaid), let stand for 3 minutes, then centrifuge at the same speed for 1 minute. The fluid containing the DNA stored in the microcentrifuge tube was stored at -4°C for a PCR template.

#### b. PCR (Polymerase Chain Reaction) Amplification

PCR is an enzymatic synthesis process to multiply a specific nucleotide sequence in vitro (in a tube). The process includes three stages, namely denaturation, annealing, and extension. This procedure is performed on isolated DNA samples. PCR mix is inserted into the PCR tube:

#### Table 1. Composition of PCR Mix

Reaction	(µL)
Nuclease Free Water	20
2x KAPA2G Fast Ready Mix	25
63F (Forward primer)	2.5
1387R (Reverse primer)	2.5
DNA template	5
Total PCR mix	55

Comment [evika2]: Sebutkan sekuens prime forward and reverse

The total volume of PCR Mix was 55  $\mu$ L for each sample, then entered into the PCR machine. Amplification is done using a PCR (DNA thermal cycler) machine. For PCR amplification, the initial stage of pre-denaturation is at 95°C for 3 minutes, then denaturation at 95°C for 10 seconds, annealing at 55°C for 10 seconds, extension at 72°C for 15 seconds for 35 cycles followed by a final extension at 72°C for 1 minute and finally hold 4°C.

#### c. Electrophoresis

2% agarose is prepared by dissolving 2 g of agarose in 100 mL of 10X Tris borate EDTA. Then heat until boiling and dissolve. Next, 1  $\mu$ L of ethidium bromide (0.2  $\mu$ g / mL) was added and put in a gel mold that had been fitted with a comb. After the agarose has solidified (about 30 minutes), then it is put into an electrophoresis tank containing a 0.5% TBE solution. Entering the DNA of the amplified sample as much as 5  $\mu$ L, then to find out the size of the PCR amplification product, a 100 bp marker was inserted in the first well and followed by the amplified DNA sample. The electrodes are connected to the power supply then turn on for 40 minutes 100 volts. After that, the electrophoresis device is turned off then the gel from the tool is taken. The gel was transferred to the UV-transilluminator, and then the results were observed on a computer.

# d. Sequencing

PCR samples and the forward primer 63F (5'-CAGGCCTAACACATGCAAGTC-3') and reverse primer 1387R (5'-GGGCGGTGTGTACAAGGC-3') were sent to 1st BASE Malaysia for sequencing. The result is a  $\pm$  1,300bp long nucleotide sequence. These nucleotides were entered in the online BLAST (Basic Local Alignment Search Tool) program on the website NCBI (National Center for Biotechnology Information) https://www.ncbi.nlm.nih.gov. to match species data in the Gen Bank. Sequences were aligned using the Clustal W program, to see the level of kinship between species. Furthermore, phylogenetic tree construction was carried out using the neighbor-joining method using the MEGA 6 program.

# **RESULT AND DISCUSSION**

## Chitinolytic bacteria

The results of isolation and selection of chitinolytic bacteria from shrimp processing waste obtained 10 chitinolytic bacterial isolates as shown in the table below:

Table 2. Results of	Isolation and Selection of Ch	itinolytic Bacteria from Shr	imp Processing Waste
Isolate Code	Halo Zone Diameter (mm)	Bacterial Colony Diameter (mm)	Chitinolitic Index
KLA-1	12	8	1,5
KLA-2	45	10	4,5
KLA-3	15	8	1,9
KLA-4	75	5	15
KLA-5	13	5	2,6
KLA-7	7	5	1,4
KLA-8	15	13	1,2
KLA-9	11	5	2,2
KLA-10	26	7	3,7
KLA-14	9	7	1,3

Comment [evika3]: Judul tabel lebih tepat tentang hasil pengukuran halo zone...bacterial colony...chitinolitic index/ pengujian in vitro beberapa isolate bakteri

Based on the data obtained, it is known that the isolates that have the highest chitinolytic index are KLA-2 isolates, which have chitinolytic index 4.5, and KLA-4, which have chitinolytic index 15. Furthermore, the KLA-2 and KLA-4 isolates were determined as potential producing chitinase enzyme isolates originating from shrimp processing waste, which can be utilized in various fields because it has the most extensive chitinolytic index of all isolates. The chitinase activity of the two isolates can be seen in the figure below:



Figure 1. The chitinolytic activity of KLA-2 and KLA-4 isolates formed a clear zone around the bacterial colony.

The halo zone, as in Figure 1, the color of the transparent media is caused by the chitinase enzyme released into the media, which is a colorless metabolite (forming a clear zone around the bacterial colony) (Noreen et al., 2018), chitinase is an extracellular enzyme produced by chitinolytic bacteria which plays an essential role in hydrolyzing chitin (Asif et al., 2020). Extracellular enzymes are produced in cells but are released into the growth medium (Z. Zhao et al., 2020).

Each had a clear zone diameter, and colony diameter measured using a ruler on the third day of incubation from the ten isolates. The measurement results obtained can be seen again in Table 2. The size of the clear zone depends on bacteria's ability to produce chitinase, which varies widely. The difference in chitin degradation rate can be indicated by the clear area's appearance around the colony on agar containing chitin (Yavari-Bafghi et al., 2019). These differences may be due to small differences in the genes encoding them, only the *chi18D* gene on *Cellvibrio japonicus* bacteria has fundamental roles in chitin degradation substrate (Monge et al., 2018).

This study used chitin agar as a selective medium to isolate and select chitinolytic bacteria, it contains colloidal chitin, colloidal chitin is chitin dissolved in concentrated hydrochloric acid as studied by Hsu and Lockwood (1975) as a selective medium for obtaining actinomycetes from water and soil (Hsu & Lockwood, 1975), colloidal chitin is the result of the chemical degradation of chitin polymer and contains chitin oligomer (Tran et al., 2019). The substrate is used in screening as a component in chitin media to induce chitinolytic enzyme groups, such as exochitinase and endochitinase (Drewnowska et al., 2020). The screening method using chitin media was chosen because it was quite practical, this method can screen large amounts of chitinolytic isolates quickly, and the results of the screening can be seen qualitatively and semiquantitatively (Shamshina, 2019).

# Colony Macroscopic and Microscopic Observations of Bacterial Cells

Macroscopic observations are carried out directly, while microscopic of observations of cells are carried out with gram staining, which is a differential stain, and endospore staining, which is a particular stain.

**Comment [evika4]:** Pembahasan mengacu p data tabel apakah ada hubungan antara halo zon bacterial colony dengan chitinolitic index?

		Cole	ony Macrosc	opic		
Isolate Code	Size	Shape	Color	Elevation	Edge	Surface
KLA-1	Moderate	Irregular	Milky white	Flat	Rhizoid	Smooth shiny
KLA-2	Moderate	Irregular	Milky white	Flat	Rhizoid	Smooth shiny
KLA-3	Moderate	Irregular	Milky white	Flat	Entire	Rough
KLA-4	Moderate	Circular	Milky white	Flat	Entire	Smooth shiny
KLA-5	Moderate	Circular	Milky white	Flat	Entire	Smooth shiny
KLA-7	Moderate	Irregular	Milky white	Flat	Entire	Smooth shiny
KLA-8	Large	Filamentous	Milky white	Flat	Rhizoid	Rough
KLA-9	Moderate	Circular	Milky white	Flat	Entire	Smooth shiny
KLA-10	Moderate	Circular	Milky white	Flat	Entire	Smooth shiny
KLA-14	Moderate	Filamentous	Milky white	Flat	Rhizoid	Rough

# Table 4. Microscopic Observation of Chitinolytic Bacterial Isolates

	Microscopic Cell				
Isolate Code	Cell shape	Gram	Endospore		
KLA-1	Bacil	+	+		
KLA-2	Bacil	+	+		
KLA-3	Coccus	-	-		
KLA-4	Bacil	+	+		
KLA-5	Bacil	+	+		
KLA-7	Bacil	+	+		
KLA-8	Bacil	-	-		
KLA-9	Bacil	+	+		
KLA-10	Coccus	-	-		
KLA-14	Coccus	-	-		



Figure 2. The results of gram staining of two isolates that had the highest chitinolytic index

Colonies that grow on agar plates need to pay attention to size, color, transparency, edges, surface properties (elevation), and shape to obtain from macroscopic bacterial colonies (Logan and Vos, 2015).

The results obtained from bacterial macroscopic observations of the 10 chitinolytic isolates can be seen in table 3, while the results obtained from microscopic observations of bacterial cells can be seen in table 4. Colony macroscopic observations for KLA-2 and KLA-4 isolates which were designated as potential producing isolates because it has the highest chitinolytic of all filled isolates. Microscopic observations of cells for KLA-2 and KLA-4 isolates had the same

Comment [evika5]: Tidak ada di metode tiba tiba muncul data ini

Comment [evika6]: Tidak ada di metode tiba tiba muncul data ini cell microscopic characters; *bacil* cell shape (rod), gram-positive bacteria, and has *endospore* (table 4 and figure 2). According to Sizar et al., stated that in 1884 Hans Christian Gram created Gram Staining methode; gram-positive bacteria on gram stain were purplish-blue due to the crystal violet-iodine dye complex maintained even though it was given a bleaching solution of acetone alcohol. Meanwhile, gram-negative color is red because the complex dissolves when giving acetone alcohol and takes on a safranin color. The color difference indicates the difference in the structure of the

bacterial cell wall (Budin et al., 2012; Gram Positive Bacteria - StatPearls - NCBI Bookshelf, 2020).

Molecular Identification of Chitinase Enzyme Producing Bacteria



Figure 3. Electrophoresis results from the 16S-rRNA gene amplification product

Comment [evika7]: Gambar terlalu besar

From the results of the two samples' electrophoresis, it was found that there were bands that were separated and parallelled to the markers of around 1,300bp. It indicates that the amplified gene fragment is  $\pm$  1.300bp in size, so it can be concluded that the amplification process of the two chitinolytic bacterial isolates was successful. The targets that have been achieved are sufficient to identify these bacterial species. Furthermore, the results are sent to PT. Genetics Science Indonesia will then be sent to 1st BASE INT sequencing in Malaysia for mapping the base pairs that were successfully amplified. The results obtained from Malaysia were then analyzed on Gen Bank using BLAST analysis.

	_	
Isolat KLA-2		
TGGCCGCTTGAGCTTGCTCCTTTGACGTTAGCGGCGGACGGGTGAGTAACACGTGGGCAA	. [	60]
CCTACCTTATAGTTTGGGATAACTCCGGGGAAACCGGGGGCTAATACCGAATAATCTGTTTC	: [	120]
ACCTCATGGTGAAATATTGAAAGACGGTTTCGGCTGTCGCTATAGGATGGGCCCGCGGCC	; [	180]
CATTAGCTAGTTGGTGAGGTAACGGCTCACCAAGGCGACGATGCGTAGCCGACCTGAGAG	;[	240]
GGTGATCGGCCACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTAG	; [	300]
GAATCTTCCACAATGGGCGAAAGCCTGATGGAGCAACGCCGCGTGAGTGA	1	360]
CGGTTCGTAAAACTCTGTTGTAAGGGAAGAACAAGTACAGTAGTAACTGGCTGTACCTTC	; [	420]
ACGGTACCTTATTAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAG	; [	480]
TGGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGCGCGCGC	; [	540]
ATGTGAAAGCCCACGGCTCAACCGTGGAGGGTCATTGGAAACTGGGAGACTTGAGTGCAC	; [	600]
AAGAGGATAGTGGAATTCCAAGTGTAGCGGTGAAATGCGTAGAGATTTGGAGGAACACCA	· [	660]
GTGGCGAAGGCGACTATCTGGTCTGTAACTGACACTGAGGCGCGAAAGCGTGGGGGAGCAA	. [	720]
ACAGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAGTGCTAAGTGTTAGGGGGG	[	780]
TTCCGCCCCTTAGTGCTGCAGCTAACGCATTAAGCACTCCGCCTGGGGAGTACGGTCGCA	· [	840]
AGACTGAAACTCAAAGGAATTGACGGGGGCCCGCACAAGCGGTGGAGCATGTGGTTTAAT	] [	900]
TCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCCCGTTGACCACTGTAGAGATAT	[	960]
GGTTTCCCCTTCGGGGGGCAACGGTGACAGGTGGTGCATGGTTGTCTTCACCTCGGGTCGA	. [	1020]
GAGATGTTGGGTTAAGTCCCGCAACGAGCGCAACCCTTGATCTTAGTTGCCATCATTTAG	] ;	1080]
TTGGGCCCTCTAAGGTGACTGCCGGTGACAAACCTGAAGAAGGTGGGGGGATGACGTCAAA	. [	1140]
TCATCATGCCCCTTAAGACCGGGGTTAACACGGTGCTCAATGGTAGAAACCAACGGTTGC	: [	1200]
CACCCCCGGGAAGGAACCTATCCTAAAAAGTCCTTCCTTTTCGGATTGTAGGGGCACCCC	[	1260]
CCCCAGTGAACCCGGAATCCCTTAAATTCCGGAAAAAATGGCCGGGAAAAATTTCCCGG	;[	1320]
CATTTTGAAACACGCCAATTATTTTTG	[	1347]

Figure 4. Electrophoresis results from the 16S-rRNA gene amplification product

Isolat KLA-4		
GTGGGACCCTACGGAGCTAGCGGCGGACGGGTGAGTACACGTAGGCAACCTGCCTCTCAT	[	60]
${\tt ACTGGGATAACATAGGGAAACTTATGCTAATACCGGATAGGTTTTTGGACCGCATGGTCC}$	[	120]
GAAAAGAAAAGATGGCTTCTGCTATCACTGGGAGATGGGCCTGCGGCGCATTAGCTAGTT	[	180]
GGTGGGGTAACGGCCTACCAAGGCGACGATGCGTAGCCGACCTGAGAGGGTGACCGGCCA	]	240]
${\tt CACTGGGACTGAGACACGGCCCATACTCCTACGGGAGGCAGCAGTAGGGAATTTTCCACA}$	[	300]
${\tt ATGGACGAAAGTCTGATGGAGCAACGCCGCGTGAACGATGAAGGTCTTCGGATTGTAAAG}$	]	360]
${\tt TTCTGTTGTTAGGGACGAATAAGTACCGTTCGAATAGGGCGGTACCTTGACGGTACCTGA$	]	420]
CGAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAGCGTT	]	480]
GTCCGGATTTATTGGGCGTAAAGCGCGCGCGCAGGCGGCTATGTAAGTCTGGTGTTAAAGCC	]	540]
CGGAGCTCAACTCCGGTTCGCATCGGAAACTGTGTAGCTTGAGTGCAGAAGAGGAAAGCG	]	600]
GTATTCCACGTGTAGCGGTGAAATGCGTAGAGATGTGGAGGAACACCAGTGGCGAAAGCG	]	660]
GCTTTCTGGTCTGTAACTGACGCTGAGGCGCGAAAGCGTGGGGAGCAAACAGGATTAGAT	[	720]
ACCCTGGTAGTCCACGCCGTAAACGATGAGTGCTAGGTGTTGGAGGTTTCAATACCCTCA	E	780]
GTGCCGCAGCTAACGCAATAAGCACTCCGCCTGGGGAGTACGCTCGCAAGAGTGAAACTC	]	840]
AAAGGAATTGACGGGGGCCCGCACAAGCGGTGGAGCATGTGGTTTAATTCCAAGCAACGC	[	900]
GAACAACCTTACCAGGTCTTGACATCCCGCTGACCGCTCTGGAGACAGAGCTTCCCTTCG	[	960]
GGGCATCGGTGAACAGGTGGTGCATTGGTTGTCGTCAGCTCGTGTCCTGAGATGTTGGGT	[1	L020]
TTAAGTCCCGCAACCAGCGCCACCCCTTATCTTTTAGTTCCCAGCATTCAGTTGGGCAAC	[]	[080]
TCTTAAAAAGACTGCCCTTCAACAAAACCGCAGGAAAGGCGGGGGATGACGTCCAATTCA	[1	L140]
${\tt TCATGGCCCCTATGAACCTGGGCCTACCAACGGTGCTAACAATGGATTGGGTACAACCGG}$	[1	L200]
GGACCCTTAACCCCCCAAGGGTTTGGCCAATTCCCTTTAAAAGCCAATTTTCGATTTTCG	[1	L260]
GAATTGTTAGGTTGCCATCTCTCCCTAACTGGAAATCCCGGAAATCCCGGTATTAATTCC	[]	L320]
CGGGAATAACCAAGGGCCCGGGGGGAAAAACTTTTCCCGGGAATTTCTTATTTACCCCCG	[]	1380]
CCCAACAATGGTTTCAGGGCCCGCTCGAAAG	[]	[411]

Figure 5. The nucleotide base sequences of KLA-2 and KLA-4 isolates were sequenced

Alignments Download GenBank Graphics Distance tree of results						
Description	Max score	Total score	Query cover	E value	Ident	Accession
Lysinibacillus fusiformis strain WH22 16S ribosomal RNA gene, complete sequence	2124	2124	94%	0.0	97%	FJ418643.1
Lysinibacillus sp. RAB1 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KR055710.1
Bacillus sp. B-3-35 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583528.1
Bacillus sp. B-3-30 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583523.1
Lysinibacillus sp. A-3-10 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583488.1
Lysinibacillus sp. C-2-31 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583477.1
Lysinibacillus sp. C-2-16B 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583464.1
Lysinibacillus sp. B-2-20 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583447.1
Lysinibacillus sp. C-1-4 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583379.1
Lysinibacillus sp. C-1-2 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583377.1
Lysinibacillus sp. A-1-22 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT583328.1
Lysinibacillus fusiformis strain M12 16S ribosomal RNA gene, partial sequence	2122	2122	94%	0.0	97%	KT345669.1

Figure 6. BLAST analysis results for KLA-2 isolates

	Alignments Download GenBank Graphics Distance tree of results						
	Description	Max score	Total score	Query cover	E value	Ident	Accession
	Brevibacillus reuszeri strain CMB-15 16S ribosomal RNA gene, partial sequence	1958	1958	89%	0.0	95%	KJ186078.1
	Brevibacillus reuszeri strain NA-9 16S ribosomal RNA gene, partial sequence	1947	1947	89%	0.0	95%	KU254671.1
1	Brevibacillus reuszeri strain NA-8 16S ribosomal RNA gene, partial sequence	1947	1947	89%	0.0	95%	KU254670.1
	Brevibacillus reuszeri partial 16S rRNA gene, strain L71	1947	1947	89%	0.0	95%	LN890067.1
	Paenibacillus sp. DB14824 16S ribosomal RNA gene, partial sequence	1947	1947	89%	0.0	95%	KP670284.1
1	Uncultured bacterium clone PSJGSASTRA 16S ribosomal RNA gene, partial sequence	1947	1947	89%	0.0	95%	KR013217.1
21	Brevibacillus reuszeri strain BCX-22 16S ribosomal RNA gene, partial sequence	1947	1947	89%	0.0	95%	KM378576.4

Figure 7. BLAST analysis results for KLA-4 isolates

The BLAST analysis results of the two chitinolytic isolates, as shown in Figure 6, show that the KLA-2 isolate species is *Lysinibacillus fusiformis strain WH22* with a maximum score of 2124, a total score of 2124, query coverage of 94%, and identities of 97%. Meanwhile, the KLA-4 isolate species was the *B. reuszeri* strain CMB-15 with a maximum score of 1958, a total score of 1958, a query coverage of 89% and 95% identities (Figure 7).

Lysinibacillus fusiformis strain WH22 16S ribosomal RNA gene, complete sequence Sequence ID: <u>gb[FJ418643.1]</u> Length: 1544 Number of Matches: 1							
Range 1	70 to 1	44 GenBank Graphics				🔻 Next Match 🔺 Previous Match	
Score 2124 b	its(115	))	Expect 0.0	Identities 1237/1278(97%)	Gaps 10/1278(0%)	Strand Plus/Plus	
Query	9	TGAGCTTGCTCCTTTGA	CGTTAGCGGCGGA	CGGGTGAGTAACACGTGGGCAJ	CCTACCTTATAGTTTGGGATAACTCCGGGAA	accggggctartaccgartartctgtttcacctcat	G 128
Sbjet	70	TGAGCITGCTCCTTTGA	GTTAGCGGCGGA	CGGGTGAGTAACACGTGGGCAJ	CCTACCTTATAGTTTGGGATAACTCCGGGAA	accggggctaataccgaataatctgtttcacctcat	G 189
Query	129	GTGAAATATTGAAAGAC	GTTTCGGCTGTC	SCTATAGGATGGGCCCGCGGC	CATTAGCTAGTTGGTGAGGTAACGGCTCACC.	AAGGCGACGATGCGTAGCCGACCTGAGAGGGTGATC	G 248
Sbjct	190	GIGAAATATIGAAAGAC	GTTTCGGCTGTC	SCTATAGGATGGGCCCGCGGC	CATTAGCTAGTTGGTGAGGTAACGGCTCACC	AAGGCGACGATGCGTAGCCGACCTGAGAGGGTGATC	G 309
Query	249	GCCACACTGGGACTGAG.	ACACGGCCCAGAC	ICCTACGGGAGGCAGCAGTAG	GAATCTTCCACAATGGGCGAAAGCCTGATGG	AGCAACGCCGCGTGAGTGAAGAAGGATTTCGGTTCG	T 368
Sbjct	310	GCCACACTGGGACTGAG.	ACACGGCCCAGAC	ICCTACGGGAGGCAGCAGTAG	GAATCTTCCACAATGGGCGAAAGCCTGATGG	AGCAACGCCGCGTGAGTGAAGAAGGATTTCGGTTCG	I I 429
Query	369	AAAACTCTGTTGTAAGG	GAAGAACAAGTAC	AGTAGTAACTGGCTGTACCTT	ACGGTACCTTATTAGAAAGCCACGGCTAACT.	ACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAG	C 488
Sbjct	430	AAAACTCTGTTGTAAGG	GAAGAACAAGTAC	AGTAGTAACTGGCTGTACCTT	ACGGTACCTTATTAGAAAGCCACGGCTAACT.	ACGTGCCAGCAGCCGCCGTAATACGTAGGTGGCAAG	C 549
Query	489	GTTGTCCGGAATTATTG	GCGTAAAGCGCG	CGCAGGTGGTTTCTTAAGTCT	ATGTGAAAGCCCACGGCTCAACCGTGGAGGG	TCATTGGAAACTGGGAGACTTGAGTGCAGAAGAGGA	T 608
Sbjet	550	GTTGTCCGGAATTATTG	GCGTAAAGCGCG	CGCAGGTGGTTTCTTAAGTCT	ATGTGAAAGCCCACGGCTCAACCGTGGAGGG	TCATTGGAAACTGGGAGACTTGAGTGCAGAAGAGGA	E 669
Query	609	AGTGGAATTCCAAGTGT.	AGCGGTGAAATGC	GTAGAGATTTGGAGGAACACC/	GTGGCGAAGGCGACTATCTGGTCTGTAACTG	ACACTGAGGCGCGAAAGCGTGGGGAGCAAACAGGAT	T 728
Sbjct	670	AGTGGAATTCCAAGTGT.	AGCGGTGAAATGC	STAGAGATITGGAGGAACACC/	GTGGCGAAGGCGACTATCTGGTCTGTAACTG	ACACTGAGGCGCGAAAGCGTGGGGAGCAAACAGGAT	I I 789
Query	729	AGATACCCTGGTAGTCC.	ACGCCGTAAACGA	IGAGIGCTAAGIGITAGGGGGI	TICCGCCCCTTAGTGCTGCAGCTAACGCATT.	AAGCACTCCGCCTGGGGAGTACGGTCGCAAGACTGA	A 848
Sbjet	790	AGATACCCTGGTAGTCC.	ACGCCGTAAACGA	IGAGIGCTAAGIGITAGGGGGI	TTCCGCCCCTTAGTGCTGCAGCTAACGCATT.	AAGCACTCCGCCTGGGGAGTACGGTCGCAAGACTGA	A 909
Query	849	ACTCAAAGGAATTGACG	GGGCCCGCACAA	SCGGTGGAGCATGTGGTTTAA	TCGAAGCAACGCGAAGAACCTTACCAGGTCT	TGACATCCCGTTGACCACTGTAGAGATATGGTTTCC	C 968
Sbjct	910	ACTCAAAGGAATTGACG	GGGCCCGCACAA	SCGGTGGAGCATGTGGTTTAA	TCGAAGCAACGCGAAGAACCTTACCAGGTCT	TGACATCCCGTTGACCACTGTAGAGATATGGTTTCC	C 1029
Query	969	CTTCGGGGGGCAACGGTG.	ACAGGTGGTGCAT	GGTTGTCTTCACCTCGGGTCG	GAGATGTTGGGTTAAGTCCCGCAACGAGCGC	aaccettgatettagttgecateatttagttgggec	C 1088
Sbjct	1030	CTTCGGGGGGCAACGGTG.	ACAGGTGGTGCAT	GTTGTCGTCAGCTCGTGTCG	GAGATGTTGGGTTAAGTCCCGCAACGAGCGC	AACCCTTGATCTTAGTTGCCATCATTTAGTTGGGCA	C 1149
Query	1089	TCTAAGGTGACTGCCGG	IGACAAACCTGAA	SAAGGTGGGGGGATGACGTCAAJ	TCATCATGCCCCTTAAGACCGGGGGTTA-ACA	CGGTGCT-CAATGGTAGAAACCAACGGTTGCCACCC	C 1206
Sbjct	1150	TCTAAGGTGACTGCCGG	IIIIIIIIIIIIIII	GAAGGT-GGGGATGACGTCAA	ITCATCATGCCCCTTATGACCTGGGCTACACA	CG-TGCTACAATGGACGATACAAACGGTTGCCAACT	l C 1267
Query	1207	-CGGGAAGGAACCTATC	TAAAAAGTCCTT	C-CTTTTCGGATTGTAGGG-GG	Accoccce-cagtgaacccggaatc 127	9	
Sbjct	1268	GCGAGAGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	GATAAAGTCGTT	CTCAGTTCGGATTGTAGGCTG	CAACTCGCCTACA-TGAAGCCGGAATC 134	4	

Figure 8. Comparison of the nucleotide sequence of KLA-2 isolates (Query) with L. fusiformis bacteria strain WH22 (Sbjct)

Sequence	e ID: <u>qbl</u>	J186078.1 Length: 1	416 Number of	Matches: 1				
Range 1	: 19 to 12	9 GenBank Graphics				Next Match Previous Match		
Score			Expect	Identities	Gaps	Strand		
1958 b	its(1060		0.0	1204/1268(95%)	31/1268(2%)	Plus/Plus		
Query	12	CGGA <sup>1</sup> GCTAGCGGCGGA	CGGGTGAGT <sup>®</sup> A	CACGTAGGCAACCTGCCTCTCATAC	GGGATAACATAGGGAAACTTATGCTAATA	CCGGATAGGTTTTTGGACCGCATGGTCCGA	AAAGAAA	129
sojec	1.5	CEERGECIRECEGCEER	Ceceleaeiaa	CAUGIAGGUARCEIGCEICICAGAC.	GGGATAACATAGGGAAACTTATGCTAATA	CCGGATAGGIIIIIGGACCGCAIGGICCGA	AAAGAAA	190
Query	130 139	AGATGGCTTCTGCTATC	ACTGGGAGATG	GGCCTGCGGCGCATTAGCTAGTTGG GGCCTGCGGCGCATTAGCTAGTTGG	GGGGTAACGGCCTACCAAGGCGACGATGC	GTAGCCGACCTGAGAGGGTGACCGGCCACH	CTGGGAC	249
	250	TOLONOLOGOCOUTIC	TCOTACCCAC		CLOCLE CTOTOL TOCLOCL COCCOCCT	CARCONCONCEPTORON TTOTA A 8 CT	CTOTOT	260
Sbjet	259	TGAGACACGGCCCAGAC	TCCTACGGGAG	GCAGCAGTAGGGAATTTTCCACAAT	GACGAAAGTCTGATGGAGCAACGCCGCGT	GAACGATGAAGGTCTTCGGATTGTAAAGTT	CTGTTGT	378
Query	370	TAGGGACGAATAAGTAC	CGTTCGAATAG	GGCGGTACCTTGACGGTACCTGACG	GAAAGCCACGGCTAACTACGTGCCAGCAG	CCGCGGTAATACGTAGGTGGCAAGCGTTGI	CCGGATT	489
Sbjet	379	TAGGGACGAATAAGTAC	CGTTCGAATAG	GCCGGTACCTTGACGGTACCTGACG	GAAAGCCACGGCTAACTACGTGCCAGCAG	CCGCGGTAATACGTAGGTGGCAAGCGTTGT	CCGGATT	498
Query	490	TATTGGGCGTAAAGCGC	GCGCAGGCGGC	TATGTAAGTCTGGTGTTAAAGCCCG	AGCTCAACTCCGGTTCGCATCGGAAACTG	TGTAGCTTGAGTGCAGAAGAGGAAAGCGGI	ATTCCAC	609
Sbjet	499	TATTGGGCGTAAAGCGC	GCGCAGGCGGC	TATGTAAGTCTGGTGTTAAAGCCCG	AGCTCAACTCCGGTTCGCATCGGAAACTG	TGTAGCTTGAGTGCAGAAGAGGAAAGCGGI	ATTCCAC	618
Query	610	GTGTAGCGGTGAAATGC	GTAGAGATGTG	GAGGAACACCAGTGGCGAAAGCGGC	TTCTGGTCTGTAACTGACGCTGAGGCGCG	AAAGCGTGGGGAGCAAACAGGATTAGATAC	CCTGGTA	729
Sbjet	619	GTGTAGCGGTGAAATGC	GTAGAGATGTG	GAGGAACACCAGTGGCGAAGGCGGC	TTCTGGTCTGTAACTGACGCTGAGGCGCG	AAAGCGTGGGGGAGCAAACAGGATTAGATAC	CCTGGTA	738
Query	730	GICCACGCCGIAAACGA	TGAGTGCTAGG	IGTIGGAGGTITCAATACCCICAGI	CCGCAGCTAACGCAATAAGCACTCCGCCT	GGGGAGTACGCTCGCAAGAGTGAAACTCAA	AGGAATT	849
Sbjet	739	GTCCACGCCGTAAACGA	TGAGIGCIAGG	IGTIGGGGGTTTCAATACCCTCAGT	CCGCAGCTAACGCAATAAGCACTCCGCCT	GGGGAGTACGCTCGCAAGAGTGAAACTCAA	AGGAATT	858
Query	850	GACGGGGGGCCCGCACAA	GCGGTGGAGCA	igiggittaattccaagcaacgcgaj	CAACCTTACCAGGTCTTGACATCCCGCTG	ACCGCTCTGGAGACAGAGCTTCCCTTCGGG	GCATCGG	969
Sbjct	859	GACGGGGGGCCCGCACAA	GCGGTGGAGCA	IGTGGTTTAATTCGAAGCAACGCGAJ	GAACCTTACCAGGTCTTGACATCCCGCTG	ACCGCTCTGGAGACAGAGCTTCCCTTCGG	GCAGCGG	978
Duerv	970	TGAACAGGTGGTGCATT	GGTTGTCGTCA	SCTOSTSTCCTSAGATGTTSGCTTT	AGTCCCGCAACCAGCGCCACCCCTTATCT	TTTAGTTCCCAGCATTCAGTTGGGCAACTC	TTAAAAA	1089
Sbjct	979	TG-ACAGGTGGTGCATG	GGTTGTCGTCA	GCTCGTGTCGTGAGATGTTGGG-TT	AGTCCCGCAACGAGCGCAACCC-TTATC-	TTTAGTTGCCAGCATTCAGTTGGGCA-CTC	-TAGAGA	1092
Query	1090	GACTGCCCTTCAACAAA	ACCGCAGGAAA	GCCGGGGGATGACGTCCAATTCATC	ATGGCCCCT-ATGAACCTGGGCCTACCAAC	COTOCTAACAATGGATTGGGTACAACCGGG	GACCCTT	1208
Sbjet	1093	GACTGCCGT-CGACAAG	ACGG-AG-AAA	GGC-GGGGATGACGTCAAA-TCATC	TGCCCCCTTATGACCTTGGGC-TACCAAC	G-TGCTA-CAATGG-TIGG-TACAAC-GGG	-ACGCT-	1199
Query	1209	AACCCCCCAAGGGTTTG	CCAATTCCCT	TTAAAAGCCAATTT-TCGATTTTCG	AATTGTTAGGTTGC 1275			

Figure 9. Comparison of the nucleotide sequence of KLA-4 isolates (Query) with the *B. reuszeri* bacteria *strain CMB-15* (Sbjct)



Figure 10. The phylogenetic trees of both isolates and relative species used the neighbor-joining method based on the results of 16S-rRNA analysis

After the KLA-2 and KLA-4 isolate species were identified, phylogenetic tree analysis was carried out to precisely construct the relationships between organisms and estimate the differences between one ancestor to the offspring (Amit Roy, 2014). The 16S-rRNA gene is widely used for phylogenetic tree construction because the nucleotide base sequence between rRNA molecules can be compared precisely (Amit Roy, 2014; Woese, 1987), making it easier to identify the diversity, the 16S-rRNA gene in bacteria has a high level of similarity because it is present in all prokaryotes (Tremblay et al., 2015; Woese, 1987).

Phylogenetic tree analysis in two stages using the Clustal W program combined with the MEGA 6 (Molecular Evolutionary Genetics Analysis) program (Horiike, 2016). The first stage uses the Clustal W program to align the second sequence of chitinolytic bacterial isolates with Gen Bank sequences, which have sequences similar to KLA-2 and KLA-4 isolates. The MEGA 6 program will then create its phylogenetic tree using the "Neighbor Joining" approach, 1000x bootstrap, and the p-distance method (Horiike, 2016; Tamura et al., 2013).

The phylogenetic analysis results using 16S-rRNA coding DNA sequences as seen in Figure 10, it is known that the KLA-2 isolate is the closest relative to *L. fusiformis* strain WH22 with a genetic distance of 0.04 and a bootstrap value of 100. The greater the bootstrap value, the higher the bootstrap value. The level of trustworthiness of the reconstructed tree topology (Chang et al., 2019; Reddy et al., 2017). The KLA-4 isolate had the closest relationship with the *B. reuszeri* strain CMB-15 with a genetic distance of 0.07 and a bootstrap value of 100 compared to other types of bacteria in the Gen Bank.

#### Lysinibacillus fusiformis

*Lysinibacillus fusiformis* was initially known as *Bacillus fusiformis* before 2007; after 2007, *L. fusiformis* was transferred to the genus *Lysinibacillus*, along with Lysinibacillus sphaericus because based on the results of the molecular analysis using the 16S-rRNA gene sequence alignments, it has different sequences of the nucleus of genus *Bacillus*. The meaning of "lysine" in members of the genus *Lysinibacillus* indicates lysine, alanine, glutamic acid, and aspartic acid in the cell wall peptidoglycan layer (Ahmed et al., 2007).

Singh (2012) has successfully isolated chitinolytic bacteria from the rhizosphere of peanut plants in India; after being identified morphologically, biochemically, and molecularly. The chitinolytic bacteria 16S-rDNA gene was identified as *L. fusiformis B-CM18*. Purification of the chitinase enzyme with gel filtration and ion exchange Sephadex G-100 obtained a molecular weight of 20 kDa, a purity of 7.1 and a specific chitinase activity of 0.22 U / mg and an optimum temperature of 35° C. After finding the Pure chitinase enzyme, He conducted an in vitro antifungal activity test of chitinase crude extract against various plant pathogenic fungi and the result was the highest zone of inhibition recorded against *Fusarium oxysporum f. sp. Ciceri* (4.6 mm), *Fusarium solani* (4.1 mm), followed by *Fusarium oxysporum f. sp. lycopersici* (3.8 mm), and *Macrophomina phaseolina* (3.4 mm) (R. K. Singh et al., 2013).

This study proves that the bacteria *L. fusiformis* can produce the chitinase enzyme, an enzyme used by bacteria to degrade chitin, and chitinase is also used by bacteria to assimilate chitin as a source of carbon and nitrogen (Asif et al., 2020). Apart from being chitinolytic bacteria, the bacteria *L. fusiformis* can also produce protease enzymes (Verma et al., 2016), indole acetic acid, salysil acid, siderophores, can dissolve phosphate, and can fix nitrogen (Singh RJ, Kumar DP, 2013). These bacteria also can detoxify factory wastewater (He et al., 2011), biotransformation of isoeugenol into vanillin (L. Zhao et al., 2016), degradation of quorum-sensing molecules of N-acyl-homoserine lactone (Yong et al., 2015), and as a biotrickling filter or gas filter. Chlorobenzene (Z. X. Li et al., 2014) as well as a biocontrol agent (Wang et al., 2013).

#### Brevibacillus reuszeri

Brevibacillus reuszeri was initially known as Bacillus reuszeri before 1996; after 1996, B. reuszeri was transferred to

the genus *Brevibacillus*, along with 9 other species because based on the results of the molecular analysis using the 16SrRNA gene sequence alignments have different nucleotide sequences with members of the genus *Bacillus* (Shida et al. 1996). *B. reuszeri* is a gram-positive, motile bacterium with flagella and ellipsoidal spores. These bacteria have the shape of stem cells measuring about 0.5-0.9  $\mu$ m to 2.0-5.0  $\mu$ m. Colonies are flat, smooth, circular, and even. It is obligate aerobic, which means that it can utilize oxygen to metabolize various sugars and other carbohydrates (Johnson & Dunlap, 2019).

Prasanna (2013) has succeeded in isolating new strains of bacteria belonging to the genus *Brevibacillus* from mangrove swamplands in India, which show entomopathogenic and chitinolytic activity (Prasanna et al., 2013), not from *B. reuszeri* but *B. laterosporus*.

Several studies have reported the abilities possessed by the *B. reuszeri* bacteria, including being able to produce extracellular metalloprotease/thermolysin inhibitors (Kobayashi et al., 2004), which can produce L-methionine-N-carbamoylase, which is a potential biocatalyst for the production of L-amino acids (Nandanwar et al., 2013), can dissolve phosphate and be used in plant growth-promoting rhizobacteria (PGPR) plants (Pradesh et al., 2019; Prasad et al., 2019; Yildirim et al., 2011), as well as the production of antimicrobial bioactive compounds (Gholizadeh et al., 2013; Yang & Yousef, 2018).

This research provides evidence of new findings that the *B. reuszeri* bacteria isolated from shrimp processing waste can produce chitinase enzymes.

# REFERENCE

- A. Veliz, E., Martínez-Hidalgo, P., & M. Hirsch, A. 2017. Chitinase-producing bacteria and their role in biocontrol. AIMS Microbiology, 3(3), 689–705. DOI:10.3934/microbiol.2017.3.689.
- Ahmed, I., Yokota, A., Yamazoe, A., & Fujiwara, T. 2007. Proposal of Lysinibacillus boronitolerans gen. nov. sp. nov., and transfer of Bacillus fusiformis to Lysinibacillus fusiformis comb. nov. and Bacillus sphaericus to Lysinibacillus sphaericus comb. nov. International Journal of Systematic and Evolutionary Microbiology, 57(5), 1117–1125. DOI:10.1099/ijs.0.63867-0.
- Amit Roy, S. R. 2014. Molecular Markers in Phylogenetic Studies-A Review. Journal of Phylogenetics & Evolutionary Biology, 02(02). DOI:10.4172/2329-9002.1000131.
- Asif, T., Javed, U., Zafar, S. B., Ansari, A., Ul Qader, S. A., & Aman, A. 2020. Bioconversion of Colloidal Chitin Using Novel Chitinase from Glutamicibacter uratoxydans Exhibiting Anti-fungal Potential by Hydrolyzing Chitin Within Fungal Cell Wall. Waste and Biomass Valorization, 11(8), 4129–4143. DOI:10.1007/s12649-019-00746-2.
- Benhabiles, M. S., Salah, R., Lounici, H., Drouiche, N., Goosen, M. F. A., & Mameri, N. 2012. Antibacterial activity of chitin, chitosan and its oligomers prepared from shrimp shell waste. Food Hydrocolloids, 29(1), 48–56. DOI:10.1016/j.foodhyd.2012.02.013.
- Berini, F., Katz, C., Gruzdev, N., Casartelli, M., Tettamanti, G., & Marinelli, F. 2018. Microbial and viral chitinases: Attractive biopesticides for integrated pest management. Biotechnology Advances, 36(3), 818–838. DOI:10.1016/j.biotechadv.2018.01.002.
- Budin, G., Chung, H. J., Lee, H., & Weissleder, R. 2012. A Magnetic Gram Stain for Bacterial Detection. Angewandte Chemie, 124(31), 7872–7875. DOI:10.1002/ange.201202982.
- Chang, J.-M., Floden, E. W., Herrero, J., Gascuel, O., Di Tommaso, P., & Notredame, C. 2019. Incorporating alignment uncertainty into Felsenstein's phylogenetic bootstrap to improve its reliability. Bioinformatics, February, 1–9. DOI:10.1093/bioinformatics/btz082.
- Das, S., Roy, D., & Sen, R. 2016. Utilization of Chitinaceous Wastes for the Production of Chitinase. In Advances in Food and Nutrition Research (1st ed., Vol. 78). Elsevier Inc. DOI:10.1016/bs.afnr.2016.04.001.
- Dliyauddin, M. O. H., Ardyati, T. R. I., & Suharjono. 2020. Evaluation of enzymatic activity and identification of potent proteolytic and chitinolytic bacteria isolated from crab shell waste. Biodiversitas, 21(1), 211–218. DOI:10.13057/biodiv/d210127.
- Drewnowska, J. M., Fiodor, A., Barboza-Corona, J. E., & Swiecicka, I. 2020. Chitinolytic activity of phylogenetically diverse Bacillus cereus sensu lato from natural environments. Systematic and Applied Microbiology, 43(3), 126075. DOI:10.1016/j.syapm.2020.126075.
- Elieh-Ali-Komi, D., & Hamblin, M. R. 2016. Chitin and Chitosan: Production and Application of Versatile Biomedical Nanomaterials. International Journal of Advanced Research, 4(3), 411–427. http://www.ncbi.nlm.nih.gov.
- Gholizadeh, S. S., Baserisalehi, M., & Bahador, N. 2013. Study on bioactive compounds produced by soil origin Brevibacillus spp. Nature Environment and Pollution Technology, 12(2), 209–214.
- Ghorbel-Bellaaj, O., Younes, I., Maâlej, H., Hajji, S., & Nasri, M. 2012. Chitin extraction from shrimp shell waste using Bacillus bacteria. International Journal of Biological Macromolecules, 51(5), 1196–1201. DOI:10.1016/j.ijbiomac.2012.08.034.
- Gram Positive Bacteria StatPearls NCBI Bookshelf. 2020. Retrieved February 21, 2021, from https://www.ncbi.nlm.nih.gov.

Gupta, M. M., Chourasiya, D., & Sharma, M. P. 2019. Diversity of arbuscular mycorrhizal fungi in relation to sustainable

plant production systems. In Microbial Diversity in Ecosystem Sustainability and Biotechnological Applications: Volume 2. Soil & Agroecosystems (Vol. 2). DOI:10.1007/978-981-13-8487-5\_7.

Halder, S. K., Maity, C., Jana, A., Das, A., Paul, T., Das Mohapatra, P. K., Pati, B. R., & Mondal, K. C. 2013. Proficient biodegradation of shrimp shell waste by Aeromonas hydrophila SBK1 for the concomitant production of antifungal chitinase and antioxidant chitosaccharides. International Biodeterioration and Biodegradation, 79, 88–97. DOI:10.1016/j.ibiod.2013.01.011.

Haliza, W. 2016. Karakteristik Kitinase Dari Mikrobia. Buletin Teknologi Pasca Panen, 8(1), 1-14.

- He, M., Li, X., Liu, H., Miller, S. J., Wang, G., & Rensing, C. 2011. Characterization and genomic analysis of a highly chromate resistant and reducing bacterial strain Lysinibacillus fusiformis ZC1. Journal of Hazardous Materials, 185(2–3), 682–688. DOI:10.1016/j.jhazmat.2010.09.072.
- Horiike, T. 2016. an Introduction To Molecular Phylogenetic Analysis. Reviews in Agricultural Science, 4(0), 36–45. DOI:10.7831/ras.4.36.
- Hsu, S. C., & Lockwood, J. L. 1975. Powdered Chitin Agar as a Selective Medium for Enumeration of Actinomycetes in Water and Soil1. Applied Microbiology, 29(3), 422–426. DOI:10.1128/am.29.3.422-426.1975.
- Johnson, E. T., & Dunlap, C. A. 2019. Phylogenomic analysis of the Brevibacillus brevis clade: a proposal for three new Brevibacillus species, Brevibacillus fortis sp. nov., Brevibacillus porteri sp. nov. and Brevibacillus schisleri sp. nov. Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology, 112(7), 991–999. DOI:10.1007/s10482-019-01232-4.
- Junianto, J., Wahyuntari, B., & Setyahadi, S. 2013. Selection of Methods for Microbiological Extraction of Chitin from Shrimp Shells. Microbiology Indonesia, 7(2), 75–83. DOI:10.5454/mi.7.2.5.
- Kim, Y. H., Park, S. K., Hur, J. Y., & Kim, Y. C. 2017. Purification and characterization of a major extracellular chitinase from a biocontrol bacterium, Paenibacillus elgii HOA73. Plant Pathology Journal, 33(3), 318–328. DOI:10.5423/PPJ.FT.01.2017.0022.
- Kobayashi, T., Nakagawa, N., Imada, C., Hamada-Sato, N., & Watanabe, E. 2004. Occurrence of a metalloprotease (thermolysin) inhibitor among Brevibacillus species and purification of such inhibitor from Brevibacillus reuszeri IFO 15719T. Fisheries Science, 70(2), 299–305. DOI:10.1111/j.1444-2906.2003.00804.x.
- Le, B., & Yang, S. H. 2019. Microbial chitinases: properties, current state and biotechnological applications. World Journal of Microbiology and Biotechnology, 35(9), 1–12. DOI:10.1007/s11274-019-2721-y.
- Li, Y., Lei, X., Zhu, H., Zhang, H., Guan, C., Chen, Z., Zheng, W., Fu, L., & Zheng, T. 2016. Chitinase producing bacteria with direct algicidal activity on marine diatoms. Scientific Reports, 6(February), 1–13. DOI:10.1038/srep21984.
- Li, Z. X., Yang, B. R., Jin, J. X., Pu, Y. C., & Ding, C. 2014. The operating performance of a biotrickling filter with Lysinibacillus fusiformis for the removal of high-loading gaseous chlorobenzene. Biotechnology Letters, 36(10), 1971–1979. DOI:10.1007/s10529-014-1559-5.
- Logan, N. A., & Vos, P. De. 2015. Bacillus . In Bergey's Manual of Systematics of Archaea and Bacteria. DOI:10.1002/9781118960608.gbm00530.
- Monge, E. C., Tuveng, T. R., Vaaje-Kolstad, G., Eijsink, V. G. H., & Gardner, J. G. 2018. Systems analysis of the glycoside hydrolase family 18 enzymes from Cellvibrio japonicus characterizes essential chitin degradation functions. Journal of Biological Chemistry, 293(10), 3849–3859. DOI:10.1074/jbc.RA117.000849.
- Nandanwar, H. S., Vohra, R. M., & Hoondal, G. S. 2013. Trimeric l-N-carbamoylase from newly isolated Brevibacillus reuszeri HSN1: A potential biocatalyst for production of l-α-amino acids. Biotechnology and Applied Biochemistry, 60(2), 219–230. DOI:10.1002/bab.1066.
- Noreen, N., Ramzan, N., Perveen, Z., Shahzad, S., Noreen, N., & Al, E. T. 2018. Assessing The Enzymatic Activities of Compost Associated Mesophilic, Thermotolerant And Thermophilic Bacteria And Fungi. 15, 815–825.

Om, H. 2020. Role of Chitinases as a waste management to control global crisis. 303–313.

- Parnell, J. J., Berka, R., Young, H. A., Sturino, J. M., Kang, Y., Barnhart, D. M., & Dileo, M. V. 2016. From the lab to the farm: An industrial perspective of plant beneficial microorganisms. Frontiers in Plant Science, 7(AUG2016), 1–12. DOI:10.3389/fpls.2016.01110.
- Pradesh, H., Yadav, N., Pradesh, U., Kumar, A., South, R. G., Pradesh, U., Meena, V. S., & Division, C. P. 2019. Plant Growth Promoting Rhizobacteria for Agricultural Sustainability. In Plant Growth Promoting Rhizobacteria for Agricultural Sustainability. DOI:10.1007/978-981-13-7553-8.
- Prasad, M., Srinivasan, R., Chaudhary, M., Choudhary, M., & Jat, L. K. 2019. Plant Growth Promoting Rhizobacteria (PGPR) for Sustainable Agriculture. In PGPR Amelioration in Sustainable Agriculture. Elsevier Inc. DOI:10.1016/b978-0-12-815879-1.00007-0.
- Prasanna, L., Eijsink, V. G. H., Meadow, R., & Gåseidnes, S. 2013. A novel strain of Brevibacillus laterosporus produces chitinases that contribute to its biocontrol potential. Applied Microbiology and Biotechnology, 97(4), 1601–1611. DOI:10.1007/s00253-012-4019-y.
- Reddy, S., Kimball, R. T., Pandey, A., Hosner, P. A., Braun, M. J., Hackett, S. J., Han, K. L., Harshman, J., Huddleston, C. J., Kingston, S., Marks, B. D., Miglia, K. J., Moore, W. S., Sheldon, F. H., Witt, C. C., Yuri, T., & Braun, E. L. 2017. Why do phylogenomic data sets yield conflicting trees? Data type influences the avian tree of life more than taxon sampling. Systematic Biology, 66(5), 857–879. DOI:10.1093/sysbio/syx041.

- Shamshina, J. L. 2019. Chitin in ionic liquids: Historical insights into the polymer's dissolution and isolation. A review. Green Chemistry, 21(15), 3974–3993. DOI:10.1039/c9gc01830a.
- Shida Osamu, Hiroaki Takagi, Kiyoshi Kadowaki, K. K. 1996. Proposal for Two New Genera, Brevibacillus gen. nov. and Aneurinibacillus gen. nov. International Journal of Systematic Bacteriology, 46(4), 939–946.
- Singh, R. K., Kumar, D. P., Solanki, M. K., Singh, P., Srivastva, A. K., Kumar, S., Kashyap, P. L., Saxena, A. K., Singhal, P. K., & Arora, D. K. 2013. Optimization of media components for chitinase production by chickpea rhizosphere associated Lysinibacillus fusiformis B-CM18. Journal of Basic Microbiology, 53(5), 451–460. DOI:10.1002/jobm.201100590.
- Singh, R., Upadhyay, S. K., Singh, M., Sharma, I., Sharma, P., Kamboj, P., Saini, A., Voraha, R., Sharma, A. K., Upadhyay, T. K., & Khan, F. 2020. Chitin, chitinases and chitin derivatives in biopharmaceutical, agricultural and environmental perspective. Biointerface Research in Applied Chemistry, 11(3), 9985–10005. DOI:10.33263/BRIAC113.998510005.
- Singh RJ, Kumar DP, and S. M. 2013. Multifarious plant growth promoting characteristics of chickpea rhizosphere associated Bacilli help to suppress soil-borne pathogens. Plant Growth Regulation, Springer, 73(1), 91–101.
- Soeka, Y. S., & Sulistiani, S. 2012. Seleksi, Karakterisasi, dan Identifikasi Bakteri Penghasil Kitinase yang Diisolasi dari Gunung Bromo Jawa Timur. Jurnal Natur Indonesia, 13(2), 155. DOI:10.31258/jnat.13.2.155-161.
- Suryadi, Y., Priyatno, T. P., Samudra, I. M., Susilowati, D. N., Lawati, N., & Kustaman, E. 2016. Pemurnian Parsial dan Karakterisasi Kitinase Asal Jamur Entomopatogen Beauveria bassiana Isolat BB200109. Jurnal AgroBiogen, 9(2), 77. DOI:10.21082/jbio.v9n2.2013.p77-84.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. Molecular Biology and Evolution, 30(12), 2725–2729. DOI:10.1093/molbev/mst197.
- Tran, T. N., Doan, C. T., Nguyen, V. B., Nguyen, A. D., & Wang, S. L. 2019. The isolation of chitinase from Streptomyces thermocarboxydus and its application in the preparation of chitin oligomers. Research on Chemical Intermediates, 45(2), 727–742. DOI:10.1007/s11164-018-3639-y.
- Tremblay, J., Singh, K., Fern, A., Kirton, E. S., He, S., Woyke, T., Lee, J., Chen, F., Dangl, J. L., & Tringe, S. G. 2015. Primer and platform effects on 16S rRNA tag sequencing. Frontiers in Microbiology, 6(AUG), 1–15. DOI:10.3389/fmicb.2015.00771.
- Verma, P., Yadav, A. N., Khannam, K. S., Kumar, S., Saxena, A. K., & Suman, A. 2016. Molecular diversity and multifarious plant growth promoting attributes of Bacilli associated with wheat (Triticum aestivum L.) rhizosphere from six diverse agro-ecological zones of India. Journal of Basic Microbiology, 56(1), 44–58. DOI:10.1002/jobm.201500459.
- Wang, K., Yan, P. sheng, Ding, Q. long, Wu, Q. xi, Wang, Z. bo, & Peng, J. 2013. Diversity of culturable rootassociated/endophytic bacteria and their chitinolytic and aflatoxin inhibition activity of peanut plant in China. World Journal of Microbiology and Biotechnology, 29(1), 1–10. DOI:10.1007/s11274-012-1135-x.
- Widhyastuti, N. 2007. Produksi Kitinase Ekstraseluler Aspergilus rugulosus 501 Secara Optimal pada Media Cair. Berita Biologi, 8(6), 547–553.
- Yan, Q., & Fong, S. S. 2015. Bacterial chitinase: Nature and perspectives for sustainable bioproduction. Bioresources and Bioprocessing, 2(1). DOI:10.1186/s40643-015-0057-5
- Yang, X., & Yousef, A. E. 2018. Antimicrobial peptides produced by Brevibacillus spp.: structure, classification and bioactivity: a mini review. World Journal of Microbiology and Biotechnology, 34(4), 0. DOI:10.1007/s11274-018-2437-4.
- Yavari-Bafghi, M., Babavalian, H., & Amoozegar, M. A. 2019. Isolation, screening and identification of haloarchaea with chitinolytic activity from hypersaline lakes of Iran. Archives of Biological Sciences, 71(1), 71–81. DOI:10.2298/ABS180525049Y.
- Yildirim, E., Karlidag, H., Turan, M., Dursun, A., & Goktepe, F. 2011. Promoition of Broccoli by Plant Growth Promoting Rhizobacteria. Hort Science, 46(6), 932–936. http://hortsci.ashspublications.org.
- Yong, D., Ee, R., Lim, Y. L., Chang, C. Y., Yin, W. F., & Chan, K. G. 2015. Insights on quorum-quenching properties of Lysinibacillus fusiformis strain RB21, a Malaysian municipal solid-waste landfill soil isolate, via complete genome sequence analysis. Genome Announcements, 3(3), 2014–2015. DOI:10.1128/genomeA.00409-15.
- Yurnaliza, Y., Margino, S., & Sembiring, L. 2012. Kemampuan Kitinase Streptomyces RKt5 sebagai Antijamur terhadap Patogen Fusarium oxysporum. Jurnal Natur Indonesia, 14(1), 42. DOI:10.31258/jnat.14.1.42-46.
- Zhao, L., Bao, G., Geng, B., Song, J., & Li, Y. 2016. Draft genome sequence of Lysinibacillus fusiformis strain SW-B9, a novel strain for biotransformation of isoeugenol to vanillin. Genome Announcements, 3(2), 9–10. DOI:10.1128/genomeA.00289-15.
- Zhao, Z., Baltar, F., & Herndl, G. J. 2020. Linking extracellular enzymes to phylogeny indicates a predominantly particleassociated lifestyle of deep-sea prokaryotes. Science Advances, 6(16), 1–11. DOI:10.1126/sciadv.aaz4354.



Mashuri Masri <mashuri.masri@uin-alauddin.ac.id>

# Mashuri Masri-UIN Alauddin makassar

8 pesan

Mashuri Masri <mashuri.masri@uin-alauddin.ac.id> Kepada: translate.sedekah@gmail.com

assalamualaikum

kami mashuri masri, teman bu kurnia dosen biologi UIN alauddin makassar. mohon bantuannya utk proof reading naskah kami, seperti naskah bu kurnia beberapa bulan yg lalu.

terimakasih sebelumnya. salam hormat

V-Novel chitinolytic bacteria from shrimp (edit1).docx 2952K

Translate Sedekah <translate.sedekah@gmail.com> Kepada: Mashuri Masri <mashuri.masri@uin-alauddin.ac.id>

Dear Bapak Mashuri Masri,

Terima kasih atas kiriman naskahnya.

Total biaya untuk naskah anda adalah Rp. 272.108,-

dengan rincian: Proofreading 4612 kata x Rp. 59/kata (Tanpa Daftar Pustaka)

Estimasi waktu penyelesaian : Sabtu, 10 April 2021. (Harap Konfirmasi melalui wa/membalas email ini jika Bapak/Ibu sepakat dengan biaya translate dan waktu penyelesaian agar naskah segera kami proses).

Jika Bapak/lbu membutuhkan waktu lebih cepat dari estimasi waktu yang kami tawarkan, maka opsi percepatan bisa digunakan dengan tambahan biaya percepatan sebesar + Rp. 65/ kata (mohon konfirmasi kembali)

Sebelum kami kirimkan kembali naskahnya, diharapkan membayar tagihan di atas. Tagihan dapat dibayarkan melalui transfer ke rekening

# Mandiri an. Agista Rosiana (1330012114542).

Apabila biaya tersebut sudah ditransfer, mohon mengirimkan bukti/struk ke alamat email ini atau no CP kami. Kelebihan jumlah transfer akan didonasikan untuk Komunitas Kongkrit.(Silahkan mengunjungi akun resmi instagram kami di @komunitaskongkrit\_ untuk mengetahui kegiatan yang sudah dilakukan oleh Komunitas Kongkrit dengan menggunakan sebagian dana Translate Sedekah. Saran Anda sangat kami butuhkan untuk Translate Sedekah yang lebih baik. Terima kasih atas kerjasamanya.

Salam, Tim TS Komunitas Kongkrit Kontak : Taufiq (+62895360173913), Hijrah (+6285299322978), Ria (+62 822-3259-1306) [Kutipan teks disembunyikan]

Mashuri Masri <mashuri.masri@uin-alauddin.ac.id> Kepada: Translate Sedekah <translate.sedekah@gmail.com>

Assalamualaikum bapak ibu.

berikut kami kirimkan bukti transfer kami memilih opsi 65/kata utk percepatan ya bapak/ibu.

terimakasih atas kerjasamanya salam hormat mashuri masri [Kutipan teks disembunyikan]

## PROOF READING MASHURI MASRI.jpg

6 April 2021 05.52

5 April 2021 22.33

7 April 2021 00.12

1/3

# 4/12/2021



Translate Sedekah <translate.sedekah@gmail.com> Kepada: Mashuri Masri <mashuri.masri@uin-alauddin.ac.id>

Mohon maaf Pak,

Untuk harga percepatan +65/kata (ditambah 65rupiah dari harga normal)

Sehingga harga proofreading percepatan 59+65 per kata= 124/kata

Mohon konfirmasinya jika Bapak hendak mengambil opsi percepatan. Kami akan mengirimkan tagihan yang baru jika demikian. Namun, jika Bapak tetap dengan harga normal, kelebihan transfer akan kami kembalikan dan mohon mengirimkam no rek Bapak.

Terima kasih

[Kutipan teks disembunyikan]

Mashuri Masri <mashuri.masri@uin-alauddin.ac.id> Kepada: Translate Sedekah <translate.sedekah@gmail.com>

assalamualaikum.

oh maaf, kalo begitu sy kembali ke harga normal ya. rekening mandiri saya : mashuri masri 1030007644624

terimakasih atas kerjasamanya

[Kutipan teks disembunyikan]

Translate Sedekah <translate.sedekah@gmail.com> Kepada: Mashuri Masri <mashuri.masri@uin-alauddin.ac.id>

Dear Bapak Mashuri,

Berikut kami lampirkan bukti transfer atas dana lebih dari transfer yang telah Bapak lakukan.

Salam, Translate Sedekah

[Kutipan teks disembunyikan]

Live and a second secon

Mashuri Masri <mashuri.masri@uin-alauddin.ac.id> Kepada: Translate Sedekah <translate.sedekah@gmail.com>

## terimakasih

[Kutipan teks disembunyikan]

7 April 2021 09.20

7 April 2021 11.04

7 April 2021 11.22

7 April 2021 11.41

Translate Sedekah <translate.sedekah@gmail.com> Kepada: Mashuri Masri <mashuri.masri@uin-alauddin.ac.id>

# Dear Bapak Mashuri Masri,

Berikut kami lampirkan hasil terjemahan/proofreading kami. Mohon diperiksa kembali.

Silahkan mengunjungi akun resmi instagram kami di @komunitaskongkrit\_ untuk mengetahui kegiatan yang sudah dilakukan oleh Komunitas Kongkrit dengan menggunakan sebagian dana Translate Sedekah.

Saran Anda sangat kami butuhkan untuk Translate Sedekah yang lebih baik. Terima kasih atas kerjasamanya.

Hormat Kami,

Tim Translate Sedekah Komunitas Kongkrit Kontak (wa/telepon): Taufiq (0895360173913) Kontak : Taufiq (+62895360173913), Hijrah (+6285299322978), Ria (+62 822-3259-1306) [Kutipan teks disembunyikan]

#### 2 lampiran

elean\_V-Novel chitinolytic bacteria from shrimp (edit1) (Pr. Silmi Rpr.Tya).doc 3193K

tracked\_V-Novel chitinolytic bacteria from shrimp (edit1) (Pr. Silmi Rpr.Tya).doc 3240K

of Bio			👗 mashuribiod
U	ncorrected Proof	×	
	articinants		Libi
ems Sr	nuio Editors (editors)		
	nica Sentiasari (acentiasari)		
D	EWI NUR PRATIWI (dewinurpratiwi)		
м	ashuri Masri (mashuribiodiversitas)		
	Messages		
	Note	From	
	Dear Author(s),	dewinurpratiwi	
	Pls find attached file for an uncorrected proof (Convedited file)	2021-04-13 09:28	
	The revised manuscript is awaited. Do not worry about layout changes due to	PM	
	revision; our staff will fix it again.		
	Note: Kindly use track change when you make improvements.		
	🗋 dewinurpratiwi, chitinolytic bacteria - Masri.doc		
	<ul> <li>Dear Ahmad dwi setiawan and team</li> </ul>	mashuribiodiversitas	8-18 02:35 AM
	I have revised the manuscript and send it in OJS.	2021-04-15 08:22	
	thank you for collaboration	AM	
	thank you for conaboration		Q Search
	🗋 mashuribiodiversitas, 8019-Article Text-40637-1-18-20210413 COPYEDITING.doc		
	Add Message		
	Revisions		
	40808-1 Article Text, FINAL ARTICLE.doc		
	▶ D 40809-1 Other, PROFESSIONAL LANGUAGE EDITOR ndf		

Review Discussions				
	dewinurpratiwi 2021-04-13 09:28 PM	mashuribiodiversitas 2021-04-15 08:22 AM	1	
	dewinurpratiwi 2021-04-13 09:30 PM	dewinurpratiwi 2021-04-19 09:28 AM	2	

Uncorrected Proof	×	
Participants		
Smujo Editors (editors)		
Anisa Septiasari (aseptiasari)		
DEWI NUR PRATIWI (dewinurpratiwi)		
Mashuri Masri (mashuribiodiversitas)		
Messages		
Dear Author(s), <sub>No</sub> Pas, find attached file for an uncorrected proof (Copyedited file). The revised manuscript is awaited. Do not worry about layout changes due to revision; our staff will fix it again. Note: Kindly use track change when you make improvements.	dewinurpratiwi 2021-04-13 09:28 मिश्रम	
🗋 dewinurpratiwi, chitinolytic bacteria - Masri.doc		
<ul> <li>Dear Ahmad dwi setiawan and team</li> <li>I have revised the manuscript and send it in OJS.</li> <li>thank you for collaboration</li> </ul>	mashuribiodiversitas 2021-04-15 08:22 AM	
mashuribiodiversitas, 8019-Article Text-40637-1-18-20210413 COPYEDITING.doc  Add Message		

l of Bio <mark>lemiant Dimensit</mark>	n Tala 🗛 🖉	allah 🦚 Mara Cho	🛔 mashuribiodiver
Uncorrec	ted proof [02]	×	
<b>Participa</b> Smujo Edit Anisa Septi Mashuri Ma	<b>nts</b> ors (editors) iasari (aseptiasari) asri (mashuribiodiversitas)		Librar
Messag	jes		
Note	,	From	
Pls, fi 🗋 ed	nd attached file for 2nd uncorrected proof itors, chitinolytic bacteria - Masri=REV+ (02).doc	editors 2021-04-20 01:28 PM	
<ul> <li>thank</li> <li>I have</li> <li>I atta</li> </ul>	c you for your uncorrected proof.	nashuribiodiversitas 2021-04-20 03:54 PM	
l" ma	ashuribiodiversitas 8019-Article Text-41095-1-18-20210420 (1) FDIT doc		10.0005 AM
Add Mes	Reviewer's Attachments	April 15,	Q Search
	Image: Ward of the second state of		
	Revisions		
	► 🛃 40808-1 Article Text, FINAL ARTICLE.doc		
	▶ 🔄 40809-1 Other, PROFESSIONAL LANGUAGE EDITOR.pdf		
	Review Discussions		

Uncorrected Proof	dewinurpratiwi 2021-04-13 09:28 PM	mashuribiodiversitas 2021-04-15 08:22 AM	1	
	dewinurpratiwi 2021-04-13 09:30 PM	dewinurpratiwi 2021-04-19 09:28 AM	2	

Uncorrected proof [02]	×
Participants	
Smujo Editors (editors)	
Anisa Septiasari (aseptiasari)	
Mashuri Masri (mashuribiodiversitas)	
Messages	
Note	From
Pls, find attached file for 2nd uncorrected proof	editors
🗋 editors, chitinolytic bacteria - Masri=REV+ (02).doc	2021-04-20 01:28 PM
thank you for your uncorrected proof.	mashuribiodiversitas
I have followed your review.	2021-04-20 03:54 PM
l attached EDITED FILE	
🗅 mashuribiodiversitas. 8019-Article Text-41095-1-18-20210420 (1) EDIT.doc	

Add Message

OIC	BILLING		×	
OPEN JOURNAL SYSTEMS	Participants			Library
	DEWI NUR PRATIWI (dewinurpratiwi)			
	Mashuri Masri (mashuribiodiversitas)			
	Messages			
	Note		From	
	Dear Author(s),		dewinurpratiwi	
	Kindly find attached an invoice for	the publication of your manuscript.	2021-04-13 09:30 PM	
	Note: Paper submitted from Dece publication fee of USD 320 (IDR 4	mber 1, 2020 will be charged a 4,500,000).		
	🗋 dewinurpratiwi, 3071.MASHURI MASI	Rl.pdf		
	<b>M</b> BNI		mashuribiodiversitas 2021-04-15 08:19	
			AM	)3-18 02:35 AM
	Transa	aksi Berhasil		Q Search
	Nomor Rekening Tujuan	*****994		
	Nama Rekening Tujuan	Ibu DEWI NUR PRATIWI		
	Tanggal Transaksi	15-04-2021		
	Waktu Transaksi	15:13:34 WIB		Unload File
	Email Penerima			Text
	Bank Tujuan	BNI		
	Nama Pengirim	MA****I MA**I		
	Nomor Rekening Pengirim	******865		
	Nominal	4.500.000,00		d discussion
	Fee	0,00		1
	Total	4.500.000,00		
	Keterangan	Mashuri masri. Novel chitinolytic bacteria from the shrimp shell processing		2

			🛔 mashuribiodiversita
BILLING		×	1
Participants			
DEWI NUR PRATIWI (dewinurpratiwi)			
Mashuri Masri (mashuribiodiversitas)			
Messages			
Note Dear Author(s)		From	
		2021-04-13 09:30	
Kindly find attached an invoice for t	the publication of your manuscript.	PM	
Note: Paper submitted from Dece publication fee of USD 320 (IDR 4	mber 1, 2020 will be charged a 4,500,000).		
🗋 dewinurpratiwi, 3071.MASHURI MASI	RI.pdf		
		mashuribiodiversitas	
NI BNI		2021-04-15 08:19 AM	
Transa	aksi Berhasil		
Nomor Rekening	*****994		
Tujuan			
Nama Bakaning			
Tujuan	IDU DEWI NOK PRATIWI		
	15 0 / 0001		
langgal Iransaksi	15-04-2021		
Waktu Transaksi	15:13:34 WIB		
Email Penerima			
Bank Tuiuan	RNI		
Dalik Tujudi	DINI		
Nama Pengirim	MA***I MA**I		
Nomor Rekening	******865		
Pengirim			
Nominal	4.500.000,00		
Fee	0,00		
Total	4.500.000,00		
Keterangan	Mashuri masri. Novel		
	chitinolytic bacteria from		
	the shrimp shell processing		

Biodiversitas Journal of Biological	l Diversity Tasks 0	🥹 English 👁 View Site 🛔 masl	uribiodiversita
OPEN JOURNAL SYSTEMS	8019 / <b>Masri et al.</b> / Nov	el chitinolytic bacteria from the shrimp shell processing waste	Library
Submissions	Workflow Publicat	ion	
	Status: Published		
		This version has been published and can not be edited.	
	Title & Abstract	Title	
	Contributors	Novel chitinolytic bacteria from the shrimp shell processing waste	
	Metadata	Abstract	
	References	<b>B</b> $I \times^{z} \times_{z} \mathscr{O}$	
	Galleys	<b>Abstract.</b> <i>Masri M, Sukmawaty E, Aditia L. 2021. Novel chitinolytic bacteria from the shrimp shell processing waste. Biodiversitas 22: 2672-2681.</i> Chitinase is an enzyme that hydrolyzes chitin (?-1,4 Nsetyl-D-glucosamine polymer). Several types of microorganisms produce chitinase that contributes to basic science and application. The role of chitinase enzyme is widely utilized in various fields, such as health as an anti-tumor agent and agriculture as biological control agent <del>s</del> in plant disease caused by fungus. The purpose of this study was to isolate novel chitinolytic bacteria from shrimp shell waste in Makassar, Indonesia, using the scatter plate method on chitin access calcertive media insubsted at 27% for 2 days. The	v
			Save

Platform & workflow by OJS / PKP