

# The Development and Evaluation of Nutritional Content of Date Biscuits as Functional Food for Female Worker

*by* Hasbi Ibrahim

---

**Submission date:** 15-Jun-2023 11:20AM (UTC+0700)

**Submission ID:** 2116397644

**File name:** he\_Development\_and\_Evaluation\_of\_Nutritional\_Content\_of\_Date.pdf (361.63K)

**Word count:** 4662

**Character count:** 23787

Research Article

# The Development and Evaluation of Nutritional Content of Date Biscuits as Functional Food for Female Worker

HASBI IBRAHIM<sup>1</sup>, SYAMSUL ALAM<sup>2</sup>, YESSY KURNIATI<sup>2\*</sup>, SUTAMARA LASURDI NOOR<sup>3</sup>

<sup>1,2</sup>SKM, M. Kes, Concentration of Occupational Health, Study Program of Public Health, Universitas Islam Negeri Alauddin Makassar, Indonesia. Address : Jalan H.M. Yasin Limpo No. 36 Samata, Gowa Regency, South Sulawesi, Indonesia.

<sup>3</sup>Sutamara Lasurdi Noor, SGz, SUN Movement Secretariat, Directorate of Public Health and Nutrition, Ministry of National Development Planning, Indonesia.

\*Corresponding Author

Email ID: [yessy.kurniati@uin-alauddin.ac.id](mailto:yessy.kurniati@uin-alauddin.ac.id)

Received: 23.11.20, Revised: 16.12.20, Accepted: 08.01.21

## ABSTRACT

Dates have many benefits for human health which therefore it is potentially to be functional food. One of the ways to increase the date consumption is to add it into a biscuit's batter. This study aimed to make date biscuits and assess the nutrient contents and its acceptance. The biscuits were made into 4 batter, namely by replacing each into 0%, 10%, 20%, and 40% Ajwa dates on the flour used to make the biscuit dough. Then, we carried out the assessment on view, nutrient contents and biscuit acceptance. The results showed that when the more increasing of the number of the added dates, then the biscuits' protein and iron were rising, while however, fat, carbohydrate and energy dropped gradually. Furthermore, the most good acceptance of the biscuits was 20% Ajwa. it is necessary to test the effectiveness of date biscuits as functional food.

**Keywords :** ajwa, biscuit, development, functional, nutrition.

## INTRODUCTION

Dates (*Phoenix dactylifera* L) are one of fruits that mostly grows in the Middle-East and North Africa. Date plants are spread throughout the world, such as Iraq, Iran, Saudi Arabia, Egypt, Tunisia, Algeria, Libya, United Arab Emirates, Bahrain and Oman (Al-Jansen, 2009; Masmoudi-Allauche, F, et al, 2011). Dates, in addition, is a monocotyle plant and in the Arecaceae family, where the tree grows in medium size, around 15-25 metres in height. The leaves are 4-6 cm, and shape sharply in the back. According to the fruit, has oval-shaped fruit with the length around 3-7 cm, and its diameter is 2-7 cm, in which the colour varies from light red to light yellow, depending on the variety. This plant is dioecious, meaning it has male and female plants separately (Ateeq A, et al, 2013).

Dates are consumed as medicine broadly and herbally to overcome some diseases (Bagherzadeh A, et al, 2020; Ali, Z., et al, 2020). This is not just a tradition passed down from generation to generation because some researches have proved it has many health benefits such as its syrup having the therapeutic effect (Teleb H, Maddocks SE, Morris RK and Kenekanian AD, 2016).

Dates are affordable enough as the source of macro and micro nutrients (Sadau SL, Bako ZM, Namadina M, Ibrahim S, 2019; Nadeem, M., et al, 2019; Hussain, M, I., M, Farooq., Q, A, Syed.,, 2020). Besides, dates has phenolic (Khallouki F, Ricarte I, Brauer A and Owen RW, 2018), B Glucan and Anthocyanin (Hussain, M, I., M, Farooq., Q, A, Syed.,, 2020) which are antioxidant and antimicrobial, and also as anticancer, hepatoprotective, neuro-protective, antidiabetic, antihyperlipidemic, improving fertility and the others as antioxidant and antimicrobial (Helfar A, Shaheen HM, Maousa JSA, 2016).

Unfortunately, while there are above mentioned benefits, the daily consumption of dates is low in many Asian countries. In order to increase the number of people consuming it, transforming the dates into something else is demanding such as biscuit because this snack is eaten almost all people worldwide (Leelavanthi, R. Sri Sangamitra, 2016). Because of it, many researchers prefer to use biscuits as a form of functional foods. Biscuits are frequently substituted with another foodstuffs to increase its nutrient contents by adding Oyster mushrooms (Proadhan, U, K., et al, 2015), spinach (Galla, N.R, P.R.Pamidinghantam, et al, 2017), apple pulp (Alongi, M., S. Melchior., M. Anese, 2019) or papaya (Pathak, R., V. Thakur and

R.K.Gupta., 2018) and even there were the biscuits added with spices such as ginger (Fabusiwa, F.J.G, et al, 2018) and turmeric (Adegoke GO, Oyenkule AO and Afolabi MO, 2017).

Dates can enhance the nutrient contents within the biscuits, especially because of high of iron, 0.1-1.5 mg/100 gram (Mohan, 2013). This would be good enough with people who suffers iron-deficiency anemia because according to a research in Jogjakarta, through consuming dates and honey, it was proven in improvement of the Haemoglobin (hb) level (Hendri, 2007). It similarly happened when performing it with animal trial (Onuh SN, et al, 2012; Pravitasari, 2014); Zen ATH, Pertiwi P, Chodidjah, 2015). Likewise, research involving 15-to-18-year-old female adolescent can boost Hb-level by giving them dates (Ali S, Alam G, Samrichard, 2020).

Therefore, based on above mentioned reason, we aimed to make biscuit dough and assess the nutrient contents and its acceptance as functional food for female worker.

## MATERIALS AND METHODS

### Ingredients:

Ingredients used on the dates biscuits were Ajwa dates, wheat flour, eggs, butter, powdered sugar, baking soda, vanilla, liquid milk, and salt where those were compiled from the local market. The Ajwa dates were chosen because it was suggested to boost the female adolescents' Hb-level (Ali S, Alam G, Samrichard, 2020), whereas the basic biscuit dough used referred to pumpkin biscuit dough (Syam, A., et al, 2019).

### Biscuit making:

The biscuits were made in the Culinary Laboratory Faculty of Public Health Universitas Hasanuddin and the ingredients were on table 1. The biscuit-making procedure was the same as how to make biscuit from pumpkin seeds (Syam, A., et al, 2019). There were 4 types of biscuit batter as well as the control, 10%-ajwa, 20%-ajwa, and 40%-ajwa. Each dough was made by substituting flour used with 0%, 10%, 20%, and 40% in total. The analysis of nutrient contents, physical appearance, and organoleptic test were performed by using the fresh-from-the-oven date biscuits at the same day.

### Nutrient contents analysis:

This analysis was performed in Terpadu laboratory, Faculty of Animal Husbandry, Universitas Hasanuddin. The determination of carbohydrate, fat, and protein contents used Luff-Schoorl, Khejdahl, and Soxhlet methods, respectively, while the determination of iron contents used XRF (X-Ray Fluorescence) method.

### Organoleptic test:

Due to Covid-19 pandemic, this test has been done by involving the faculty staff in the Faculty of Public Health, Universitas Hasanuddin. The panelists were 15 people, consisting of 8 women and 7 men with 25-45 years old. There were 4 types of biscuits tested hedonically with 5-level scales of pleasure, namely from really like (5) to really dislike (1). Moreover, those untrained panelists were given questionnaires to assess the 4-type date biscuits based on colour, aroma, taste, and texture. The panelists did not know which ingredient was in the served biscuit (single blind design) and we labelled the sample with random code.

### Statistical analysis:

Data were analysed non-parametrically by performing Kruskal-Wallis test and in terms of finding out the score of each product, we analysed the data from a Likert scale.

## RESULTS

### Biscuit characteristics:

This study made date biscuits with 4 doughs, which fulfill the criteria namely the control biscuit (without Ajwa dates), A10% (10% flour : 30 gram Ajwa dates), A20% (20% flour : 60 gram Ajwa dates), and A40% (40% flour : 120 gram Ajwa dates). The biscuits were attractive and had after-baking deliciousness. It was shown when the more dates added, the colour was darker, the texture was harder, and the taste was sweeter. This is considered because the characteristic of Ajwa dates, black in colour and sweet in flavor (Table 2)

### Nutrient contents:

From laboratory assessment, we found the protein in the biscuit was more increasing by addition of the dates. In comparison to the protein, fat and carbohydrate were going to decrease during addition of the Ajwa dates. Iron (Fe) in the biscuits was more increasing along with the addition of Ajwa dates by 1.785-7.244 mg per 100 gram biscuit (Table 3).

### Organoleptic test:

The results of the organoleptic test are provided in table 4. According to the data, A20 had the highest score along with control biscuits while increase in percentage of dates up to 40% decreased the acceptability of the biscuits.

## DISCUSSION

### Biscuit characteristics:

Ajwa is a type of date that is the most expensive in comparison to other types. Ajwa is stated on a Hadith from the Messenger as a special date. This date contains 0.31±0.44 gram fat, 3.1±0.1 gram proteins, 3.4±0.2 carbohydrates and 100 gram/kcal energy (Hamid, NAA., et al 2020).

Adding certain foodstuffs into the dough can indeed affect the biscuit characteristics. Because of the nutrient contents within the dates, the nutrients inside the date biscuits are similar. Commercially, the mixed biscuits are cheaper than the multigrain biscuits (Anwar, H., K, Rajkumari., B, Anju., 2018). The papaya addition was also able to make biscuits to be functional food because it has Papain enzyme that can facilitate digestion (Pathak, R., V. Thakur and R.K.Gupta., 2018, 7(4))

#### Nutrient contents:

From laboratory assessment, we found the protein in the biscuit was more increasing by addition of the dates (table 3). On the other hand, one study showed dates are not the resource of protein where there is just only 2.08%-3.1% protein in dates. However, the assumption was because the dates used were Ajwa, as a result the protein content was fairly good. The reason is because the Ajwa dates have a higher level of protein than other types of dates (Omawunmi SA and Ayoade LA, 2015).

In comparison to the protein, fat and carbohydrate were going to decrease during addition of the Ajwa dates. This implicated the total energy within the biscuits. Hence, the more dates added to the dough, the total energy decreased. In flour form, dates were constructed of 1.96% protein, 0.5% fat, 74.98% carbohydrate, and 9.05% fibre (Dhaakhar, J, Vahistha N, and Sharma A, 2019), however when it came to the biscuits in this study, the macronutrients were 6.36-10.11% of protein, 9.02-14.64% fat, 65.23-71.7% carbohydrate and <sup>19</sup>2.54-444.0 kcal. This contrast was most likely due to the ingredients used.

Iron (Fe) in the biscuits was more increasing along with the addition of Ajwa dates by 1.785-7.244 mg per 100 gram biscuit. One of the studies showed the Fe within dates was 0.25-5.93 mg/100 gram (Khan, A.A., et al, 2019). Another study revealed the Fe contents in dates were in between from 0.46 to 2.52 mg/100 gram. The main function of Fe in blood is to contribute in transporting oxygen throughout the <sup>12</sup>body. The sustained Fe intake recommended is 10 mg/day for man and 15 mg/day for woman. In addition, the dates can act as a supplement of Fe in order to meet the daily-intake need (Nadeem, M., et al, 2019).

Biscuit was experimented in propose to increase the nutrient. For instance, one experiment showed when biscuit was combined with Oyster mushroom, it also significantly improved the protein (Korus, AD, Gumul, M., Krystyjum, L.J.J., Korus., 2017). In line with this, substituting certain foodstuff to the dough can increase the mineral,

such as by adding Psyllium powder, the mineral of the biscuits was increase (Krystyan. M., D. Gamal, A. Korus, J. Korus, and M. Sikora, 2018). Likewise, the addition of spinach powder could improve the mineral (Calcium, Fe, and Phosphor) and also anti<sup>16</sup>oxidant (Galla, NR, PR Pamidinghantam, et al, 2017). On the same hand, fortification can increase the fiber and antioxidant. Rhubarb addition can improve the level of phenolic in biscuits as well as decreasing the probability to suffer with free-radicals (H. Dogan and R Merral, 2019). The addition of C. Racemosa can increase the biscuit nutrients such as protein, fibre, phenolic, and antioxidant (Kumar, A., E. Krishnamoorthy., H.M. Devi, et al, 2017). The biscuits made by the mixture of ginger and pigeon pea have the total of phenolic and flavonoid as antioxidant and high antivitax (Fabusiwa, F.J.G, et al, 2018). The use of tea <sup>15</sup> can improve the fibre content within the biscuits (Soma G, Mahadevamma, S., Sudha M.C., 2016). Besides, the addition of turmeric evidently <sup>14</sup> affects the level of antioxidant of the biscuits (Adegoke, G.O., Oyenkule, A.O., and Afolabi M.O, 2017). In addition, the material substitution can lower the glycemic index of the biscuits such as the addition of apple pulp which can show the decrease of the glycemic index of the biscuits (Alongi, M. S. Melchior., M. Anese, 2019). The biscuits made by composite flour are rich in crude fibre, protein, and amylase, however poor in the glycemic index and sugar (J.U. Obaroakpo, I. Iwanegbe, A. Ojokuh, 2017). Meanwhile, there is an addition of specific substance but it does not increa<sup>10</sup> the nutrient content of biscuits. For example, the addition of PSC flour did not affect any fat content if compared to control biscuits (Proadhan, U, K., et al, 2015).

#### <sup>9</sup> Organoleptic test:

The results of the organoleptic test are provided in table 4. According to the data, A20 had the highest score along with control biscuits while increase in percentage of dates up to 40% decreased the acceptability of the biscuits. This addition was significantly seen in its texture, in which if the concentration was added, the texture was getting harder. This was confirmed when looking at the physical appearance of the biscuits that was indeed less crisp. Based on colour, aroma, and taste, A20 was the biscuits having the highest score. This shows that the total of dates added which was accepted by the panelists was 20%. In comparison, A40 was the lower score in all aspects as the addition until 40% turned out to make the colour darker, the aroma stronger, the taste sweeter, and the texture harder. These

caused A40 to be not really excited amongst panelists.

The increase of contents within the date flour render the colour to be darker. It is also stated biscuits with added dates significantly had darker in colour in comparison to the control biscuits (Amany, M.S. & H. A. Hussain, 2017). Darker colour was resulted by caramelising process (Ndife J., Abdulraheem Lo, Zakaria UM, 2011). Relating to the hard texture, the decrease of crispness is caused by the amount of fibre within the dough (El-Sharnoby, P.A., Aleid SM, Al Otaibi MM, 2012; Arun, KB, F. Persia, P.S. Aswathy, J. Chandran, M.S. Sajeef, P. Jayamurthy and P. Nisha, 2015). The strong hydrophilic component will absorb more water causing the biscuits to be more tough (Korus, A, D.Gumul, M. Krystyjum, L.J.J. Korus., 2017). The high fibre content causes the date biscuits to be more tough if compared to control biscuits (Becker FS, Daminai C, de Melo AAM, Borges PRS, 2014).

This finding becomes a dilemma pondering about that A40 had the highest Fe and protein than other experimented biscuits. Nevertheless, if it was less interested organoleptically, it would be hard to judge A40 as the best product. Despite this, these still need to be enlightened by an extent research involving the well-trained panelists, in which indeed have qualified competency better in assessing the sensory properties of a product.

#### CONCLUSION

This study found the dates could be made to be functional biscuits with high of Fe. The biscuits made had both protein and Fe contents that increased along with the addition of used dates. Biscuits with the best acceptance were A20 whereas A40 showed the highest protein and Fe levels, yet on the other hand, it had the lowest score in organoleptic tests. There is a need of trained panelists in terms of performing the organoleptic test also. Likewise, this study suggests to conduct further research on the effectiveness of the date biscuits in improving the hemoglobin level among anemia patients.

#### ACKNOWLEDMENT

The author would like to thank the Institute for Research and Community Service, Alauddin State Islamic University Makassar, which has provided financial support for the implementation of this research.

#### CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

#### REFERENCES

1. Adegoke, GO., Oyenkule AO and Afolabi MO. Functional biscuits from wheat, soya bean and turmeric (*Curcuma longa* : optimization of ingredient levels using response surface methodology). *Research J. of Food and Nutr* 2017; 1(1): 13-22.
2. Ahmad A, Soni DS, Singh V, & Maurya KS. *Phoenix dactylifera linn* : a review. *Int. J. of Research in Aryveda & Pharm* 2013; 4(3), 477-51.
3. Ali E-F, A, Hazeem S, Mohammad AD, Soad AJ, et al. Date palm (*Phoenix dactylifera*) protection and remedy food. *Current Trends in Nutraceuticals* 2016; 1(9).
4. Ali S, Gemini A, Samrichard. Ajwa date fruit (*Phoenix dactylifera l*) in increasing haemoglobin (hb) level to teenage girl. *Enfermeria Clin* 2020; 30 (2), 77-79.
5. Alireza B, Ashgar E, Arman Z, Mojgan M, Sayed MBF, Mustafa A, Navid RB. Clinical effects of date palm (*phoenix dactylifera l*) : a systematic review on clinical trials. *Compl. Therapies in Med* 2002; 51, <https://doi.org/10.1016/j.ctim.2020.102429>.
6. Amany MS and Hana H. Nutritional of gluten free biscuits supplemented with sweet chickpeas and date palm powder. *Int. J. of Food Sci. And Nutr* 2017; 2(1), 128-134.
7. Aminuddin S, Zaenal, Yessy K, Nur AA, Indrah PW, Marini A. Development and biochemical analysis of pupmkin seed (*Cucurbita moschata dorch*) biscuit. *Pak. J. of Nutr* 2019; 18(8), 743-746.
8. Anna K, Dorota G, Magdalena K, Leslaw J and Jaroslaw K. Evaluation of the quality, nutritional value and antioxidant activity of gluten-free biscuits made from corn-acorn flour or corn-hemp flour composites. *Eur. Food Res. and Tech* 2017; 243, 1429-1438.
9. Anuj K, Elavarasan K, Hanjaban MD, Devanand U, et al. Influence of sea grapes (*Caulerpa racemosa*) supplementation on physical, functional and antioxidant properties of semi-sweet biscuits. *J. of Appl. Phy* 2017; 30, 1393-1403.
10. Anwar H, Rajkumari K, Anju B. Development of healthy multigrain biscuits from backwheat-barley composite flours. *Asian J. of Dairy and Food Res* 2018; 37 (2), 10.18805/ajdfr.DR-1328.
11. Arun KB, Florene P, PS Aswathy, Janu C, Sajeew MS, Jayamurthy P, and Nisha P. Plantain pee-a potential source of antioxidant dietary fibre for developing functional cookies. *J. Food Sci Tecnol* 2015; 52 (10), 6355-6364.
12. Asif K, Manzoor IK, Rukhsana J, Saadullah KL and Yasir A. Study of trace elements in fruits of dates at turbat-punjugur region of balochistan with reference to medicinal and environmental pollution. *Pakistan J. of Botani* 2019; 51(5),

- 1769-1774, doi : https://dx.doi.org/10.30848/PJB2019-5(4).
13. Faiza M-A, Radhia G, Walid K, Mohamed NS, Hajer J. Investigation of genetic variability related to the in vitro floral hermaphroditism induction in date palm (*Phoenix dactylifera*). *Afr. J. of Biotech* 2011;10, 10567-10574.
  14. Farid K, Irvila R, Andrea B, Robert WO. Characterization of phenolic compounds in mature moroccan medjool dates palm fruits (*Phoenix dactylifera*) by HPLC-DAD-ESI-MS. *J. of Food Comp. and Anal* 2018; 70, 63-71, doi : https://doi.org/10.1016/j.jfca.2018.03.005.
  15. Fernanda SB, Clarissa D, Adriane AMDM, Paulo RSB, Eduardo VDBV Incorporation of buriti endocarp flour in gluten-free whole cookies as potential source of dietary fiber. *Plant Foods for Human Nutr* 2014; 69(4), 344-350.
  16. Funmilayo JG-F., Oladele EP, G. Oboh, S.A. Adefegha, A. A. Oshodi. Nutritional properties, sensory qualities and glycemic response of biscuits produced from pigeon pea-wheat composite flour. *J. of Food Biochem* 2018; 4, doi : https://doi.org/10.1111/jfbc.12505.
  17. Funmilayo JG-F., Oladele EP, G. Oboh, S.A. Adefegha, A. A. Oshodi. Polyphenol Contents and Antioxidant Activities of Biscuits Produced From Ginger-enriched Pigeon Pea-Wheat Composite Flour Blends. *J. of Food Biochem* 2018; 42(4), https://doi.org/10.1111/jfbc.12526.
  18. Gamal AE, Salah MA, Mutlaq MA. Nutritional quality of biscuits supplemented with wheat bran and date palm fruits (*Phoenix dactylifera* L). *Foods and Nutr. Sci* 2012; 3(3), 322.
  19. Hafsa D and Raciye M. The effect of locust bean gum and rhubarb on the physical and functional properties of the gluten-free biscuits. *Italian J. of Food Sci* 2019; 31(3), doi.org/10.14674/ijfs-1293.
  20. Haider MS, Iqar AK, Muhammad J, et al. Profiling of date fruits (*Phoenix dactylifera*) during different fruit maturation phases. *Pak. J. Bot* 2018; 50(3), 1069-1076.
  21. Hajer T, Sarah EM, Keith M, Ara DK. Chemical characterisation and the anti-inflammatory, anti-angiogenic and antibacterial properties of date fruit (*Phoenix dactylifera* L). *J. of Ethnopharmacol* 2016; 194, 457-468, doi : https://doi.org/10.1016/j.jep.2016.10.032.
  22. Hasker, Leelavanthi K, Radhai S, Asok M. Effect of differently treated soyabean flour on quality of biscuits. *Int. J. of Res. in Advent Tech* 2016; 4(4), 9637.
  23. Hendri F. Effect of giving dates (*Phoenix dactylifera*) and honey (*Apex dorsalis*) on hemoglobin levels in the 16-18 year age group. Yogyakarta: Digilib FK Universitas Muhammadiyah Yogyakarta. 2007
  24. Joel N, Abdulraheem, Zakaria. Evaluation of the nutritional and sensory quality of functional bread produced from whole wheat and soy bean flour blends. *Afr. J. of Food Sci* 2011; 5(8), 466-472.
  25. Jyotika V, Neha V and Asha S. Development of biscuit by partial substitution of refined wheat flour with chickpea flour and date powder. *J.of Microbiology, Biotech. and Food Sci* 2019; 8(4), 1093-1097.
  26. Magdalena K, Dorota G, Anna K, Jaroslaw K, Mare S. Physicochemical properties sensory acceptance of biscuits fortified with Plantago psyllium flour. *Emirates J. of Food and Agri* 2018; 30(9), 758-763, doi : 10.9755/ejfa.2018v30.19.1799
  27. Marilisa A, Sofia M and Monia A. Reducing the glycemic index of short dough biscuits using apple pomace as a functional ingredient. *LWT* 2019; 100, 300-305, https://doi.org/10.1016/j.lwt.2018.10.068.
  28. Mohammed SAJ. Composition of date fruit (*Phoenix dactylifera* L) from offshoots and cultured cells at different Stage. *J. of Food and Tech* 2019; 7 :102-105.
  29. Mohan, J. Health benefits of date palm : phytochemicals and their functions. Helsinki: Department of Agricultural Sciences : University of Helsinki.
  30. Muhammad IH, Muhammad F, Qamar AS Farooq. Nutritional and biological characteristics of the date palm fruit (*Phoenix dactylifera* L) : a review. *Food Bioscience* 2020; 34, https://doi.org/10.1016/j.fbio.2019100509.
  31. Muhammad N, Tahir M, Ilker U, Mian NR, Qurat UA. Mineral, vitamin and phenolic contents and sugar profiles of some prominent date palm (*Phoenix dactylifera*) varieties of Pakistan. *Pak. J. Bot* 2019; 51 (1), 171-178, DOI : 10.30848/PJB 2019-1(14).
  32. Narsing RG, Prabhakara RP, Balaswamy, Math RG, Satyanarayana A Nutritional, textural and sensory quality of biscuits supplemented with spinach (*Spinacia oleracea* L). *Int. J.of Gastronomy and Food Sci* 2017;7,20-26, https://doi.org/10.1016/j.ijgfs.2016.12.003.
  33. Nur ABH, Nur HM, Maulidiani, Ahmed M, Intan SI, et al. Quality evaluation of the physical properties, phytochemicals, biological activities and proximat analysis of nine Saudi date palm fruit varieties. *J. of The Saudi Societies of Agr. Sci* 2020; 19, 151-160.
  34. Obaroakpo, Iwanegbe, Ojokuh. The functional and sensory evaluation of biscuits produced from wheat, defatted soybean and coconut flour. *Current J. of Applied Sci. and Tech* 2017; 23(6), 1-7,
  35. Omawunmi SA and Ayoade LA. Nutritional composition of the fruit of the nigerian mild date palm (*Phoenix dactylifera*). *World J. Dairy Food Sci* 2015; 8: 196-200.
  36. Onuh, Ukaejiifo, Peter A, Silas U, Chukwugozie O, et al. Haemopoietic activity and effect of

- crude fruit extract of phoenix dactylifera on peripheral blood parameters. BiomedSci Direct Publication 2012; 3 (2), 1720-3.
37. Pravitasari. The effect of date palm extract on blood hemoglobin in vitro in male white rats. Surabaya: FK Airlangga. 2014
  38. Rupanjali P, Vandana T, Rajinder KG. Formulation and analysis of papaya fortified biscuits. J. of Pharmaco. and Phytochem 2018; 7(4), 1542-1545.
  39. Salisu BS, Zainab MB, Murtala MN, Sani I. Comparative proximate analysis of some date palm cultivars (Phoenix dactylifera l). FUDMA J. of Sci 2019; 3 (4).
  40. Sediaoetama A. Nutrition Science for Students and Professions in Indonesia. Jakarta: Dian Rakyat. 2010
  41. Soma G, Mahadevamma S, Sudha MC. Characterisation of tea fiber and its utilisation as an functional ingredient in the preparation of biscuits. Int. Food Res. J 2016; 3(6), 2525-2533.
  42. Utpal KP, Khan MMRL, Fardos A-A, Jahangir A. Development and quality evaluation of mushroom (Pleurotassajor-caju) enriched biscuit. Emirates J. of Food and Agri 2015; 27(7), 542-547, doi : 10.9755/ejfa.2015.04.082.
  43. Winarno. Food chemistry and nutrition. Jakarta: Gramedia Pustaka Utama. 1999
  44. Zen ATH, Pertiwi P, Chodidjah. The effect of giving date palm juice (Phoenix dactylifera) on hemoglobin levels: an experimental study on male white rats Wistar strain. Sains Med 2015; 5(1) 17-9.
  45. Zeshan A, Jianke L, Yuhuan Z, Naureen N, et al. Dates (Phoenix dactylifera) and date vinegar : preventive role against various disease and in vivo mechanism. Food Reviews Int 2020; <https://doi.org/10.1080/875591292020.1735411>.

**Table 1: Recipe of the date biscuit and the control biscuit**

Ingredients	Control	A 10%	A20%	A40%
Flour (gram)	300	270	240	180
Ajwa Dates (gram)	0	30	60	120
Egg	1	1	1	1
Powdered Sugar (gram)	200	200	200	200
Butter (gram)	170	170	170	170
Liquid Milk (ml)	100	100	100	100
Vanilla (gram)	4	4	4	4
Baking soda (gram)	3	3	3	3
Salt (gram)	3	3	3	3

**Table 2: The date biscuit characteristics**

Trials	Ajwa Flour :	Profil Biscuits
Control	0:100	Golden brown in colour, interesting, crispy in texture, baked evenly, tasty in aroma and taste
A10%	10:90	Slightly black, interesting, crispy, well-baked, the distinctive aroma of dates
A 20%	20:80	More black, interesting, quite tough and less crisp, well-baked, the distinctive aroma of dates
A 40%	40:60	Black in colour, interesting, more tough and less crisp, well-baked, the distinctive aroma of dates but stronger and sweeter

**Table 3: The nutrients in 100 gram of date biscuits**

	Protein (gram)	Fat (gram)	Carbohydrate (gram)	Energy (Kkal)	Fe (mg)
Control	4.35	14.42	71.16	431.82	1.103
A10%	6.36	14.64	71.7	444.00	1.785
A20%	8.82	13.09	70.07	433.37	3.511
A40%	10.11	9.02	65.23	382.54	7.244

**Table 4: The preference score among untrained panelists towards each type of biscuits**

Sample	Colour	Aroma	Taste	Texture	Total Score
Control	57	54	54	52	54.25
A10	51	54	52	49	51.5
A20	57	57	56	47	54.25
A40	48	36	38	26	37



# The Development and Evaluation of Nutritional Content of Date Biscuits as Functional Food for Female Worker

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

17%

INTERNET SOURCES

6%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.ijpronline.com">www.ijpronline.com</a> Internet Source	8%
2	<a href="http://repo.ur.krakow.pl">repo.ur.krakow.pl</a> Internet Source	2%
3	Submitted to Chabot-Las Positas CCD Student Paper	1%
4	<a href="http://journal.uin-alauddin.ac.id">journal.uin-alauddin.ac.id</a> Internet Source	1%
5	<a href="http://www.ejfa.me">www.ejfa.me</a> Internet Source	1%
6	<a href="http://repo-dosen.ulm.ac.id">repo-dosen.ulm.ac.id</a> Internet Source	1%
7	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1%
8	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	<1%
9	Submitted to Universitas Indonesia Student Paper	<1%

10	<a href="http://ejfa.me">ejfa.me</a> Internet Source	<1 %
11	<a href="http://irep.iium.edu.my">irep.iium.edu.my</a> Internet Source	<1 %
12	<a href="http://www.thehindu.com">www.thehindu.com</a> Internet Source	<1 %
13	Zeshan Ali, Jianke Li, Yuhuan Zhang, Naureen Naeem, Shoaib Younas, Farah Javeed. " Dates ( ) and Date Vinegar: Preventive Role against Various Diseases and Related Mechanisms ", Food Reviews International, 2020 Publication	<1 %
14	<a href="http://tech.ui.edu.ng">tech.ui.edu.ng</a> Internet Source	<1 %
15	<a href="http://ir.cftri.res.in">ir.cftri.res.in</a> Internet Source	<1 %
16	Hadjer Chenini Bendiab, Noureddine Djebli, Yakub Kara, Meltem Uçar, Sevgi Kolayli. "An Investigation of Algerian Dates (Phoenix dactylifera L.); Antioxidant, Anti-inflammatory Properties and Phenolic Compositons H", Emirates Journal of Food and Agriculture, 2021 Publication	<1 %
17	Heba I. Mohamed, Hossam S. El-Beltagi, S. Mohan Jain, Jameel M. Al-Khayri. "Date palm	<1 %

(*Phoenix dactylifera* L.) secondary metabolites: Bioactivity and pharmaceutical potential", Elsevier BV, 2021

Publication

---

18

Mildner-Szkudlarz, Sylwia, Joanna Bajerska, Paweł Górnaś, Dalija Segliņa, Agnieszka Pilarska, and Teofil Jesionowski. "Physical and Bioactive Properties of Muffins Enriched with Raspberry and Cranberry Pomace Powder: A Promising Application of Fruit By-Products Rich in Biocompounds", *Plant Foods for Human Nutrition*, 2016.

Publication

---

<1 %

19

[www.suaire.sua.ac.tz](http://www.suaire.sua.ac.tz)

Internet Source

---

<1 %

20

[dergipark.org.tr](http://dergipark.org.tr)

Internet Source

---

<1 %

21

Al-Farsi, Mohamed, Cesarettin Alasalvar, Anne Morris, Mark Baron, and Fereidoon Shahidi. "Compositional and Sensory Characteristics of Three Native Sun-Dried Date (*Phoenix dactylifera* L.) Varieties Grown in Oman", *Journal of Agricultural and Food Chemistry*, 2005.

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On