

Environmental Pollution and Health Problems Due to Forest Fires with Co2 Parameters

by M. Fais Satrianegara

Submission date: 27-Apr-2022 10:05PM (UTC+0700)

Submission ID: 1821919872

File name: ojsadmin,_152.pdf (439.37K)

Word count: 2504

Character count: 13500

Environmental Pollution and Health Problems Due to Forest Fires with Co2 Parameters

Anwar Mallongi¹, Rusydi Indra², Muh. Kamil Muh. Arief², M. Fais Satrianegara³

¹Senior Lecturer of Environmental Health Faculty of Public Health, University of Hasanuddin, ²Postgraduate student of Public Health, University of Hasanuddin, Makassar, ³Faculty of Medicine and Health Sciences, Alauddin State Islamic University, Makassar

Abstract

Today, air pollution is a serious problem faced by industrial countries. The impact caused by air pollution is not only a direct impact on human health, but also can damage the environment. One of the sources of air pollution are now common is the forest fires which generate a compound emissions are carbon dioxide (CO₂). Several studies in the medical literature documenting a causal link between air pollution, especially CO₂ and negative impacts on health after short-term exposure, especially the unknown long-term health consequences of exposure to pollution. Some studies related to fire management models and CO₂ emission reductions of them use the method of mapping risk of forest fires and the use of reforestation and the elimination of illegal logging is considered effective to implement.

Keyword: Air pollution, forest fires, carbon dioxide

Introduction

Today, air pollution is a serious problem faced by industrial countries. Air pollution raises a very adverse impact. The impact caused by air pollution is not only a direct impact on human health, but also can damage the environment.

One of the sources of air pollution are now common is their forest fires. One of emissions contained in air pollution as a result of the forest fires are compounds of carbon dioxide (CO₂).¹

CO₂ emissions and global warming potential to be one of the considerations biggest environmental problems today. Due to the increase global population and industrial development quite rapidly result in the concentration of CO₂ in the atmosphere has grown an average of 2 ppm / year for the last 10 years.²

Materials and Method

In this literature review, keywords used to get a reference to a related article is "air pollution", "forest fires" and "carbon dioxide". The scope of the literature review will be limited beginning in 2011 to the top with the goal of search engines can produce journals - journals

and references are more focused on development problems and solutions. Reference is also limited in the famous journal, namely Science Direct, because the journal can be accessed free of charge. Search results using keywords found about 6,200 research articles and 7,800 books. However, based on the criteria of the carbon dioxide, only 12 articles were reviewed,


Result and Discussion

Air Pollution, Forest Fire and Carbon

Air pollution can be defined an activity in the atmosphere, where the concentration of the substance - a substance that is sufficiently high and is above the ambient value and can have an impact - the impact on humans, animals, vegetation, or material. In addition to disrupting the aesthetics, small-sized particles in the air can be inhaled into the respiratory system and cause respiratory illness and lung damage. Particles that get into the respiratory system will be set aside depending on the diameter.³

One of the sources of air pollution are now common is their forest fires. One of emissions contained in air pollution as a result of the forest fires are compounds CO₂.⁴

CO₂ is actually a component compound that naturally exist in the air. Effect of CO₂ is called the greenhouse effect of CO₂ in the atmosphere which can absorb heat energy and obstruct the course of the heat energy from the atmosphere to the surface is higher.⁽⁵⁾



Picture 1

Cycle greenhouse gas

Transport, transfer and accumulation of CO₂

The oceans play an important role for the distribution of this is because the CO₂ concentration at the level of atmospheric CO₂ that is controlled by the concentration of CO₂ in the oceans. Cycle CO₂ in the oceans separated in two senses, namely CO₂ natural and anthropogenic CO₂. Since the increase of anthropogenic CO₂ in nature, the sea has a role as an absorber of anthropogenic CO₂ to maintain the balance of the Earth system. On the other hand, the increase in this concentration in the feared sea will also affect the balance mechanism existing systems in it. Therefore, the amount of CO₂ that accumulates in the water column and how its distribution among the oceans of the world, will continue to be examined from year to year due to increased concentrations of anthropogenic CO₂ in the atmosphere.⁶

CO₂ in the atmosphere is also absorbed by plants through photosynthesis berhijau leaves, but the amount of CO₂ available to that used by plants on the earth is not balanced anymore. The more CO₂ and other greenhouse gases in the atmosphere, the more infrared radiation is

absorbed, the higher the intensity of the greenhouse and consequently the temperature at the earth's surface is also higher.⁵

Diffusion of CO₂ from tissues to the bloodstream and lungs are also caused by the pressure difference of CO₂. CO₂ pressure in tissues, veins, arteries, and alveoli. Therefore, the CO₂ in the network will be transported to the alveoli in the lungs. Literature and scientific documents relating to CO₂ inhalation exposure effects on human health and psychomotor performance, linear physiological changes in the circulatory system, cardiovascular and autonomic on exposure to CO₂ concentrations in the range of 500-5000 ppm. Human experimental studies explain that short-term exposure to CO₂ begins at 1000 ppm affect cognitive performance including decision-making and problem solving. Changes in the autonomic system for low-level exposure to CO₂ may involve this effect. Further research is needed on the long-term effects of exposure to low levels of CO₂ in the autonomous system. A number of epidemiological studies show a link between low-level exposure to CO₂ ranging from 700 ppm and symptoms associated with the building. Respiratory symptoms indicated in children who are exposed to indoor CO₂ concentrations

higher than 1000 ppm. However, in other comorbidities pollutants may be involved in these effects.⁶

The amount of CO₂ that accumulates in the water column and how its distribution among the oceans of the world, will continue to be examined from year to year due to increased concentrations of anthropogenic CO₂ in the atmosphere.⁷

Impact on the environment

Population growth and the use of fossil fuels at the same time as the main energy source in the region has led to the release of more CO₂ into the atmosphere with negative consequences on the environment.⁸

CO₂ is a greenhouse gas implicated in global warming trend. Snow and ice have declined closure, the ocean temperature has increased and the level of the ocean's surface also increased over the last century. Rising sea levels could submerge many areas. Extreme weather conditions are causing droughts, floods and typhoons, as well as the distribution of disease-causing organisms may occur predictability.⁹

Health impacts caused

Several studies in the medical literature documenting a causal link between air pollution and negative health outcomes immediately after exposure, much less is known long-term health consequences of exposure to pollution. A study using Indonesia's forests in 1997 as a natural experiment, which estimates long-term effects of air pollution on health outcomes that take advantage of the properties of longitudinal from Indonesia Family Life Survey (IFLS), by collecting individual data detail about many health outcomes, both in Discovered in 1997 and 2007. the significant negative impact of pollution, which survive in the long term. Men and older people most impacted, while the children seemed to recover almost completely from this initial shock. For the whole population,¹⁰

Judging from some of the literature and scientific documents relating to CO₂ inhalation exposure effects on human health and psychomotor performance. Linear physiological changes in the circulatory system, car-diovaskular, and autonomous exposure to CO₂ concentrations in the range of 500 to 5000 ppm. Human experimental studies have suggested that exposure to

short-term CO₂ begins at 1000 ppm affect cognitive performance including decision-making and problem solving. Changes in the autonomic system for low-level exposure to CO₂ may involve this effect. Further research is needed on the long-term effects of exposure to low levels of CO₂ in the autonomous system. A number of epidemiological studies show a link between low-level exposure to CO₂ ranging from 700 ppm and symptoms associated with the building. Respiratory symptoms indicated in children who are exposed to indoor CO₂ concentrations higher than 1000 ppm.

However, in other comorbidities pollutants may be involved in these effects. In the context of a significant linear increase of the global ambient CO₂ concentrations caused by anthropogenic activities and resources, reduce CO₂ levels in the room through the vents to the ambient air is increased energy consumption in air-conditioned buildings. To control CO₂-efficient energy in the building from the surrounding air, the rise in atmospheric CO₂ concentration needs to be suppressed. In the context of a significant linear increase of the global ambient CO₂ concentrations caused by anthropogenic activities and resources, reduce CO₂ levels in the room through the vents to the ambient air is increased energy consumption in air-conditioned buildings.⁶

Handling Model That Can Be Applied

One of the research related to overcoming the problem of forest fires by using Data Base Inventory of Forest Resources, based on four aspects topographical factors, human activities, climate and forest characteristics. A conceptual framework for mapping the risk of fire is developed by integrating Database Forest Resource Inventory. Potential risks of forest fires and map the risk zone in the Chinese heritage sites was assessed using GIS.¹¹

Research presents the results of a study of absorption and reduction of carbon dioxide emissions with the use of ecosystem services in the nature reserve (reforestation) and the elimination of illegal logging in the state of the newly formed ijevan, located in Tavush region of the Republic of Armenia.¹²

Resistance

In general, the fires caused by three major factors

that fuel conditions, weather and culture, including the lack of public awareness of the importance of forest preservation. All three of the above are causing difficult implemented regulations on forest fires.⁴

Success

Improvements in the estimation of the risk of forest fires and fire risk zone mapping is essential to reduce the negative impact of fires and to facilitate the planning of the protection of forested areas. The success of the method the estimated risk of forest fires and fire risk zone mappings simply and reliably to assess the potential risk of forest fires and map the forest fire zones in a place with little historical data about the fire. This method helps provide reference for decision makers to prioritize areas for conservation of cultural resources and / or culture, and enhance the practical application of fire prevention and suppression.¹¹

Meanwhile, the results of studies related to the absorption of CO₂ emissions was found that forest ecosystems asylum state ijevan 440.7 metric can absorb carbon and can reduce CO₂ emissions each year. And the elimination of illegal logging, the number could increase carbon sequestration. As a result the amount of carbon dioxide in the atmosphere will be reduced 39.07 metric. Research on CO₂ reduction with reforestation system and the elimination of illegal logging are found it turns out the value of carbon sequestration forest ecosystem services by lead economic benefits will be surplus.¹²⁻¹⁶

Ethical Clearance- Taken from university ethical committee

Source of Funding- Self

Conflict of Interest – Nil

References

1. Ueyama, M., Iwata, H., Nagano, H., Tahara, N., Iwama, C., & Harazono, Y.. Carbon dioxide balance in early-successional forests after forest fires in the interior of Alaska. *Agricultural and Forest Meteorology*, 2019; 275 (April), 196-207. <https://doi.org/10.1016/j.agrformet.2019.05.020>
2. Zhang, H., Bucior, BJ, & Snurr, RQ. Molecular Modeling of Carbon Dioxide Adsorption in Metal-Organic Frameworks. *Modeling and Simulation in the Science of Micro- and Meso-Porous Materials*. Elsevier Inc. 2017; <https://doi.org/10.1016/B978-0-12-805057-6.00004-1>
3. Vallero, D. *The Science of Air Pollution. Fundamentals of Air. Pollution*. 2014. <https://doi.org/10.1016/b978-0-12-401733-7.00003-7>
4. Rasyid, F. Permasalahan dan Dampak Kebakaran Hutan. *Jurnal Lingkar Widya Swara*, 2014;. (4), 47–59.
5. Anjani, D. *Penerapan Model Ipat (Impact – Population-Affluence – Technology) Pada Emisi Karbon Dioksida (Co 2) Di Asean*. Universitas Airlangga. 2014,
6. Azuma, K., Kagi, N., Yanagi, U., & Osawa, H. Effects of low-level inhalation exposure to carbon dioxide in indoor environments: A short review on human health and psychomotor performance. *Environment International*, 2018;. 121 (August), 51-56. <https://doi.org/10.1016/j.envint.2018.08.059>
7. Samiaji, T. Gas Co 2 Di Wilayah Indonesia. *Berita Dirgantara*, 2011; 12(2), 68–75.
8. Adams, S., & Nsiah, C. Reducing carbon dioxide emissions; Renewable energy does matter? *Science of The Total Environment*, 2019; 693, 133 288. <https://doi.org/10.1016/j.scitotenv.2019.07.094>
9. Köster, E., Köster, K., Berninger, F., Prokushkin, A., Aaltonen, H., Zhou, X., & Pumpanen, J. Changes in fluxes of carbon dioxide and methane is the caused by fire in the Siberian boreal forest with continuous permafrost. *Journal of Environmental Management*, 2018; 228 (September), 405-415. <https://doi.org/10.1016/j.jenvman.2018.09.051>
10. Kim, Y., Knowles, S., Manley, J., & Radoias, V. Long-run health consequences of water pollution: Evidence from Indonesia's forest fires of 1997. *Economics and Human Biology*, 2017; 26, 186-198. <https://doi.org/10.1016/j.ehb.2017.03.006>
11. You, W., Lin, L., Wu, L., Ji, Z., Yu, J., Zhu, J., ... He, D. Geographical information system-based forest fire risk assessment integrating national forest inventory of data and analysis of its spatiotemporal variability. *Ecological Indicators*, 2017; 77, 176-184. <https://doi.org/10.1016/j.ecolind.2017.01.042>
12. Avetisyan, G. Assessment of carbon dioxide sequestration service: Case study of ijevan state sanctuary (Tavush 'region of Republic of Armenia). *Annals of Agrarian Science*, 2016; 14 (3), 187-190. <https://doi.org/10.1016/j.aasci.2016.07.003>

13. Hasmi, Mallongi, A. Health risk analysis of lead exposure from fish consumption among communities along Youtefa Gulf, Jayapura. *Pakistan Journal of Nutrition* 2016; Volume 15, Issue 10, 15 September Pages 929-935
14. Kayame, R., Mallongi, A. Relationships between smoking habits and the hypertension occurrence among the adults of communities in paniai regency, Papua Indonesia. *Indian Journal of Public Health Research and Development* 2018; Volume 9, Issue 1, Pages 332-336
15. Endah Yani, R.W., Mallongi, A., Andarini, S., Prijatmoko, D., Dewanti, I.R. The effect of zinc saliva on the toddlers' nutritional status. *Journal of International Dental and Medical Research*, 2016, Volume 9, Issue 1, Pages 29-32
16. Russeng, S.S., Saleh, L.M., Virani, D., Latief, A.W.L., Mallongi, A. The investigation of the lactic acid change among employee of national electrical power plan. *Indian Journal of Public Health Research and Development* 2018; Volume 9, Issue 1, Pages 361-36

Environmental Pollution and Health Problems Due to Forest Fires with Co2 Parameters

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

8 %
INTERNET SOURCES

2 %
PUBLICATIONS

4 %
STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

2%
★ www.ijop.net
Internet Source

Exclude quotes On
Exclude bibliography On

Exclude matches < 1%