



Syarfaini Suyuti <syarfaini.suyuti@uin-alauddin.ac.id>

Hemos iniciado la producción de su artículo [GACETA_1965] para Gaceta Sanitaria

1 message

C.Arora@elsevier.com <C.Arora@elsevier.com>
To: Syarfaini.suyuti@uin-alauddin.ac.id, pmc@agri.unhas.ac.id

Wed, Dec 30, 2020 at 5:02 PM

Nuestra referencia: GACETA 1965
Referencia del artículo:
Título del artículo: Risk factors preparation of stroke incidence in health institution employees who check up at the health service expo event Indonesia
Se publicará en: Gaceta Sanitaria

Estimado/a Dr. Syarfaini:

Gracias por elegir publicar en Gaceta Sanitaria. Por favor, lea este mensaje detenidamente ya que contiene información importante.

FINALIZAR LA PUBLICACIÓN DE SU ARTÍCULO:

Nos esforzamos al máximo para publicar los artículos de nuestros autores con la mayor rapidez y eficiencia posible y, por tanto, ya hemos iniciado el proceso de producción de su artículo aceptado. Para asegurar que su artículo se publica de acuerdo a sus deseos, por favor, complete los formularios que encontrará aquí:

<http://authors.elsevier.com/authorforms/GACETA1965/e1e3fd8a2e390db393ca0fd38d6238fa>

Si este enlace no funciona copie la dirección URL completa en su navegador. Tenga en cuenta que es posible que la dirección ocupe varias líneas en este mensaje. Tendrá que acceder con la contraseña de su Elsevier Profile, que debió crear en el momento en que envió su artículo.

REVISE SUS DATOS DE CONTACTO:

Por favor, revise los siguientes datos de contacto para que podamos ponernos en contacto con usted si es necesario:

Dr. Syarfaini
Alauddin State Islamic University Makassar
Teléfono: not available
Fax: not available
Correo electrónico: Syarfaini.suyuti@uin-alauddin.ac.id;pmc@agri.unhas.ac.id

SU NÚMERO DE REFERENCIA:

Por último, para que podamos ofrecerle el mejor servicio, tome nota del número de referencia de su artículo (GACETA 1965) e inclúyalo en todos sus mensajes cuando contacte con nosotros.

Muchas gracias por su colaboración. Por favor, no dude en ponerse en contacto con nosotros si tiene alguna consulta.

Atentamente,

Mr Chaitanya Arora
Data Administrator
Elsevier
Correo electrónico: C.Arora@elsevier.com

¿TIENE ALGUNA PREGUNTA O NECESITA AYUDA?

Para obtener más ayuda, visite nuestra página de Customer Support, donde puede buscar soluciones a una gran

variedad de temas y encontrar respuesta a las preguntas más frecuentes (página web en inglés con una selección de artículos en español). También puede contactar con nuestro equipo de Atención al Cliente por teléfono o por correo electrónico (en español: de lunes a viernes de 09:00 a 17:00 GMT; en inglés: de lunes a viernes, las 24 horas) y 24/7 a través del chat en vivo (solo en inglés).

Puede empezar aquí (página web en inglés): <http://service.elsevier.com/app/home/supporthub/publishing>

Copyright © 2018 Elsevier B.V. | Privacy Policy <http://www.elsevier.com/privacypolicy>

Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registration No. 1982084



Syarfaini Suyuti <syarfaini.suyuti@uin-alauddin.ac.id>

Publishing Agreement completed for your article [GACETA_1965]

1 message

Elsevier - Author Forms <Article_Status@elsevier.com>
To: syarfaini.suyuti@uin-alauddin.ac.id
Cc: Syarfaini.suyuti@uin-alauddin.ac.id, pmc@agri.unhas.ac.id

Thu, Dec 31, 2020 at 1:25 PM

Dear Dr. Syarfaini,

Thank you for completing the Publishing Agreement Form for your article *Risk factors preparation of stroke incidence in health institution employees who check up at the health service expo event Indonesia*. Please find attached a copy of the "Journal Publishing (License) Agreement" which you completed online on December 31, 2020.

If you have any questions, please do not hesitate to contact us. To help us assist you, please quote our article reference GACETA1965 in all correspondence.

Now that your article has been accepted, you will want to maximize the impact of your work. Elsevier facilitates and encourages authors to share their article responsibly. To learn about the many ways in which you can share your article whilst respecting copyright, visit: www.elsevier.com/sharing-articles.

We are committed to publishing your article as quickly as possible.

Kind regards,
Elsevier Researcher Support

Have questions or need assistance?

Please do not reply to this automated message.

For further assistance, please visit our Elsevier Support Center where you search for solutions on a range of topics and find answers to frequently asked questions.

You can also talk to our researcher support team by phone 24 hours a day from Monday-Friday and 24/7 by live chat and email.

© 2018 Elsevier Ltd | **Privacy Policy** <http://www.elsevier.com/privacypolicy>

Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registration No. 1982084. This e-mail has been sent to you from Elsevier Ltd. To ensure delivery to your inbox (not bulk or junk folders), please add article_status@elsevier.com to your address book or safe senders list.

 GACETA1965.html
20K



Syarfaini Suyuti <syarfaini.suyuti@uin-alauddin.ac.id>

Rights and Access form completed for your article [GACETA_1965]

1 message

Elsevier - Author Forms <Article_Status@elsevier.com>
To: syarfaini.suyuti@uin-alauddin.ac.id

Thu, Dec 31, 2020 at 1:25 PM

ELSEVIER

Dear Dr. Syarfaini,

Thank you for completing the Rights and Access Form for your article *Risk factors preparation of stroke incidence in health institution employees who check up at the health service expo event Indonesia* on December 31, 2020.


The Order Summary is attached to this email.

Your article will be free for everyone to read online as soon as it is published.

If you have any questions, please do not hesitate to contact us. To help us assist you, please quote our article reference GACETA1965 in all correspondence.

Now that your article has been accepted, you will want to maximize the impact of your work. Elsevier facilitates and encourages authors to share their article responsibly. To learn about the many ways in which you can share your article whilst respecting copyright, visit: www.elsevier.com/sharing-articles.

Kind regards,
Elsevier Researcher Support

**Seven strategies for you to create a brand and promote your research**

Learn how to give your research the visibility it deserves with these seven strategies.

> [Access module now](#)

Have questions or need assistance?

Please do not reply to this automated message.

For further assistance, please visit our Elsevier Support Center where you search for solutions on a range of topics and find answers to frequently asked questions.

You can also talk to our researcher support team by phone 24 hours a day from Monday-Friday and 24/7 by live chat and email.

© 2018 Elsevier Ltd | [Privacy Policy](http://www.elsevier.com/privacypolicy) <http://www.elsevier.com/privacypolicy>


Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registration No. 1982084. This e-mail has been sent to you from Elsevier Ltd. To ensure delivery to your inbox (not bulk or junk folders), please add article_status@elsevier.com to your address book or safe senders list.



Order Confirmation.html

11K

AUTHOR QUERY FORM

 ELSEVIER	Journal: GACETA	Please e-mail your responses and any corrections to:
	Article Number: 1965	E-mail:

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the ‘Q’ link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
Q1	Please confirm that given name and surname are correctly identified. The different colors indicate whether tagged as first or last name. Please note that proper identification is key for correct indexing of the article.
Q2	Correctly acknowledging the primary funders and grant IDs of your research is important to ensure compliance with funder policies. We could not find any acknowledgment of funding sources in your text. Is this correct?
Q3	Table 1 is not cited in the text. Please check that the citation suggested is in the appropriate place, and correct if necessary.
Q4	Please check and confirm “Conflict of interest”.
	<div style="border: 1px solid black; padding: 5px; margin-top: 20px;"> Please check this box or indicate your approval if you have no corrections to make to the PDF file <input type="checkbox"/> </div>

Thank you for your assistance.

Risk factors preparation of stroke incidence in health institution employees who check up at the health service expo event Indonesia

Syarfaini^{a,*}, Nildawati^b, Syahratul Aeni^b, Surahmawati^c, A. Syamsiah Adha^a, Munawir Amansyah^d

^a Science Of Nutrition, Public Health Study Program, Medical and Health Science Faculty of Alauddin State Islamic University Makassar, 92133, Indonesia

^b Epidemiology, Public Health Study Program, Medical and Health Science Faculty of Alauddin State Islamic University Makassar, 92133, Indonesia

^c Administration Of Health Policy, Public Health Study Program, Medical and Health Science Faculty of Alauddin State Islamic University Makassar, 92133, Indonesia

^d Environmental Health, Public Health Study Program, Medical and Health Science Faculty of Alauddin State Islamic University Makassar, 92133, Indonesia

ARTICLE INFO

Article history:

Received 24 August 2020

Accepted 4 December 2020

Available online xxx

Keywords:

Employee

Stroke incidence

Risk factor

ABSTRACT

Objective: To identify the risk factor of Stroke in health institution employees.

Methods: The research design was cross-sectional with 107 people who participated in this survey using purposive sampling method.

Result: The relationship between family history with the incidence of stroke obtained P value ($P < 0.05$), the relationship between smoking habits with the incidence of stroke P value = 0.012 ($P < 0.05$), the relationship between lack of physical activity with the incidence of stroke obtained $P = 0.476$ ($P > 0.05$), lack of consumption of fruits and vegetables obtained a value of $P = 0.091$ ($P > 0.05$) and alcohol consumption with the incidence of stroke obtained a value of $P = 0.616$ ($P > 0.05$).

Conclusion: Relationship between family history, smoking habits with the incidence of stroke.

© 2020 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Stroke is a non-infectious disease that causes the first disability and second death in the world.¹ Based on WHO data, stroke victims number 15 million in the world with a total of 5 million deaths and 5 million others experience permanent disabilities each year.^{2,3} Stroke is a cerebrovascular disease that is found not only in developed and developing countries. Stroke has get single of the greatest origins of disability in western nations. The incidence of stroke in the Netherlands ranks third as the cause of DALY's (Disability Adjusted Life Years).⁴

The number of stroke sufferers is greater in developing countries in Asia compared to developed countries. The incidence of stroke in China reaches 1.5–2 million people each year. Ischemic stroke is the near common case of stroke and the factors that influence it are genetic factors, environmental factors, and their relationship.⁵ The incidence of stroke in Indonesia reaches 500,000 people per year, including 125,000 people (2.5%) die and the rest are mildly or severely disabled. Ischemic stroke is around 80% of all stroke cases, while the incidence of hemorrhagic stroke is only about 20% of all stroke case.^{6,7} In Indonesia, The prevalence of stroke has expanded by 8.3/1000 population in 2007 to 12.1/1000 population in 2013.⁸

Increased body mass and the number of risk will increase the potential for the occurrence of stroke. Modifiable risk factors for stroke include a history of stroke, hypertension, heart disease, dia-

betes mellitus, asymptomatic carotid disease, transient ischemic attack, hypercholesterolemia, oral contraceptive use, narcotic use, antiphospholipid antibodies, obesity, smoking, alcoholics, hyperuricemia, increased hematocrit. While the factors that cannot be modified are age, sex, heredity, and race/ethics.⁹

Methods

This was a cross-sectional approach in South Sulawesi, Indonesia. This research is a quantitative analytic study designed with a cross sectional approach. The results of medical examinations in the Health Service EXPO Event South Sulawesi 2018 were used as secondary data in this study. The population in this study were employees of the municipal/district health office totaling 126 people. A total of 107 people on the health check form and fulfilling the inclusion criteria were sampled in this study. Secondary data was obtained from the results of a health examination conducted at communicable disease POSBINDU during the Health Service EXPO Event.

Data collected in this study include data on age, sex, family history of illness, history of self-illness, smoking history, physical activity, consumption of vegetables and fruit, and alcohol consumption. Data analysis conducted in this research is univariate analysis and bivariate analysis. This study used univariate analysis to specify the frequency of data processing, namely age, sex, history of stroke, family history of illness, physical activity, utilization of vegetables and fruits, alcohol consumption, and smoking habits. While the bivariate analysis was carried out to determine the relationship of two variables namely the dependent variable and the independent variable by using parametric test analysis namely Pearson correlation analysis with the SPSS program.

Peer-review under responsibility of the scientific committee of the 1st International Conference on Safety and Public Health (ICOS-PH 2020). Full-text and the content of it is under responsibility of authors of the article.

* Corresponding author.

E-mail addresses: Syarfaini.suyuti@uin-alauddin.ac.id, pmc@agri.unhas.ac.id (Syarfaini).

<https://doi.org/10.1016/j.gaceta.2020.12.014>

0213-9111/© 2020 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Syarfaini, et al. Risk factors preparation of stroke incidence in health institution employees who check up at the health service expo event Indonesia. Gac Sanit. 2020. <https://doi.org/10.1016/j.gaceta.2020.12.014>

Table 1
Proportion of risk factor for stroke incidence.

Variable	Frequency (n)	Percentage (%)		
Age	20-26 Years	16	15	
	27-33 Years	18	17	
	34-40 Years	24	22	
	41-47 Years	18	17	
	48-54 Years	16	15	
	55-61 Years	13	12	
	>62 Years	2	2	
Gender	Men	34	32	
	Women	73	68	
Family history	Diabetes	Yes	37	34.6
		No	70	65.4
	Hypertension	Yes	30	28
		No	77	72
	heart disease	Yes	9	8
		No	98	92
	Stroke	Yes	5	4.6
		No	102	95.4
	High cholesterol	Yes	11	10.2
		No	96	89.8
	Asthma	Yes	6	5.6
		No	101	94.4
	Cancer	Yes	1	1
		No	106	99

Source: Secondary Data, 2018.

Result

Results are research findings in the form of data (tables and figures) from observations. The results should be clear and concise (Table 1).

The subjects were employees in health institution. Most of the participant (22%) were in the 34-40 years age category. The major (68%) of participant are female. Most of them without family history of disease.

Based on Table 2, there is a relationship between heart disease in the family and the incidence of stroke ($P=0.000$). There is a related between stroke in the family and the incidence of stroke ($P=0.000$). For cancer, ($P=0.000$) was significantly associated between cancer in the family and the occurrence of stroke.

According to Table 3, there is a related between smoking habits and stroke incidence ($P=0.012$). There is no related between lack of physical activity and the incidence of stroke ($P=0.476$) in health agency employees who conduct examinations at non communicable disease POSBINDU Health Service EXPO Event.

According to Table 4, there is no statistically significant difference in the lack of consumption of vegetables and fruit with the incidence of stroke ($P=0.091$). There is no relationship between alcohol consumption and the incidence of stroke among employees of health agencies who conduct examinations at noncommunicable disease POSBINDU Health Service EXPO Event ($P=0.616$).

Discussion

Family disease history

The risk of stroke for heredity, experts say that there is a recessive gene that affects it. The gene is associated with diseases that are risk factors for stroke. Diseases associated with these genes include diabetes, hypertension, hyperuricemia, hyperlipidemia, coronary heart disease, and abnormalities in blood vessels that are hereditary. the testing using a bivariate analysis of family history based on study (history of heart disease, stroke, and family history with cancer) result a significant related with the incidence of stroke among health agency employees who carry out checks at noncommunicable diseases POSBINDU Health Service EXPO Event. This finding

was consistent with previous studies that a family history of stroke is one of the risk factors for stroke complications. Our finding was consistent with previous studies that a family history of stroke is one of the risk factors for stroke complications. The risk of suffering from an ischemic stroke increased two times more in 500 women aged 18-44 years with a family history of stroke.¹⁰

A family disease history of diabetes, asthma, hypertension, and cholesterol are no significant relationship in stroke incidence. Family history is one of the potential risk factors associated with stroke incidence. However, family history is not an independent risk factor that causes a stroke. This dominant study did not have a significant related between family disease history of illness and the stroke incidence. Increased risk of stroke in family history (heredity) is obtained through several mechanisms, namely genetic factors, genetic sensitivity, cultural/environmental factors, and lifestyle as well as interactions between genetic and environmental factors.¹¹

Smoking habits

Smoking is one of the main causes of heart problem and stroke. The incidence of stroke will increase when combined with other risk factors, especially hypertension. This study is following with research published by Aliah in Yogyakarta, it shows smoking is significantly related to the incidence of stroke and smokers are at 4.51 times the risk of stroke compared to nonsmokers ($P=0.000$; $OR=4.51$; $95\% CI=2.107-9669$).¹² Other studies state that the risk of stroke will increase by about 2-3 times in smokers compared to nonsmokers and will only disappear after stopping smoking for 5-10 years.¹³

Lack of physical activity

This study shows that there is no significant related between less physical activity and the incidence of stroke. Different findings were obtained by Perawaty (2014) which states that people who have less physical activity have a risk of 3.46 times greater stroke than those who have lots of physical activity.¹⁴ WHO report which explains that a sedentary lifestyle or physical inactivity is the main cause of death and disease. Physical activities such as sports carried out with the principle of FIT (frequency, intensity, and time)

Table 2
Analysis of relationships between family history with Stroke incidence.

Family History		Stroke Total				P
		Yes	No	N	%	
Heart	Yes	1	0	1	1	0.000
	No	4	102	106	99	
Stroke	Yes	1	0	1	1	0.000
	No	4	102	106	99	
Cancer	Yes	1	0	1	1	0.000
	No	4	102	106	99	

Source: Secondary Data, 2018.

Table 3
Analysis of the relationship between smoking habits and lack of physical with stroke incidence.

	Stroke		Total N	Total %	P
	Yes	No			
<i>Smoking habits</i>					
Yes	4	28	32	29.9	0.012
No	1	74	75	70.1	
<i>Lack of physical activity</i>					
Yes	2	26	28	26.2	0.476
No	3	76	79	73.8	

Source: Secondary Data, 2018.

Table 4
Analysis of the relationship between fruits, vegetables and alcohol consumption with the Stroke incidence.

		Stroke		Total N	Total %	P
		Yes	No			
<i>Less consumption of fruits and vegetables</i>						
Yes	4	5	64	69	64.5	0.091
No		0	38	38	35.5	
<i>Alcohol consumption</i>						
Yes		0	5	5	4.7	0.616
No		5	97	102	95.3	

Source: Secondary Data, 2018.

146 can minimize the risk of death from heart problems and stroke,
147 help prevent and reduce hypertension, and can control the risk of
148 obesity by up to 50%.² Research on stroke risk factors in American
149 blacks reports that low physical activity has the potential to be a
150 contributing factor to stroke incidence and is obtained RR: 2.07;
151 95%; CI: 1.25-3.43.¹⁵

152 *Less consumption of fruits and vegetables*

153 The results of this study pointed that that there was no relation-
154 ship significantly between consuming fewer fruits and vegetables
155 with stroke incidence among health agency workers who exam-
156 ined at noncommunicable disease POSBINDU Health Service EXPO
157 Event. In contrast to previous studies in Japan, fruits and also green
158 and yellow vegetable associates with the risk of death from stroke in
159 men and women. Eating vegetables and fruits reduces the potential
160 of stroke by 20-40% in both women and men.¹⁶

161 Vegetables and fruits contain various nutrients including
162 vitamins, minor minerals, fiber, and various biologically active
163 components. This component is known as phytochemicals which
164 have complementary and (complementary) mechanisms in antiox-
165 idant activity, reduce thrombotic aggregation, alter cholesterol
166 metabolism, and reduce blood pressure.¹⁷

167 *Alcohol consumption*

168 The results of this research are in accordance with previous
169 study that examining the risk factors for stroke in the black race of

170 America and the result is that the consumption of alcohol in small
171 amounts in the acute or chronic categories after being controlled
172 by smoking and hypertension factors showed no significant rela-
173 tionship, OR: 1.68; 95% CI: 0.84-3.35.¹⁵ Another study conducted
174 in Japan which states that there is a difference in the risk of stroke
175 between drinkers every week and those who only consume alcohol
176 occasionally.¹⁸ This study is also in accordance with a cohort study
177 conducted on 38,156 men in health professions in the United States.
178 It is found that individuals who consume large amounts of alcohol
179 and more than 2 times/day is a main factor in the occurrence of
180 ischemic stroke.¹⁹

181 In one of the hadiths, the Messenger of Allah explained that there
182 is no better favor after the true belief except healthy favors (al-
183 fityah). Therefore, he always advised his friends to multiply one of
184 the best prayers, namely to always be given forgiveness (al-'afw)
185 and safety (al-'afiyah) from all kinds of dangers, including illness,
186 both physical and mental illness.²⁰

187 **Conclusion**

188 Based on 11 independent variables studied, 4 variables were
189 found that had a relationship significantly with stroke incidence.
190 There is a relationship between family history (Heart, stroke, can-
191 cer) with stroke incidence. Smoking habits with the incidence
192 of stroke have a significant relationship. There is no relationship
193 between, consumption of vegetables and fruit, physical activity,
194 and consumption of alcohol, with the incidence of stroke.

195 Suggestions for similar research, which need to be improved
196 in the measurement instrument (questionnaire). Because of the
197 weaknesses in this study, there are several risk factors that are not
198 in accordance with the calculation standards, such as consumption
199 of fruits and vegetables that are not measured based on 24-h food
200 recall and physical activity measurements that are not according to
201 WHO standards.

202 Conflict of interests

203 204 The authors declare that they have no conflict of interest.

204 References

205 1. Feigin VL. Stroke in developing countries: can the epidemic be
206 stopped and outcomes improved? *Lancet Neurol.* 2007;6:94-7,
207 [http://dx.doi.org/10.1016/S1474-4422\(07\)70007-8](http://dx.doi.org/10.1016/S1474-4422(07)70007-8).
208 2. World Health Organization. Global strategy on diet, phys-
209 ical activity and health; 2004. [https://www.who.int/
210 dietphysicalactivity/strategy/eb11344/strategy_english_web.pdf?ua=1](https://www.who.int/dietphysicalactivity/strategy/eb11344/strategy_english_web.pdf?ua=1)
211 3. World Health Organization. Global burden of stroke; 2010.
212 4. Janssen AWM, de Leeuw FE, Janssen MCH. Risk factors for ischemic stroke and
213 transient ischemic attack in patients under age 50. *J Thromb Thrombolysis.*
214 2011;31:85-91, <http://dx.doi.org/10.1007/s11239-010-0491-3>.
215 5. Xu C, Wang F, Wang B, et al. Minor allele C of chromosome 1p32 sin-
216 gle nucleotide polymorphism rs11206510 confers risk of ischemic
217 stroke in the Chinese Han population. *Stroke.* 2010;41:1587-92,
218 <http://dx.doi.org/10.1161/STROKEAHA.110.583096>.
219 6. Suwanwela NC. Stroke epidemiology in Thailand. *J Stroke.* 2014;16:1,
220 <http://dx.doi.org/10.5853/jos.2014.16.1.1>.
221 7. Farhoudi M, Mehrvar K, Sadeghi-Bazargani H, et al. Stroke subtypes, risk fac-
222 tors and mortality rate in northwest of Iran. *Iran J Neurol.* 2017;16:112-7.
<https://pubmed.ncbi.nlm.nih.gov/29114365>

223 8. Badan Penelitian dan Pengembangan Kesehatan. Laporan Riset Kesehatan Dasar;
224 2007. <http://kesga.kemkes.go.id/images/pedoman/Riskesdas2007Nasional.pdf>
225 9. Perhimpunan Dokter Spesialis Saraf Indonesia. Guideline Stroke Tahun 2011.
226 (Misbach Y, Lamsudin R, Allah A, Basyiruddin, eds.). Pokdi Stroke Perdossi; 2011.
227 <http://skybooks.club/download/books.php?Guideline+stroke-2011>
228 10. Muliawati R. Beberapa Faktor Risiko Yang Berpengaruh Terhadap Kejadian
229 Stroke Iskemik Pada Penderita Diabetes Mellitus Tipe 2 (Studi Kasus di RSUD Dr
230 Kariadi Semarang); 2015. Published online. <http://eprints.undip.ac.id/46429/>
231 11. Goldstein LB, Robert A, Alberts MJ, et al. Primary pre-
232 vention of ischemic stroke. *Stroke.* 2006;37:1583-633,
233 <http://dx.doi.org/10.1161/01.STR.0000223048.70103.F1>.
234 12. Aliah A, Widjaja D. Faktor Risiko Stroke pada Beberapa Rumah Sakit di Makasar
235 (Januari-September 2000). *J Med Nus.* 2004;25:1-6.
236 13. Shah RS, Cole JW. Smoking and stroke: the more you smoke the
237 more you stroke. *Expert Rev Cardiovasc Ther.* 2010;8:917-32,
238 <http://dx.doi.org/10.1586/erc.10.56>.
239 14. Perawaty P, Dahlan P, Astuti H. Pola makan dan hubungannya dengan keja-
240 dian stroke di RSUD dr. Doris Sylvanus Palangka Raya. *J Gizi dan Diet Indones.*
241 2014;2:51-61, [http://dx.doi.org/10.21927/ijnd.2014.2\(2\).51-61](http://dx.doi.org/10.21927/ijnd.2014.2(2).51-61).
242 15. Gillum RF. Risk factors for stroke in blacks: a critical review. *Am J Epidemiol.*
243 1999;150:1266-74, <http://dx.doi.org/10.1093/oxfordjournals.aje.a009957>.
244 16. Sauvaget C, Nagano J, Allen N, Kodama K. Vegetable and fruit intake and stroke
245 mortality in the Hiroshima/Nagasaki Life Span Study. *Stroke.* 2003;34:2355-60,
246 <http://dx.doi.org/10.1161/01.STR.0000089293.29739.97>.
247 17. Lipoeto NI. Zat Gizi Dan Makanan Pada Penyakit Kardiovaskuler. Padang:
248 Andalas University Press; 2006.
249 18. Hiroyasu I, Shunroku B, Toshifumi M, et al. Alcohol consumption and
250 risk of stroke among middle-aged men: the JPHC study cohort I. *Stroke.*
251 2004;35:1124-9, <http://dx.doi.org/10.1161/01.STR.0000124459.33597.00>.
252 19. Mukamal KJ, Ascherio A, Mittleman MA, et al. Alcohol and
253 risk for ischemic stroke in men: the role of drinking pat-
254 terns and usual beverage. *Ann Intern Med.* 2005;142:11-9,
255 <http://dx.doi.org/10.7326/0003-4819-142-1-200501040-00007>.
256 20. Hanbal A ibn MI. English Translation of Musnad Imam
257 Ahmad Bin Hanbal. (Khattab H, ed.). Darussalam; 2012.
258 <https://books.google.co.id/books?id=CHuBMwEACAAJ>

UNCORRECTED