




## ORIGINAL ARTICLE

# Dietary *Dracocephalum kotschy* essential oil improved growth, haematology, immunity and resistance to *Aeromonas hydrophila* in rainbow trout (*Oncorhynchus mykiss*)

Hafsan Hafsan<sup>1</sup>  | Dmitry Bokov<sup>2,3</sup> | Walid Kamal Abdelbasset<sup>4,5</sup> |  
 Mustafa M. Kadhim<sup>6,7,8</sup> | Wanich Suksatan<sup>9</sup>  | Hasan Sh. Majdi<sup>10</sup> |  
 Gunawan Widjaja<sup>11,12</sup> | Abduladheem Turki Jalil<sup>13,14</sup>  | Maytham T. Qasim<sup>15,16</sup> |  
 Mahboubeh Balvardi<sup>17</sup>

<sup>1</sup>Biology Department, Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Gowa, Indonesia

<sup>2</sup>Institute of Pharmacy, Sechenov First Moscow State Medical University, Moscow, Russian Federation

<sup>3</sup>Laboratory of Food Chemistry, Federal Research Center of Nutrition, Biotechnology and Food Safety, Moscow, Russian Federation

<sup>4</sup>Department of Health and Rehabilitation Sciences, College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Al Kharj, Saudi Arabia

<sup>5</sup>Department of Physical Therapy, Kasr Al-Aini Hospital, Cairo University, Giza, Egypt

<sup>6</sup>Department of Dentistry, Kut University College, Kut, Iraq

<sup>7</sup>College of Technical Engineering, The Islamic University, Najaf, Iraq

<sup>8</sup>Department of Pharmacy, Osol Aldeen University College, Baghdad, Iraq

<sup>9</sup>Faculty of Nursing, HRH Princess Chulabhorn College of Medical Science, Chulabhorn Royal Academy, Bangkok, Thailand

<sup>10</sup>Department of Chemical Engineering and Petroleum Industries, Al-Mustaqbal University College, Babylon, Iraq

<sup>11</sup>Faculty of Public Health, Universitas Indonesia, Depok, Indonesia

<sup>12</sup>Faculty of Law, Universitas Krisnadwipayana Indonesia, Jatiwaringin, Indonesia

<sup>13</sup>Faculty of Biology and Ecology, Yanka Kupala State University of Grodno, Grodno, Belarus

<sup>14</sup>College of Technical Engineering, The Islamic University, Najaf, Iraq

<sup>15</sup>Ministry of Education, Directorate Thi-Qar Education, Thi-Qar, Iraq

<sup>16</sup>Department of Anesthesia, College of Health and Medical Technology, Al-Ayen University, Thi-Qar, Iraq

<sup>17</sup>Department of Biology, Payame Noor University, Tehran, Iran

## Correspondence

Hafsan Hafsan, Biology Department,  
 Faculty of Science and Technology,  
 Universitas Islam Negeri Alauddin  
 Makassar, Sultan Alauddin Street, Gowa  
 92118, Indonesia.  
 Email: [hafsan.bio@uin-alauddin.ac.id](mailto:hafsan.bio@uin-alauddin.ac.id)

## Abstract

In the present study, the effect of *Dracocephalum kotschy* essential oil (DKE) was investigated on growth, haematology, immune and antioxidant defence system and resistance of rainbow trout juveniles to bacterial infection (*Aeromonas hydrophila*). For this purpose, the fish were fed a diet containing different concentrations of DKE including 0 (control), 0.2, 0.25 and 0.3 mg/kg diet in three replicates for 60 days. After feeding trial, the fish were challenged with a pathogenic dose of *A. hydrophila*. Based on the results, immune components in plasma (alternative complement activity [ACH<sub>50</sub>], IgM content, lysozyme activity, total protein and total albumin) and mucus (protease activity, IgM content and lysozyme activity) significantly elevated in fish fed diet containing 0.2 and 0.25 mg DKE/kg diet compared with other groups ( $p < 0.01$ ). The plasma bactericidal activity increased in all DKE-supplemented fish. Supplementation of fish with 0.3 mg DKE/kg diet depressed the mucosal lysozyme activity and IgM content.

The liver antioxidant enzymes, glutathione peroxidase (GPx), catalase (CAT) and superoxide dismutase (SOD) showed increased activity in response to 0.25 and 0.3 mg DKE/kg diet compared with other experimental diets ( $p < 0.01$ ). Dietary DKE changed the haematology of the fish. The haematocrit, red blood cell count (RBC), white blood cell (WBC), haemoglobin content and mean erythrocyte cell haemoglobin content (MCHC) increased in response to DKE. In contrast, the mean corpuscular haemoglobin (MCH) decreased in fish fed 0.2 and 0.25 mg DKE/kg diet compared with other diets ( $p < 0.01$ ). The supplementation of fish with 0.2 and 0.25 mg DKE/kg diet decreased the expression of cytokine-related genes, TNF- $\alpha$  and IL-8. In contrast, the DKE up-regulated the expression of the immune-related genes, C3 and lysozyme. DKE at concentration of 0.3 mg/kg diet depressed the C3 and lysozyme gene expressions. The DKE supplementation decreased the mortality rate of the fish after bacterial challenge compared with non-supplemented ones ( $p < 0.01$ ). In conclusion, the findings of this study revealed the immune-stimulating effects of DKE at optimized dietary concentrations. In addition, DKE decreased the mortality induced by *A. hydrophila* infection.

#### KEYWORDS

bacterial, *Dracocephalum kotschy*, essential oil, fish, immunity

## 1 | INTRODUCTION

Aquaculture is known as one of the most important ways to provide the protein needs of human (Pradeepkiran, 2019). However, the development of intensive aquaculture and super-intensive aquaculture has increased the risk of fish diseases as well as environmental pollutions (Cao et al., 2007; Cole et al., 2009; Jackson, 1978; Páez-Osuna, 2001; Romero et al., 2012). Today, antibiotics are widely used to treat fish diseases, but there are still some problems with these chemicals that may limit their use (Lulijwa et al., 2020; Romero et al., 2012). Continued use of antibiotics may create resistant bacterial strains, and in addition, the accumulation of these chemicals and their derivatives in fish meat can threaten human health as a consumer (Brown et al., 2021; Lulijwa et al., 2020). Furthermore, the release of antibiotics into natural aquatic environments can have a negative impact on the flora and fauna and ultimately the entire food web of that environment (Liu et al., 2017; Lulijwa et al., 2020). Owing to these problems, it seems necessary to replace antibiotics with safe alternatives. Dietary supplements, including plant-based supplements, are currently used to keep health and to treat many diseases in humans and animals (Fattepur et al., 2018; Gondo, 2021; Othman et al., 2019; Roslan & Shariff, 2021; Shalaby, 2015, 2018; Shalaby, Sakoury, Harthi, et al., 2020; Shalaby, Sakoury, Kholif, et al., 2020; You et al., 2021; Zharif et al., 2018). During the last decade, many studies have reported the medicinal, antioxidant, immunogenic and antibacterial properties of medicinal plants and their derivatives in fish (Ahmadifar et al., 2019; Farsani et al., 2019; Rashidian, Bahrami Gorji, et al., 2020; Van Doan et al., 2020; Bibi et al., 2021; DeiviArunachalam et al., 2021; Ghafarifarsani Kachuei et al., 2021;

Ghafarifarsani, Hoseinifar, Adorian, et al., 2021; Ghafarifarsani, Hoseinifar, Talebi, et al., 2021; Ghafarifarsani, Rashidian, et al., 2021; Qahir et al., 2021; Rashidian et al., 2021; Yousefi, Farsani, et al., 2021; Yousefi, Ghafarifarsani et al., 2021; Raissy et al., 2022; Yousefi et al., 2022). However, the identification and introduction of the new species with immunogenic and antibacterial properties can be useful for aquaculture enhancement. *Dracocephalum* with more than 60 species is native to temperate regions of Asia and Europe, which is consumed in many parts of the world due to its high medicinal properties (Heydari et al., 2019). *Dracocephalum kotschy* Boiss plant is one of the well-known species of *Dracocephalum*, which is used in treatment of infectious diseases due to its medicinal properties and active compounds. The chemical composition of *Dracocephalum kotschy* includes bioactive compounds with anti-cancerous, antioxidant, antibacterial, anti-nociceptive, anti-hyperlipidaemic, anti-spasmodic, cytotoxic and immunostimulatory effects (Heydari et al., 2019; Moradi et al., 2020; Sadraei et al., 2017; Talari et al., 2014). In a study by Ghavam et al. (2021), the antibacterial properties of *D. kotschy* essential oil (DKE) were attributed to some bioactive compounds in the biochemical composition of DKE including  $\alpha$ -pinene, terpinen-4-ol, limonene, neral and geraniol. In addition, the results of Asghari et al. (2015) showed that the methanol extracts of *D. kotschy* inhibit the growth of *Mycobacterium tuberculosis*. The extracts of *D. kotschy* showed antibacterial effects against Gram-positive bacteria (*Staphylococcus aureus*, *Bacillus cereus* and *Listeria Monocytogenes*) and Gram-negative bacteria (*Salmonella enterica*, *Escherichia coli* and *Enterica Aerogenes*) (Kamali et al., 2015). According to a literature review, we found no data regarding the immunogenic, antioxidant and antibacterial

effects of *D. kotschy* in fish. Therefore, the present study was conducted to evaluate the effects of DKE on immune and antioxidant defence system and resistance of rainbow trout juveniles to infection induced by *A. hydrophila*. As an opportunistic bacterium, *A. hydrophila* has cosmopolitan distribution in water bodies of the world. *A. hydrophila* causes the bacterial haemorrhagic septicaemia disease in many species of warm and cold water fish including rainbow trout (Constantino Casas et al., 1997; Harikrishnan & Balasundaram, 2005; Nya & Austin, 2010). The results of the present study may suggest a natural way to enhance the immunity and to control infection induced by *A. hydrophila* in rainbow trout.

## 2 | MATERIALS AND METHODS

### 2.1 | Fish and feeding trial

Rainbow trout fish with an average weight of  $55 \pm 5.6$  g were purchased from a local cold water fish farm, and then transported using a tank with continuous oxygenation to lab and finally distributed into the 12 polyethylene tanks (containing 400 L disinfected and chlorinated water) for 7 days acclimation. During acclimation, the fish were fed three times a day at a rate of 4% of body weight with a commercial rainbow trout diet (Faradaneh Co., GFT1; diet form: pellet, pellet size:  $5 \pm 0.4$  mm; diet composition: total crude protein: 40%, total crude lipid: 14%, total crude ash: 9% and total crude fibre: 3%).

After the adaptation to lab conditions, fish were distributed into the 12 tanks at density of 30 fish per tank and fed DKE at dietary levels of 0 (control), 0.2, 0.25 and 0.3 mg/kg diet in three replicates. Feeding was carried out daily at a rate of 4% of fish body weight. The dietary concentrations of the essential oil in this study were selected based on the in vitro inhibitory effect (or disk diffusion assay) of the essential oil on the growth of *A. hydrophila* according to the guidelines of Clinical & Laboratory Standards Institute (CLSI), formerly known as NCCLS (Wayne, 2007), with some modifications. Briefly, the sterile Petri dishes (9 cm in diameter) were prepared with 20 ml of Mueller–Hinton medium. Then, a 100 µl of the bacterial suspension ( $10^8$  CFU/ml) was spread on

the plates. After 6 min, a sterile filter paper disc (6 mm) containing different concentrations of DKE (0, 0.05, 0.1, 0.15, 0.2, 0.25 and 0.3 mg) was placed on the surface of the plates at 4°C for 1 h followed by incubation at 37°C for 30 h. The diameters of the inhibition zones (mm) were measured, including the diameter of the disks. Gentamycin (30 µg/disk) was used as positive control. All tests were done in three replicates. A disk diffusion assay was conducted to determine the highest antibacterial activity for DKE (Table 1). In this regard, the highest antibacterial activity was observed in treatments, 20, 25 and 30 which these concentrations were selected for feeding trial. During feeding experiment, water was daily renewed by 90% and also the water quality indices (temperature:  $15.3 \pm 0.3^\circ\text{C}$  [using thermometer, ZEAL, UK], oxygen:  $6.8 \pm 0.11$  mg/L [using portable oxygen meter: Oxyguard Polaris Dissolved Oxygen Meter, Dynamic Aqua Supply Ltd, Canada], total ammonia:  $0.008 \pm 0.002$  mg/L and pH:  $7.5 \pm 0.13$  [using portable pH meter, Model AE-PH501]) were monitored daily.

### 2.2 | GC-Mass spectrometry of *Dracocephalum kotschy* essential oil

The gas chromatography–mass spectrometry (Shimadzu QP2010 Ultra series instrument, flow rate 0.7 ml/min, temperature rate:  $3^\circ\text{C}/\text{min}$  and final temperature:  $330^\circ\text{C}$ ) was used to analyse the chemical composition of DKE.

### 2.3 | Growth and survival parameters

The growth and survival parameters were measured after feeding trial by the following equations (Ghafarifarsani, Hoseinifar, Javahery, et al., 2022):

$$\text{Weight gain (WG) (\%)} = \frac{\text{final weight} - \text{initial weight}}{\text{initial weight}} \times 100$$

$$\text{Specific growth rate (SGR) (\%/d)} = \left( \frac{\ln \text{final wt(g)} - \ln \text{initial wt(g)}}{\text{days}} \right) \times 100$$

Genes	Primers	Primer sequences	ID number
Lysozyme	Lysozyme Forward	TTCCCCAGGTATCCCATGAT	AB027305
	Lysozyme Reverse	GTGTCTGATGTGGCTGTGCT	
Complement	C3 Forward	AGCTTGCTGACTGGCTTTGT	100136951
	C3 Reverse	TCATAAACGGTGACCCCAAC	
TNF-α	TNF-α Forward	CATCGGCCCTGCACCTATA	X99303
	TNF-α Reverse	TGTGGAACACAGCAGGGTTGG	
IL-8	IL-8 Forward	AGAATGTACGCCAGCCTTGT	AJ279069
	IL-8 Reverse	TCTCAGACTCATCCCCTCAGT	
β-Actin	β-Actin Forward	ACTGCACAGCCAAGAGAGTTCA	AB039726
	β-Actin Reverse	GTTATTAAAGCGGCCGATATGC	

TABLE 1 Sequences of oligonucleotide primers for real-time PCR

The gene efficacy for target and reference genes was 1.93–1.98.

$$\text{Feed conversion rate (FCR)} = \frac{\text{total feed consumed(g)}}{\text{weight gain(g)}}$$

$$\text{Survival rate (SR) (\%)} = \left( \frac{\text{final fish numbers}}{\text{initial fish numbers}} \right) \times 100$$

## 2.4 | Sampling

The blood, mucus and tissue samples were taken from 10 fish per tank after the feeding trial. For this purpose, fish were starved for 24 h, anesthetized with 20 mg/L clove oil and then few ( $n = 3$ ) were placed in a nylon bag for collection of mucus based on the method of Subramanian et al. (2007). The blood samples were collected from remainders ( $n = 7$ ) from caudal vein using 2-ml syringe. The plasma part of blood was separated by centrifuging at 4°C for 5 min. In addition, a part of blood was allocated for measurement of the haematological indices.

For tissue sampling, the fish were first sacrificed with high dose of ethyl 3-aminobenzoate methanesulphonate (100 g/L) to take liver sample. The plasma and liver samples were finally stored in liquid nitrogen until biochemical analysis.

## 2.5 | Biochemical assays

### 2.5.1 | Immune parameters

The activity of lysozyme in plasma and mucus was assayed using lyophilized *Micrococcus luteus* according to Ellis (1990). The immunoglobulin (IgM) content of plasma and mucus was determined using polyethylene glycol according to Siwicki and Anderson (1993). The plasma alternative complement activity ( $\text{ACH}_{50}$ ) was assayed using the method of Ortuno et al. (2001) using *haemolysis* of rabbit red blood cells. The plasma bactericidal activity was evaluated by mixing plasma samples with bacterial suspension (*A. hydrophila*, OD: 0.5 at 546 nm) and finally calculating bacterial colony forming units (CFUs) created on nutrient agar plates after 24 h incubation at 36°C (Rao et al., 2006).

The total protein and albumin levels in plasma were spectrophotometrically determined using commercial assay kits (Sigma-Aldrich Co., USA) according to manufacturers' instructions. Protease activity in mucus was determined using the azocasein hydrolysis method, as described by Ross et al. (2000). Mucosal alkaline phosphatase (ALP) activity was measured using a commercial assay kit (Sigma-Aldrich Co., USA) according to manufacturers' instructions.

### 2.5.2 | Liver antioxidant enzymes

All antioxidant enzymes were determined using commercial assay kits, as instructed by manufacturer (Sigma-Aldrich CO, USA). Catalase (CAT) activity was assayed at 240 nm using catalysis of hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) according to Claiborne (1985). Superoxide dismutase

(SOD) activity was colourimetrically determined at 440 nm through oxidation of xanthine to superoxide radicals (SR) and generation of a red formazan dye upon reaction of WST-1 reagent with the SR (Marklund & Marklund, 1974). Glutathione peroxidase (GPX) activity was measured through oxidation of glutathione by GPx and assaying oxidized glutathione at 340 nm. The lipid peroxidation was spectrophotometrically (at 532 nm) assayed upon reaction of malondialdehyde (MDA) with thiobarbituric acid (Utley et al., 1967).

### 2.5.3 | Haematological parameters

The total count of red and white blood cells was measured using a Neubauer haemocytometer slide at a magnification of 400× (Harikrishnan et al., 2009). Haematocrit (Hct) was measured using the microhaematocrit method (Blaxhall & Daisley, 1973). The haemoglobin (Hb) content was determined using the cyanohaemoglobin method (Oriakpono et al., 2012). The haematological indices including mean corpuscular haemoglobin (MCH), mean corpuscular volume (MCV) and corpuscular haemoglobin concentration (MCHC) were estimated according to the following equations (Blaxhall & Daisley, 1973):

$$\text{MCHC} = \text{Hb} \times \frac{10}{\text{Hct}}; \text{MCV} = \text{Hct} \times \frac{10}{\text{RBC}(\text{million})};$$

$$\text{MCH} = \text{Hb} \times \frac{10}{\text{RBC}(\text{million})}$$

## 2.6 | Gene expression assay

### 2.6.1 | RNA extraction

The total RNA content was extracted from liver tissues using acid guanidinium thiocyanate-phenol-chloroform based on the method described by Awad et al. (2013) with some modifications. The quantity and quality of the extracted RNA (ER) was estimated by determining the absorbance at 260 nm, whereas the quality of ER was evaluated using electrophoresis according to Kwasek et al. (2014).

### 2.6.2 | RT-PCR

One microgram of ER was allocated to produce first-strand cDNAs using a Fermentas cDNA synthesis kit for reverse transcription polymerase chain reaction (RT-PCR) according to the manufacturer's protocol and a set of oligo-dT as primer. The real-time PCR primers were synthesized based on the sequences of DNA from GenBank (accessions numbers listed in Table 1) and using Gene Runner (version 6) software. The  $\beta$ -actin gene was considered as reference gene to evaluate the gene expressions. The real-time PCR was done using an iCycler (BioRad, USA) and commercial kit (Bio flux-Bioer Technology Co., China) following the manufacturer's protocols. The fold change in TNF- $\alpha$ , IL-8, complement and lysozyme relative to mRNA expression was estimated using the  $2^{-\Delta\Delta\text{Ct}}$  method (Livak &

Schmittgen, 2001) and the data were analysed using iQ5 optical system software version 2.1 (Bio-Rad).

## 2.7 | Bacterial challenge

After 60 days of feeding experiment, the remaining fish in the tanks were exposed to *A. hydrophila* bacterium by injection. The stock of the bacterium (Strain number: IBRC-M 10814) was cultured on agar medium for 24 h at 37°C. A sub-lethal dose of bacteria ( $1.3 \times 10^7$  cells/ml phosphate buffered saline [PBS]) was used for bacterial challenge, which was calculated through determining the 7-day LD<sub>50</sub> ( $3 \times 10^7$  cells/ml) in a previous exposure experiment by LaPatra et al. (2010). The cumulative fish mortality was recorded daily for 10 days. In addition, *A. hydrophila*-induced mortality was confirmed through re-isolating the organism from the liver on at least 30% of the dead fish using conventional methods.

## 2.8 | Statistical analysis

All statistical analysis was done using SPSS software. The normality of data (mean  $\pm$  standard deviation) was evaluated using Kolmogorov–Smirnov test. The Arcsin transformation was used on proportional and per cent data before running analysis of variance. One-way analysis of variance was applied to find statistical differences. Finally, the comparison of means was done using Tukey test ( $p < 0.01$ ).

## 3 | RESULTS

### 3.1 | GC-Mass spectrometry of *Dracocephalum kotschy* essential oil

The chemical composition of DKE is presented in Table 2.  $\delta$ -3-Carene (10.5%), limonene (9.8%), carvacrol (9.2%), 1,8-cineole (7.3%) and p-cymene (4.2%) composed the main compounds in the chemical composition of DKE (Table 2).

### 3.2 | Growth and mortality parameters

The growth parameters (weight gain per cent, specific growth rate and food conversion ratio) and mortality rate showed no significant differences between the treatments throughout 10 weeks of feeding trial (Table 3,  $p > 0.01$ ).

### 3.3 | Plasma and mucus immune components

The activity of plasma complement and lysozyme and the levels of IgM, total protein and total albumin significantly increased in

TABLE 2 Chemical compositions of *Dracocephalum kotschy* essential oil; the compounds are listed in order of elution time

Compounds	Kovats constant	%
3Z-Hexenol	871	0.3
$\alpha$ -Thujene	935	0.2
$\alpha$ -Pinene	946	2.8
Sabinene	980	2.1
$\beta$ -Pinene	985	2.7
3-Octanol	1000	1.4
$\delta$ -3-Carene	1021	10.5
p-Cymene	1026	4.2
Limonene	1035	9.8
1,8-Cineole	1041	7.3
$\gamma$ -Terpinene	1065	3.1
Terpinolene	1101	3.2
p-Cymenene	1112	2.9
Linalool	1121	2.1
$\beta$ -Thujone	1125	2.5
trans-Pinocarveol	1145	2.2
Camphor	1151	2.4
Pinocarvone	1172	1.8
Isopulegone	1182	1.9
Terpinen-4-ol	1186	3.1
Thuj-3-en-10-al	1189	1.9
trans-Carveol	1222	3.8
Pulegone	1245	2.5
Carvone	1250	5.3
Geraniol	1255	1.4
Limonen-10-ol	1296	1.6
Carvacrol	1305	9.2
Eugenol	1369	1.1
Neryl acetone	1441	0.2
Germacrene D	1491	1.5
Spathulenol	1585	1.4
Total		96.4

response to 0.2 and 0.25 mg DKE/kg diet compared with control and those fed 0.3 mg DKE/kg diet (Table 4,  $p < 0.01$ ). However, the treatment with 0.3 mg DKE/kg diet showed higher activity in plasma complement, lysozyme and total protein compared with control (Table 4,  $p < 0.01$ ). The DKE supplementation increased the bactericidal activity, as the number of CFUs decreased in DKE-supplemented fish (Table 4,  $p < 0.01$ ). In mucus, the protease activity, IgM content and lysozyme activity significantly increased in response to 0.2 and 0.25 mg DKE/kg diet compared with control and fish of 0.3 mg DKE/kg diet treatment (Table 4,  $p < 0.01$ ). However, the mucosal lysozyme activity and IgM content significantly decreased in fish fed 0.3 mg DKE/kg diet compared with other experimental groups (Table 4,  $p < 0.01$ ).

**TABLE 3** Growth and survival of the fish (rainbow trout) after 70 days of supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschy* essential oil (DKE)

	Treatments			
	0 mg DKE (control)	0.2 mg DKE	0.25 mg DKE	0.3 mg DKE
FW (g)	252.5 ± 8.5	260.2 ± 11.2	255.8 ± 9.4	264.1 ± 12.3
WG (%)	358.1 ± 9.5	372.5.5 ± 11.2	365.4 ± 8.7	381.8 ± 10.9
SGR (%/d)	2.16 ± 0.05	2.21 ± 0.04	2.19 ± 0.03	2.23 ± 0.06
FCR	1.2 ± 0.04	1.18 ± 0.07	1.16 ± 0.05	1.21 ± 0.04
SR (%)	86.5 ± 2.5 <sup>a</sup>	90.4 ± 2.3 <sup>a</sup>	92.2 ± 1.5 <sup>a</sup>	70.2 ± 3.6 <sup>b</sup>

Note: Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ). Abbreviations: FCR, Feed conversion rate; FW, Final weight; SGR, Specific growth rate (%/d); SR, Survival rate; WG, Weight gain.

**TABLE 4** The plasma and mucus immune component changes in rainbow trout in response to 70 days supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschy* essential oil (DKE)

	Treatments			
	0 mg DKE (control)	0.2 mg DKE	0.25 mg DKE	0.3 mg DKE
Plasma immune components				
Lysozyme activity (IU/ml)	60.6 ± 12.1 <sup>a</sup>	180.8 ± 15.3 <sup>b</sup>	241.5 ± 20.5 <sup>c</sup>	132.5 ± 13.2 <sup>d</sup>
ACH <sub>50</sub> activity (IU/ml)	1.6 ± 0.3 <sup>a</sup>	4.9 ± 0.5 <sup>b</sup>	5.2 ± 0.3 <sup>b</sup>	3.7 ± 0.6 <sup>c</sup>
IgM (mg/ml)	2.33 ± 0.3 <sup>a</sup>	3.84 ± 0.7 <sup>b</sup>	4.46 ± 0.4 <sup>b</sup>	2.55 ± 0.11 <sup>a</sup>
Bactericidal activity (No. of CFUs)	235.5 ± 22.51 <sup>a</sup>	152.4 ± 19.4 <sup>b</sup>	131.5 ± 12.5 <sup>b</sup>	160.6 ± 20.1 <sup>b</sup>
Total protein (g/dL)	2.2 ± 0.21 <sup>a</sup>	4.3 ± 0.12 <sup>b</sup>	4.5 ± 0.15 <sup>b</sup>	3.1 ± 0.1 <sup>c</sup>
Total albumin (g/dL)	0.73 ± 0.25 <sup>a</sup>	1.42 ± 0.14 <sup>b</sup>	2.65 ± 0.21 <sup>c</sup>	0.9 ± 0.2 <sup>a</sup>
Mucosal immune components				
Lysozyme activity (IU/ml)	21.5 ± 6.3 <sup>a</sup>	42.3 ± 10.2 <sup>b</sup>	50.8 ± 12.7 <sup>b</sup>	32.9 ± 11.5 <sup>ab</sup>
IgM (mg/ml)	0.42 ± 0.12 <sup>a</sup>	0.95 ± 0.1 <sup>b</sup>	1.2 ± 0.15 <sup>b</sup>	0.66 ± 0.18 <sup>a</sup>
Protease activity (IU/ml)	12.5 ± 5.3 <sup>a</sup>	25.2 ± 4.1 <sup>b</sup>	30.3 ± 7.4 <sup>b</sup>	14.6 ± 5.4 <sup>a</sup>
Alkaline phosphatase activity (IU/ml)	2.77 ± 1.5 <sup>a</sup>	8.22 ± 2.02 <sup>b</sup>	10.41 ± 1.1 <sup>b</sup>	4.32 ± 1.9 <sup>a</sup>

Note: Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

**TABLE 5** The haematological alternations of rainbow trout after 70 days of supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschy* essential oil (DKE)

	Treatments			
	0 mg DKE (control)	0.2mg DKE	0.25mg DKE	0.3mg DKE
RBC (×10 <sup>6</sup> /ml)	0.86 ± 0.07 <sup>a</sup>	1.43 ± 0.08 <sup>b</sup>	1.58 ± 0.06 <sup>c</sup>	0.9 ± 0.04 <sup>a</sup>
WBC (×10 <sup>3</sup> /ml)	3.34 ± 0.25 <sup>a</sup>	3.62 ± 0.15 <sup>a</sup>	5.12 ± 0.2 <sup>b</sup>	3.7 ± 0.3 <sup>a</sup>
Hct (%)	26.4 ± 2.5 <sup>a</sup>	30.2 ± 1.42 <sup>b</sup>	31.2 ± 1.4 <sup>b</sup>	28.5 ± 2.1 <sup>a</sup>
Hb (g/dl)	8.92 ± 1.81 <sup>a</sup>	9.34 ± 2.52 <sup>a</sup>	13.61 ± 1.13 <sup>b</sup>	10.12 ± 1.21 <sup>a</sup>
MCV (fl)	212.4 ± 13.5 <sup>a</sup>	221.2 ± 10.1 <sup>a</sup>	268.3 ± 12.4 <sup>b</sup>	230.5 ± 18.1 <sup>a</sup>
MCH (pg)	104.6 ± 6.2 <sup>a</sup>	65.7 ± 7.1 <sup>b</sup>	86.1 ± 5.2 <sup>c</sup>	112.5 ± 8.8 <sup>a</sup>
MCHC (%)	3.1 ± 0.06 <sup>a</sup>	3.12 ± 0.03 <sup>a</sup>	3.24 ± 0.05 <sup>b</sup>	3.06 ± 0.11 <sup>a</sup>

Note: Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

### 3.4 | Haematological parameters

After feeding trial, the values of haematocrit and red blood cell count significantly increased in fish fed with 0.2 and 0.25 mg DKE/kg diet (Table 5,  $p < 0.01$ ). Meanwhile, the mean erythrocyte cell

volume (MCV), haemoglobin content and mean erythrocyte cell haemoglobin content (MCHC) significantly increased in fish fed 0.25 mg DKE/kg diet than those of fish fed the other diets (Table 5,  $p < 0.01$ ). The mean corpuscular haemoglobin (MCH) showed significant decreases in fish fed 0.2 and 0.25 mg DKE/kg diet than compared with



those fed the other diets (Table 5,  $p < 0.01$ ). The values of white blood cell (WBC) significantly increased in response to 0.25 mg DKE/kg diet compared with other diets (Table 5,  $p < 0.01$ ). Also, there were no significant differences in WBC count between control and fish fed 0.3 mg DKE/kg diet (Table 5,  $p > 0.01$ ).

### 3.5 | Liver antioxidant enzymes

The activity of glutathione peroxidase (GPX), catalase (CAT) and superoxide dismutase (SOD) significantly increased in response to 0.25 and 0.3 mg DKE/kg diet compared with control and those fed diet containing 0.2 mg DKE/kg diet (Table 6,  $p < 0.01$ ). In the 0.2 mg DKE/kg diet treatment, only SOD showed significant increases compared with control (Table 6,  $p < 0.01$ ).

### 3.6 | Lipid peroxidation index

As a lipid peroxidation marker, the levels of malondialdehyde (MDA) showed no significant differences between all experimental groups after 60 days feeding trial (Figure 1,  $p > 0.01$ ).

### 3.7 | Gene expressions

After feeding trial, the expression of inflammatory genes, TNF- $\alpha$  and IL-8, significantly decreased in response to 0.2 and 0.25 mg DKE/kg diet (Figure 2,  $p < 0.01$ ). In contrast, these treatments exhibited higher expression of the immune-related genes (C3 and lysozyme) in response to 0.2 and 0.25 mg dietary DKE compared with control and fish from 0.3 mg DKE/kg diet treatment (Figure 3,  $p < 0.01$ ). The expression of C3 and lysozyme showed significant decreases in fish fed 0.3 mg DKE/kg diet compared with control (Figure 3,  $p < 0.01$ ).

### 3.8 | Bacterial challenge

After 10 days of bacterial injection, the cumulative fish mortality (CFM) showed different values among the treatments (Figure 4,  $p < 0.01$ ). The CFM was significantly lower in fish fed DKE compared with control after 10 days bacterial challenge (Figure 4,  $p < 0.01$ ).

## 4 | DISCUSSION

The findings of this study revealed that dietary DKE, especially at concentrations of 0.2 and 0.25 mg DKE/kg diet, can improve the immune and antioxidant system in the rainbow trout. Therefore, DKE can be considered as an immunostimulant, although this role has not yet been reported in fish. The antioxidant properties of *D. kotschyi* are mostly attributed to the flavonoids such as terpinolene, p-Cymene,  $\gamma$ -terpinene, carvone, carvacrol, eugenol and limonene (Ahmad & Beg, 2013; Aydin et al., 2013; Foti & Ingold, 2003; Gülçin, 2011; de Oliveira et al., 2015; Pombal et al., 2017; Yanishlieva et al., 1999).

It is recognized that the compound luteolin plays the main role in the antioxidant activity of the plant (Fattahi et al., 2013; Gohari et al., 2003; Heydari et al., 2019). In this study, the dietary DKE also stimulated the production of antioxidant enzymes (SOD, CAT and GPX) in liver, which clearly indicates the enhancing effects of DKE on the antioxidant defense in the fish. However, we found no changes in the MDA levels as the main indicator of oxidative stress (Del Rio et al., 2005) in response to dietary DKE. In this study, DKE at dietary levels of 0.2 and 0.25 mg DKE/kg elevated immune components in plasma (ACH<sub>50</sub>, IgM content, lysozyme activity, total protein and total albumin) and mucus (protease activity, IgM content and lysozyme activity) and also the expression of some immune-related genes (lysozyme and C3). However, we found no report regarding the immunogenic effects of *D. kotschyi* in the literatures. Some studies have reported a significant decline in secretion of IL-1 $\beta$  from the cells, which may be resulting from the presence of anti-inflammatory compounds in the biochemical composition of *D. kotschyi* (Amirghofran et al., 2011).

The results of the present study support this assumption because we observed down-regulation of the inflammatory genes, TNF- $\alpha$  and IL-8, after supplementation with 0.2 and 0.25 mg DKE/kg diet. The anti-inflammatory effects of *Dracocephalum* sp. has been reported in some studies (Kalantar et al., 2018; Nie et al., 2021; Sadraei et al., 2017; Toshmatov et al., 2019). Some studies have reported the antimicrobial activity of *D. kotschyi* essential oil, which may be related to some compounds such as geranial  $\alpha$ -pinene, geraniol, neral, geraniol acetate and limonene in the biochemical composition of *D. kotschyi* (Dorman & Deans, 2000; Kamali et al., 2015; Singh et al., 2012).

In general, haematology is an important tool to evaluate the fish health in relation to diseases, stress, nutrition and changes in environmental conditions (Burgos-Aceves et al., 2019; Enayat Gholampour et al., 2020; Fazio, 2019; Ghafarifarsani, Hoseinifar,

	Treatments			
	0 mg DKE (control)	0.2 mg DKE	0.25 mg DKE	0.3 mg DKE
SOD (IU/mg)	0.411 $\pm$ 0.08 <sup>a</sup>	0.693 $\pm$ 0.06 <sup>b</sup>	0.752 $\pm$ 0.12 <sup>bc</sup>	0.981 $\pm$ 0.09 <sup>cd</sup>
CAT (IU/mg)	71.23 $\pm$ 13.3 <sup>a</sup>	58.33 $\pm$ 16.45 <sup>a</sup>	112.54 $\pm$ 8.5 <sup>b</sup>	132.5 $\pm$ 11.2 <sup>c</sup>
GPx (IU/mg)	0.121 $\pm$ 0.08 <sup>a</sup>	0.163 $\pm$ 0.1 <sup>a</sup>	0.462 $\pm$ 0.16 <sup>b</sup>	0.714 $\pm$ 0.11 <sup>c</sup>

Note: Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

TABLE 6 The liver antioxidant enzyme changes in the rainbow trout in response to 70 days supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschyi* essential oil (DKE)




FIGURE 1 The changes in lipid peroxidation index (MDA: malondialdehyde levels) in the rainbow trout in response to 70 days of supplementation with diet containing different levels of *Dracocephalum kotschyi* essential oil (DKE). Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ )




FIGURE 2 The expression of cytokine-related genes, TNF- $\alpha$  and IL-8, in the rainbow trout in response to 70 days of supplementation with diet containing different levels of *Dracocephalum kotschyi* essential oil (DKE). Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ )

Aftabgard, et al., 2022; Rashidian, Kajbaf, et al., 2020; Sula et al., 2020). In this study, dietary DKE caused significant increases in the values of Hct, RBCs, MCV, Hb content and MCHC. However, MCH decreased in fish fed 0.2 and 0.25mg DKE/kg diet, which may be due to the depressing effect of DKE on the formation of Hb in red blood cells. In addition, the values of WBCs significantly increased in response to 0.25mg DKE/kg diet, which may be attributed to the immunostimulatory role of DKE. Generally, the effects of plants and their derivatives on fish haematology are different, which may be dependent on plant species, exposure concentration, exposure duration, experimental condition, etc.

In the present study, the supplementation of fish with 0.3mg DKE/kg diet reduced the innate immune responses compared with the 0.2 and 0.25mg DKE/kg diet treatments. Furthermore, the expression of lysozyme and C3 genes in the 0.3mg DKE/kg diet treatment was lower than control. These results may be due to the immunotoxic




FIGURE 3 The expression of immune-related genes, complement (C3) and lysozyme in the rainbow trout in response to 70 days of supplementation with diet containing different levels of *Dracocephalum kotschyi* essential oil (DKE). Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ )




FIGURE 4 Cumulative mortality of rainbow trout, *Oncorhynchus mykiss* ( $n = 15$ /tank), fed different concentration of DKE in the diet throughout 10-day challenge with *Aeromonas hydrophila*. Significant differences are indicated with different lower-case letters (Tukey test,  $p < 0.01$ ). Data: Mean  $\pm$  SD

effects of DKE at high dietary concentrations. The cytotoxic effects of *D. kotschyi* have been previously reported by Ashrafi et al. (2017), Sani et al. (2017) and Shaabani et al. (2020). Ashrafi et al. (2017) indicated that DKE at concentrations of 26.4  $\mu\text{g/ml}$  and 4266.7  $\mu\text{g/ml}$  has toxic effects on HeLa cells and human lymphocytes over in vitro condition. Similar results were observed in the study of Sani et al. (2017), when calu-6 and mehr-80 lung cancer cell lines were exposed to 12.5 200  $\mu\text{g/ml}$  of methanolic extract of *D. kotschyi*. In addition, Shaabani et al. (2020) reported the cytotoxic effects of 400  $\mu\text{g/ml}$  *D. kotschyi* extract on glioblastoma U87 cells over in vitro condition.

In the present study, dietary DKE reduced the mortality induced by *A. hydrophila* infection. However, this reduction was not so considerable. Generally, it is found that the antibacterial effect of DKE is stronger on Gram-positive bacteria (Ashrafi et al., 2017), as it is reported for other plant essential oils (Ballester-Costa et al., 2013; Shakeri et al., 2014). Due to the lack of cell wall in Gram-positive bacteria, the phospholipid bilayer of these bacteria is more



receptive to hydrophobic compounds of essential oils than Gram-negative bacteria (Nikaido, 2003). The contact of essential oils with cell membrane results in disruptions of permeability and bacterial enzyme systems (Wendakoon & Sakaguchi, 1993). Therefore, in this study, the low antibacterial effect of DKE may be due to the Gram-negative bacteria nature of *A. hydrophila*. However, the antibacterial properties of DKE may be due to some compounds including terpinolene,  $\beta$ -Pinene, p-cymene, carvone, carvacrol, eugenol and limonene in its chemical composition (Devi et al., 2010; Han et al., 2020; Marchese et al., 2017; Porfírio et al., 2017; Rondon et al., 2006; Salehi et al., 2019; Xu et al., 2008).

## 5 | CONCLUSION

The finding of the present study indicated that DKE at dietary may improve the immunity and antioxidant defence in the rainbow trout. However, DKE had no appropriate efficiency in reducing the fish mortality induced by *A. hydrophila*.

## ACKNOWLEDGEMENT

The authors are grateful to all the researchers and those who helped to conduct this research and publish the results.

## CONFLICT OF INTEREST

Authors have no conflict of interest to declare for the publication of the present work.

## ETHICAL APPROVAL

All applicable international, national and/or institutional guidelines for the care and use of animals were followed.

## AUTHOR CONTRIBUTIONS

Hafsan Hafsan, Dmitry Bokov and Walid Kamal Abdelbasset contributed to experimental design. Mustafa M. Kadhim, Wanich Suksatan and Mahboubeh Balvardi contributed to methodology and data analysis. Hasan Sh. Majdi, Gunawan Widjaja, Abduladheem Turki Jalil and Maytham T. Qasim contributed to paper writing.

## DATA AVAILABILITY STATEMENT

The datasets in this study are available from the corresponding author on reasonable request. All data and materials are available for publication.

## ORCID

Hafsan Hafsan  <https://orcid.org/0000-0001-5821-0164>

Wanich Suksatan  <https://orcid.org/0000-0003-1797-1260>

Abduladheem Turki Jalil  <https://orcid.org/0000-0001-8403-7465>

## REFERENCES

Ahmad, S., & Beg, Z. H. (2013). Hypolipidemic and antioxidant activities of thymoquinone and limonene in atherogenic suspension fed rats.

*Food Chemistry*, 138(2–3), 1116–1124. <https://doi.org/10.1016/j.foodchem.2012.11.109>

Ahmadifar, E., Sheikhzadeh, N., Roshanaei, K., Dargahi, N., & Faggio, C. (2019). Can dietary ginger (*Zingiber officinale*) alter biochemical and immunological parameters and gene expression related to growth, immunity and antioxidant system in zebrafish (*Danio rerio*)? *Aquaculture*, 507, 341–348. <https://doi.org/10.1016/j.aquaculture.2019.04.049>

Amirghofran, Z., Malek-Hosseini, S., Golmoghaddam, H., Kalantar, F., & Shabani, M. (2011). Inhibition of nitric oxide production and proinflammatory cytokines by several medicinal plants. *Iranian Journal of Immunology*, 8(3), 159–169.

Asghari, G., Nasr Esfahani, B., & Paydar, P. (2015). Evaluating the effect of *Dracocephalum kotschy* methanol extract on *Mycobacterium tuberculosis*. *Research Journal of Pharmacognosy*, 2(3), 31–36.

Ashrafi, B., Ramak, P., Ezatpour, B., & Talei, G. R. (2017). Investigation on chemical composition, antimicrobial, antioxidant, and cytotoxic properties of essential oil from *Dracocephalum kotschy* Boiss. *African Journal of Traditional, Complementary and Alternative Medicines*, 14(3), 209–217. <https://doi.org/10.21010/ajtcam.v14i3.23>

Awad, E., Austin, D., & Lyndon, A. R. (2013). Effect of black cumin seed oil (*Nigella sativa*) and nettle extract (Quercetin) on enhancement of immunity in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquaculture*, 388, 193–197. <https://doi.org/10.1016/j.aquaculture.2013.01.008>

Aydin, E., Türkez, H., & Taşdemir, Ş. (2013). Anticancer and antioxidant properties of terpinolene in rat brain cells. *Archives of Industrial Hygiene and Toxicology*, 64(3), 415–424. <https://doi.org/10.2478/10004-1254-64-2013-2365>

Ballester-Costa, C., Sendra, E., Fernández-López, J., Pérez-Álvarez, J. A., & Viuda-Martos, M. (2013). Chemical composition and in vitro antibacterial properties of essential oils of four *Thymus* species from organic growth. *Industrial Crops and Products*, 50, 304–311. <https://doi.org/10.1016/j.indcrop.2013.07.052>

Bibi, M., Khan, S., Kakar, A.-U.-R., Khan, N., Tareen, A. H., Bibi, S., & Taj, H. (2021). Determination of essential and non-essential elements in *Xylanthemum macropodium* of Balochistan, Pakistan. *Baghdad Journal of Biochemistry and Applied Biological Sciences*, 2(02), 94–103. <https://doi.org/10.47419/bjbabs.v2i02.39>

Blaxhall, P. C., & Daisley, K. W. (1973). Routine haematological methods for use with fish blood. *Journal of Fish Biology*, 5(6), 771–781. <https://doi.org/10.1111/j.1095-8649.1973.tb04510.x>

Brown, R., Moore, L., Mani, A., Patel, S., & Salinas, I. (2021). Effects of ploidy and salmonid alphavirus infection on the skin and gill microbiome of Atlantic salmon (*Salmo salar*). *PLoS One*, 16(2), e0243684. <https://doi.org/10.1371/journal.pone.0243684>

Burgos-Aceves, M. A., Lionetti, L., & Faggio, C. (2019). Multidisciplinary haematology as prognostic device in environmental and xenobiotic stress-induced response in fish. *Science of the Total Environment*, 670, 1170–1183. <https://doi.org/10.1016/j.scitotenv.2019.03.275>

Cao, L., Wang, W., Yang, Y., Yang, C., Yuan, Z., Xiong, S., & Diana, J. (2007). Environmental impact of aquaculture and countermeasures to aquaculture pollution in China. *Environmental Science and Pollution Research-International*, 14(7), 452–462. <https://doi.org/10.1065/espr2007.05.426>

Claiborne, A. (1985). Catalase activity. In R. A. Greenwald (Ed.), *CRC handbook of methods in oxygen radical research* (pp. 283–284). CRC Press.

Cole, D. W., Cole, R., Gaydos, S. J., Gray, J., Hyland, G., Jacques, M. L., Powell-Dunford, N., Sawhney, C., & Au, W. W. (2009). Aquaculture: Environmental, toxicological, and health issues. *International Journal of Hygiene and Environmental Health*, 212(4), 369–377. <https://doi.org/10.1016/j.ijheh.2008.08.003>

Constantino Casas, F., Armijo Ortiz, A., Osorio Sarabia, D., & Chavez Soriano, L. A. (1997). Infection due to *Aeromonas hydrophila* and

- Icthyophthirius multifiliis* in rainbow trout (*Oncorhynchus mykiss*, Walbaum) and tilapia (*Oreochromis aureus*, L) in Morelos, Mexico. Pathological study and review of treatment. *Veterinaria Mexico*, 28(1), 59–62.
- de Oliveira, T. M., de Carvalho, R. B. F., da Costa, I. H. F., de Oliveira, G. A. L., de Souza, A. A., de Lima, S. G., & de Freitas, R. M. (2015). Evaluation of p-cymene, a natural antioxidant. *Pharmaceutical Biology*, 53(3), 423–428.
- Deivi Arunachalam, K., Kuruva, J. K., Pradhoshini, K. P., Musthafa, M. S., & Faggio, C. (2021). Antioxidant and antigenotoxic potential of *Morinda tinctoria* Roxb. leaf extract succeeding cadmium exposure in Asian catfish, *Pangasius sutchi*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 249, 109149.
- Del Rio, D., Stewart, A. J., & Pellegrini, N. (2005). A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutrition, Metabolism and Cardiovascular Diseases*, 15(4), 316–328. <https://doi.org/10.1016/j.numecd.2005.05.003>
- Devi, K. P., Nisha, S. A., Sakthivel, R., & Pandian, S. K. (2010). Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhi* by disrupting the cellular membrane. *Journal of Ethnopharmacology*, 130(1), 107–115. <https://doi.org/10.1016/j.jep.2010.04.025>
- Dorman, H. D., & Deans, S. G. (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, 88(2), 308–316. <https://doi.org/10.1046/j.1365-2672.2000.00969.x>
- Ellis, A. I. (1990). Lysozyme assays. *Techniques in Fish Immunology*, 1, 101–103.
- Enayat Gholampour, T., Fadaei Raieni, R., Pouladi, M., Larijani, M., Pagano, M., & Faggio, C. (2020). The dietary effect of Vitex agnus-castus hydroalcoholic extract on growth performance, blood biochemical parameters, carcass quality, sex ratio and gonad histology in Zebrafish (*Danio rerio*). *Applied Sciences*, 10(4), 1402. <https://doi.org/10.3390/app10041402>
- Farsani, M. N., Hoseinifar, S. H., Rashidian, G., Farsani, H. G., Ashouri, G., & Van Doan, H. (2019). Dietary effects of *Coriandrum sativum* extract on growth performance, physiological and innate immune responses and resistance of rainbow trout (*Oncorhynchus mykiss*) against *Yersinia ruckeri*. *Fish & Shellfish Immunology*, 91, 233–240. <https://doi.org/10.1016/j.fsi.2019.05.031>
- Fattahi, M., Nazeri, V., Torras-Claveria, L., Sefidkon, F., Cusido, R. M., Zamani, Z., & Palazon, J. (2013). Identification and quantification of leaf surface flavonoids in wild-growing populations of *Dracocephalum kotschy* by LC–DAD–ESI–MS. *Food Chemistry*, 141(1), 139–146. <https://doi.org/10.1016/j.foodchem.2013.03.019>
- Fattepur, S., Nilugal, K. C., Rajendran, R., Asmani, F., & Yusuf, E. (2018). Anti-hyperlipidemic activity of methanolic extract of *Boesenbergia pandurata* (finger root) in experimental induced hypercholesterolemic Sprague Dawley rats. *Asian Journal of Pharmaceutical and Clinical Research*, 11(3), 8–12. <https://doi.org/10.22159/ajpcr.2018.v11s3.29962>
- Fazio, F. (2019). Fish hematology analysis as an important tool of aquaculture: A review. *Aquaculture*, 500, 237–242. <https://doi.org/10.1016/j.aquaculture.2018.10.030>
- Foti, M. C., & Ingold, K. U. (2003). Mechanism of inhibition of lipid peroxidation by  $\gamma$ -terpinene, an unusual and potentially useful hydrocarbon antioxidant. *Journal of Agricultural and Food Chemistry*, 51(9), 2758–2765. <https://doi.org/10.1021/jf020993f>
- Ghafariarsani, H., Hoseinifar, S. H., Adorian, T. J., Ferrigolo, F. R. G., Raissy, M., & Van Doan, H. (2021). The effects of combined inclusion of *Malva sylvestris*, *Origanum vulgare*, and *Allium hirtifolium* Boiss for common carp (*Cyprinus carpio*) diet: Growth performance, antioxidant defense, and immunological parameters. *Fish & Shellfish Immunology*, 119, 670–677. <https://doi.org/10.1016/j.fsi.2021.10.014>
- Ghafariarsani, H., Hoseinifar, S. H., Aftabgard, M., & Van Doan, H. (2022). The improving role of savory (*Satureja hortensis*) essential oil for Caspian roach (*Rutilus caspicus*) fry: Growth, haematological, immunological, and antioxidant parameters and resistance to salinity stress. *Aquaculture*, 548, 737653. <https://doi.org/10.1016/j.aquaculture.2021.737653>
- Ghafariarsani, H., Hoseinifar, S. H., Javahery, S., Yazici, M., & Van Doan, H. (2022). Growth performance, biochemical parameters, and digestive enzymes in common carp (*Cyprinus carpio*) fed experimental diets supplemented with vitamin C, thyme essential oil, and quercetin. *Italian Journal of Animal Science*, 21(1), 291–302.
- Ghafariarsani, H., Hoseinifar, S. H., Talebi, M., Yousefi, M., Van Doan, H., Rufchaei, R., & Paolucci, M. (2021). Combined and singular effects of ethanolic extract of persian shallot (*Allium hirtifolium* Boiss) and synbiotic Biomin® IMBO on growth performance, serum-and mucus-immune parameters and antioxidant defense in Zebrafish (*Danio rerio*). *Animals*, 11(10), 2995. <https://doi.org/10.3390/ani1102995>
- Ghafariarsani, H., Kachuei, R., & Imani, A. (2021). Dietary supplementation of garden thyme essential oil ameliorated the deteriorative effects of aflatoxin B<sub>1</sub> on growth performance and intestinal inflammatory status of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 531, 735928. <https://doi.org/10.1016/j.aquaculture.2020.735928>
- Ghafariarsani, H., Rashidian, G., Sheikhlar, A., Naderi Farsani, M., Hoseinifar, S. H., & Van Doan, H. (2021). The use of dietary oak acorn extract to improve haematological parameters, mucosal and serum immunity, skin mucus bactericidal activity, and disease resistance in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Research*, 52(6), 2518–2527.
- Ghavam, M., Manconi, M., Manca, M. L., & Bacchetta, G. (2021). Extraction of essential oil from *Dracocephalum kotschy* Boiss. (Lamiaceae), identification of two active compounds and evaluation of the antimicrobial properties. *Journal of Ethnopharmacology*, 267, 113513.
- Gohari, A. R., Saeidnia, S., Matsuo, K., Uchiyama, N., Yagura, T., Ito, M., Kiuchi, F., & Honda, G. (2003). Flavonoid constituents of *Dracocephalum kotschy* growing in Iran and their trypanocidal activity. *Natural Medicines*, 57(6), 250–252.
- Gondo, H. K. (2021). The effect of spirulina on apoptosis through the caspase-3 pathway in a Preeclamptic Wistar rat model. *Journal of Natural Science, Biology and Medicine*, 12(3), 280–284.
- Gülçin, İ. (2011). Antioxidant activity of eugenol: A structure–activity relationship study. *Journal of Medicinal Food*, 14(9), 975–985. <https://doi.org/10.1089/jmf.2010.0197>
- Han, Y., Sun, Z., & Chen, W. (2020). Antimicrobial susceptibility and antibacterial mechanism of limonene against *Listeria monocytogenes*. *Molecules*, 25(1), 33.
- Harikrishnan, R., & Balasundaram, C. (2005). Modern trends in *Aeromonas hydrophila* disease management with fish. *Reviews in Fisheries Science*, 13(4), 281–320.
- Harikrishnan, R., Balasundaram, C., Dharaneedharan, S., Moon, Y. G., Kim, M. C., Kim, J. S., & Heo, M. S. (2009). Effect of plant active compounds on immune response and disease resistance in *Cirrhina mrigala* infected with fungal fish pathogen, *Aphanomyces Invadans*. *Aquaculture Research*, 40(10), 1170–1181.
- Heydari, P., Yavari, M., Adibi, P., Asghari, G., Ghanadian, S. M., Dida, G. O., & Khamesipour, F. (2019). Medicinal properties and active constituents of *Dracocephalum kotschy* and its significance in Iran: A systematic review. *Evidence-Based Complementary and Alternative Medicine*, 2019, 1–14.
- Jackson, P. B. (1978). Health and disease in intensive aquaculture. *Journal of the South African Veterinary Association*, 49(1), 57–59.
- Kalantar, K., Gholijani, N., Mousaei, N., Yazdani, M., & Amirghofran, Z. (2018). Investigation of *Dracocephalum kotschy* plant extract on the effective inflammatory transcription factors and mediators in activated macrophages. *Anti-Inflammatory & Anti-Allergy Agents*

- in *Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Inflammatory and Anti-Allergy Agents)*, 17(1), 39–49. <https://doi.org/10.2174/1871523017666180608081656>
- Kamali, M., Khosroyar, S., & Mohammadi, A. (2015). Antibacterial activity of various extracts from *Dracocephalum kotschyi* against food pathogenic microorganisms. *International Journal of PharmTech Research*, 8(9), 158–163.
- Kwasek, K., Terova, G., Lee, B. J., Bossi, E., Saroglia, M., & Dabrowski, K. (2014). Dietary methionine supplementation alters the expression of genes involved in methionine metabolism in salmonids. *Aquaculture*, 433, 223–228. <https://doi.org/10.1016/j.aquaculture.2014.05.031>
- LaPatra, S. E., Plant, K. P., Alcorn, S., Ostland, V., & Winton, J. (2010). An experimental vaccine against *Aeromonas hydrophila* can induce protection in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 33(2), 143–151.
- Liu, S., Zhao, H., Lehmler, H. J., Cai, X., & Chen, J. (2017). Antibiotic pollution in marine food webs in Laizhou Bay, North China: Trophodynamics and human exposure implication. *Environmental Science & Technology*, 51(4), 2392–2400. <https://doi.org/10.1021/acs.est.6b04556>
- Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta CT$  method. *Methods*, 25(4), 402–408.
- Lulijwa, R., Rupia, E. J., & Alfaro, A. C. (2020). Antibiotic use in aquaculture, policies and regulation, health and environmental risks: A review of the top 15 major producers. *Reviews in Aquaculture*, 12(2), 640–663. <https://doi.org/10.1111/raq.12344>
- Marchese, A., Arciola, C. R., Barbieri, R., Silva, A. S., Nabavi, S. F., Tsetegho Sokeng, A. J., Izadi, M., Jafari, N. J., Suntar, I., Daglia, M., & Nabavi, S. M. (2017). Update on monoterpenes as antimicrobial agents: A particular focus on p-cymene. *Materials*, 10(8), 947. <https://doi.org/10.3390/ma10080947>
- Marklund, S., & Marklund, G. (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*, 47(3), 469–474. <https://doi.org/10.1111/j.1432-1033.1974.tb03714.x>
- Moradi, H., Ghavam, M., & Tavili, A. (2020). Study of antioxidant activity and some herbal compounds of *Dracocephalum kotschyi* Boiss. in different ages of growth. *Biotechnology Reports*, 25, e00408.
- Nie, L., Li, R., Huang, J., Wang, L., Ma, M., Huang, C., Wu, T., Yan, R., & Hu, X. (2021). Abietane diterpenoids from *Dracocephalum moldavica* L. and their anti-inflammatory activities in vitro. *Phytochemistry*, 184, 112680.
- Nikaido, H. (2003). Molecular basis of bacterial outer membrane permeability revisited. *Microbiology and Molecular Biology Reviews*, 67(4), 593–656. <https://doi.org/10.1128/MMBR.67.4.593-656.2003>
- Nya, E. J., & Austin, B. (2010). Use of bacterial lipopolysaccharide (LPS) as an immunostimulant for the control of *Aeromonas hydrophila* infections in rainbow trout *Oncorhynchus mykiss* (Walbaum). *Journal of Applied Microbiology*, 108(2), 686–694.
- Oriakpono, O., Aduabobo, H., & Wokoma, E. (2012). Biomarker response of a Cichlid Fish *Tilapia guineensis* to crude oil exposure. *International Journal of Modern Cellular and Molecular Biology*, 1(1), 10–18.
- Ortuno, J., Esteban, M. A., & Meseguer, J. (2001). Effects of short-term crowding stress on the gilthead seabream (*Sparus aurata* L.) innate immune response. *Fish & Shellfish Immunology*, 11(2), 187–197. <https://doi.org/10.1006/fsim.2000.0304>
- Othman, Z., Khalep, H. R. H., Abidin, A. Z., Hassan, H., & Fattapur, S. (2019). The anti-angiogenic properties of *Morinda citrifolia* L. (Mengkudu) Leaves using chicken chorioallantoic membrane (CAM) assay. *Pharmacognosy Journal*, 11(1), 12–15.
- Páez-Osuna, F. (2001). The environmental impact of shrimp aquaculture: A global perspective. *Environmental Pollution*, 112(2), 229–231. [https://doi.org/10.1016/S0269-7491\(00\)00111-1](https://doi.org/10.1016/S0269-7491(00)00111-1)
- Pombal, S., Hernández, Y., Diez, D., Mondolis, E., Mero, A., Moran-Pinzon, J., Guerrero, E. I., & Rodilla, J. M. (2017). Antioxidant activity of carvone and derivatives against superoxide ion. *Natural Product Communications*, 12(5), 1934578X1701200502. <https://doi.org/10.1177/1934578X1701200502>
- Porfírio, E. M., Melo, H. M., Pereira, A. M. G., Cavalcante, T. T. A., Gomes, G. A., Carvalho, M. G. D., Costa, R. A., & Júnior, F. E. A. C. (2017). In vitro antibacterial and antibiofilm activity of *Lippia alba* essential oil, citral, and carvone against *Staphylococcus aureus*. *The Scientific World Journal*, 2017, 4962707.
- Pradeepkiran, J. A. (2019). Aquaculture role in global food security with nutritional value: A review. *Translational Animal Science*, 3(2), 903–910. <https://doi.org/10.1093/tas/txz2012>
- Qahir, A., Kakar, A.-R., Khan, N., Khan, S., Hakeem, A., Kamal, R., & Fazal-Ur-Rehman. (2021). The antioxidant, antimicrobial, and clinical effects with elemental contents of pomegranate (*Punica granatum*) peel extracts: A review. *Baghdad Journal of Biochemistry and Applied Biological Sciences*, 2(01), 21–28. <https://doi.org/10.47419/bjbabs.v2i01.33>
- Raissy, M., Ghafarifarsani, H., Hoseinifar, S. H., El-Haroun, E. R., Naserabadi, S. S., & Van Doan, H. (2022). The effect of dietary combined herbs extracts (oak acorn, coriander, and common mallow) on growth, digestive enzymes, antioxidant and immune response, and resistance against *Aeromonas hydrophila* infection in common carp, *Cyprinus Carpio*. *Aquaculture*, 546, 737287. <https://doi.org/10.1016/j.aquaculture.2021.737287>
- Rao, Y. V., Das, B. K., Jyotirmayee, P., & Chakrabarti, R. (2006). Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 20(3), 263–273. <https://doi.org/10.1016/j.fsi.2005.04.006>
- Rashidian, G., Bahrami Gorji, S., Farsani, M. N., Prokić, M. D., & Faggio, C. (2020). The oak (*Quercus brantii*) acorn as a growth promoter for rainbow trout (*Oncorhynchus mykiss*): growth performance, body composition, liver enzymes activity and blood biochemical parameters. *Natural Product Research*, 34(17), 2413–2423.
- Rashidian, G., Boldaji, J. T., Rainis, S., Prokić, M. D., & Faggio, C. (2021). Oregano (*Origanum vulgare*) extract enhances zebrafish (*Danio rerio*) growth performance, serum and mucus innate immune responses and resistance against *Aeromonas hydrophila* challenge. *Animals*, 11(2), 299. <https://doi.org/10.3390/ani11020299>
- Rashidian, G., Kajbaf, K., Prokić, M. D., & Faggio, C. (2020). Extract of common mallow (*Malva sylvestris*) enhances growth, immunity, and resistance of rainbow trout (*Oncorhynchus mykiss*) fingerlings against *Yersinia ruckeri* infection. *Fish & Shellfish Immunology*, 96, 254–261. <https://doi.org/10.1016/j.fsi.2019.12.018>
- Romero, J., Feijóo, C. G., & Navarrete, P. (2012). Antibiotics in aquaculture—use, abuse and alternatives. In E. D. Carvalho, G. S. David, & R. J. Silva (Eds.), *Health and environment in aquaculture* (p. 159). IntechOpen.
- Rondon, M., Velasco, J., Hernandez, J., Pecheneda, M., Rojas, J., Morales, A., Carmona, J., & Diaz, T. (2006). Chemical composition and antibacterial activity of the essential oil of *Tagetes patula* L. (Asteraceae) collected from the Venezuela Andes. *Revista Latinoamericana De Química*, 34(1/3), 32.
- Roslan, N., & Shariff, H. (2021). Potential pharmacotherapy of *Olea europaea* (Olive) fruit oil against neurotoxicity induced methamphetamine. *Journal of Cellular and Molecular Anesthesia*, 4(2), 67–68.
- Ross, N. W., Firth, K. J., Wang, A., Burka, J. F., & Johnson, S. C. (2000). Changes in hydrolytic enzyme activities of naive Atlantic salmon *Salmo salar* skin mucus due to infection with the salmon louse *Lepeophtheirus salmonis* and cortisol implantation. *Diseases of Aquatic Organisms*, 41(1), 43–51.
- Sadraei, H., Asghari, G., Khanabadi, M., & Minaiyan, M. (2017). Anti-inflammatory effect of apigenin and hydroalcoholic extract of *Dracocephalum kotschyi* on acetic acid-induced colitis in rats.



- Research in Pharmaceutical Sciences, 12(4), 322. <https://doi.org/10.4103/1735-5362.212050>
- Salehi, B., Upadhyay, S., Erdogan Orhan, I., Kumar Jugran, A., L.D. Jayaweera, S., A. Dias, D., Sharopov, F., Taheri, Y., Martins, N., Baghalpour, N., C. Cho, W., & Sharifi-Rad, J. (2019). Therapeutic potential of  $\alpha$ - and  $\beta$ -pinene: A miracle gift of nature. *Biomolecules*, 9(11), 738. <https://doi.org/10.3390/biom9110738>
- Sani, T. A., Mohammadpour, E., Mohammadi, A., Memariani, T., Yazdi, M. V., Rezaee, R., Calina, D., Docea, A. O., Goumenou, M., Etemad, L., & Shahsavand, S. (2017). Cytotoxic and apoptogenic properties of *Dracocephalum kotschyi* aerial part different fractions on calu-6 and mehr-80 lung cancer cell lines. *Farmacía*, 65(2), 189–199.
- Shaabani, M., Mousavi, S. H., Azizi, M., & Jafari, A. A. (2020). Cytotoxic and apoptogenic effects of *Dracocephalum kotschyi* Boiss. extracts against human glioblastoma U87 cells. *Avicenna Journal of Phytomedicine*, 10(6), 594.
- Shakeri, A., Khakdan, F., Soheili, V., Sahebkar, A., Rassam, G., & Asili, J. (2014). Chemical composition, antibacterial activity, and cytotoxicity of essential oil from *Nepeta ucrainica* L. spp. kopetdaghensis. *Industrial Crops and Products*, 58, 315–321. <https://doi.org/10.1016/j.indcrop.2014.04.009>
- Shalaby, M. N. (2015). Effects of protein hydrolysates on physical performance and immunity in male soccer players. *The International Scientific Journal of Physical Education and Sport Sciences*, 2(2), 1–8.
- Shalaby, M. N. (2018). The effect of whey protein (Natural Nanoparticle) on muscle strength, GH, IGF, T. Protein and body composition. *International Journal of Pharmaceutical Research & Allied Sciences*, 7(1), 126–132.
- Shalaby, M., Sakoury, M. M. A., Harthi, S. M., Alshalawi, F. M., Alhajji, M. M., Alshaikh, Z. H., & Aljaber, A. H. (2020). Vitamin D3 for health and muscle functions of athletes. *Systematic Reviews in Pharmacy*, 11(9), 851–854.
- Shalaby, M. N., Sakoury, M. M., Kholif, M. A., & Alsayed, N. I. (2020). The role of Amino Acids in improving immunity and growth factors of Volleyball players. *Journal of Advanced Pharmacy Education and Research*, 10(4), 140–144.
- Singh, S. K., Vishnoi, R., Dhingra, G. K., & Kishor, K. (2012). Antibacterial activity of leaf extracts of some selected traditional medicinal plants of Uttarakhand, North East India. *Journal of Applied and Natural Science*, 4(1), 47–50. <https://doi.org/10.31018/jans.v4i1.220>
- Siwicki, A. K., & Anderson, D. P. (1993). Nonspecific defense mechanisms assay in fish. II. Potential killing activity of neutrophils and macrophages, lysozyme activity in serum and organs and total immunoglobulin level in serum. In A. K. Siwicki, D. P. Anderson, & J. Waluga (Eds.), *Fish disease diagnosis and prevention methods* (pp. 105–112). Wydawnictwo Instytutu Rybactwa Strodoladowego.
- Subramanian, S., MacKinnon, S. L., & Ross, N. W. (2007). A comparative study on innate immune parameters in the epidermal mucus of various fish species. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 148(3), 256–263. <https://doi.org/10.1016/j.cbpb.2007.06.003>
- Sula, E., Aliko, V., Pagano, M., & Faggio, C. (2020). Digital light microscopy as a tool in toxicological evaluation of fish erythrocyte morphological abnormalities. *Microscopy Research and Technique*, 83(4), 362–369. <https://doi.org/10.1002/jemt.23422>
- Talari, M., Seydi, E., Salimi, A., Mohsenifar, Z., Kamalinejad, M., & Pourahmad, J. (2014). *Dracocephalum*: novel anticancer plant acting on liver cancer cell mitochondria. *BioMed Research International*, 1–10. <https://doi.org/10.1155/2014/892170>
- Toshmatov, Z. O., Li, J., Eshbakova, K. A., Tang, D., Xin, X., & Aisa, H. A. (2019). New monoterpene glucosides from *Dracocephalum komarovii* and their anti-inflammatory activity. *Phytochemistry Letters*, 33, 102–105. <https://doi.org/10.1016/j.phytol.2019.08.005>
- Utley, H. G., Bernheim, F., & Hochstein, P. (1967). Effect of sulfhydryl reagents on peroxidation in microsomes. *Archives of Biochemistry and Biophysics*, 118(1), 29–32. [https://doi.org/10.1016/0003-9861\(67\)90273-1](https://doi.org/10.1016/0003-9861(67)90273-1)
- Van Doan, H., Hoseinifar, S. H., Ringø, E., Ángeles Esteban, M., Dadar, M., Dawood, M. A., & Faggio, C. (2020). Host-associated probiotics: A key factor in sustainable aquaculture. *Reviews in Fisheries Science & Aquaculture*, 28(1), 16–42. <https://doi.org/10.1080/23308249.2019.1643288>
- Wayne, P. (2007). *Implementation guide of POCT for HEALTH care providers* (pp. 1–37). CLSI.
- Wendakoon, C. N., & Sakaguchi, M. (1993). Combined effect of sodium chloride and clove on growth and biogenic amine formation of *Enterobacter aerogenes* in mackerel muscle extract. *Journal of Food Protection*, 56(5), 410–413. <https://doi.org/10.4315/0362-028X-56.5.410>
- Xu, J., Zhou, F., Ji, B. P., Pei, R. S., & Xu, N. (2008). The antibacterial mechanism of carvacrol and thymol against *Escherichia coli*. *Letters in Applied Microbiology*, 47(3), 174–179.
- Yanishlieva, N. V., Marinova, E. M., Gordon, M. H., & Raneva, V. G. (1999). Antioxidant activity and mechanism of action of thymol and carvacrol in two lipid systems. *Food Chemistry*, 64(1), 59–66. [https://doi.org/10.1016/S0308-8146\(98\)00086-7](https://doi.org/10.1016/S0308-8146(98)00086-7)
- You, S. H., Yoon, M. Y., & Moon, J. S. (2021). Antioxidant and anti-inflammatory activity study of fulvic acid. *Journal of Natural Science, Biology and Medicine*, 12(3), 285–289.
- Yousefi, M., Farsani, M. N., Ghafarifarsani, H., Hoseinifar, S. H., & Van Doan, H. (2021). The effects of dietary supplementation of mistletoe (*Viscum album*) extract on the growth performance, antioxidant, and innate, immune responses of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 536, 736385. <https://doi.org/10.1016/j.aquaculture.2021.736385>
- Yousefi, M., Ghafarifarsani, H., Hoseini, S. M., Hoseinifar, S. H., Abtahi, B., Vatnikov, Y. A., Kulikov, E. V., & Van Doan, H. (2022). Effects of dietary thyme essential oil and prebiotic administration on rainbow trout (*Oncorhynchus mykiss*) welfare and performance. *Fish & Shellfish Immunology*, 120, 737–744. <https://doi.org/10.1016/j.fsi.2021.12.023>
- Yousefi, M., Ghafarifarsani, H., Hoseinifar, S. H., Rashidian, G., & Van Doan, H. (2021). Effects of dietary marjoram, *Origanum majorana* extract on growth performance, hematological, antioxidant, humoral and mucosal immune responses, and resistance of common carp, *Cyprinus carpio* against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 108, 127–133. <https://doi.org/10.1016/j.fsi.2020.11.019>
- Zharif, N., Santosh, F., Kiran, C. N., Fadli, A., Ibrahim, A., & Nizam, G. (2018). Synergistic effect of ethanolic extract of *Melastoma malabataricum* leaves and antibiotics. *International Journal of Medical Toxicology & Legal Medicine*, 21(3and4), 167–170. <https://doi.org/10.5958/0974-4614.2018.00059.1>

**How to cite this article:** Hafsan, H., Bokov, D., Abdelbasset, W. K., Kadhim, M. M., Suksatan, W., Majdi, H. S., Widjaja, G., Jalil, A. T., Qasim, M. T., & Balvardi, M. (2022). Dietary *Dracocephalum kotschyi* essential oil improved growth, haematology, immunity and resistance to *Aeromonas hydrophila* in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Research*, 53, 3164–3175. <https://doi.org/10.1111/are.15829>

---

## Aquaculture Research - Account Modified in ScholarOne Manuscripts

2 messages

---

**Aquaculture Research** <onbehalf@manuscriptcentral.com>  
Reply-To: AREditorialOffice@wiley.com  
To: hafsan.bio@uin-alauddin.ac.id

Mon, Feb 14, 2022 at 1:07 PM

14-Feb-2022

Dear Dr. Hafsan:

This e-mail is a notification that your account on Aquaculture Research - ScholarOne Manuscripts (previously known as Manuscript Central) site has been modified. Your USER ID for your account at <https://mc.manuscriptcentral.com/are> is as follows:

USER ID: [hafsan.bio@uin-alauddin.ac.id](mailto:hafsan.bio@uin-alauddin.ac.id)

PASSWORD: If you are unsure of your password you can click the link below which will take you directly to the option for setting a new password.

[https://mc.manuscriptcentral.com/are?URL\\_MASK=a1ccf51c94784ec6aa301aaf4dc3e1ef](https://mc.manuscriptcentral.com/are?URL_MASK=a1ccf51c94784ec6aa301aaf4dc3e1ef)

Thank you for your participation.

Yours Sincerely,

Editorial Office  
Aquaculture Research

---

**Aquaculture Research** <onbehalf@manuscriptcentral.com>  
Reply-To: AREditorialOffice@wiley.com  
To: hafsan.bio@uin-alauddin.ac.id

Mon, Feb 14, 2022 at 1:12 PM

14-Feb-2022

Dear Dr. Hafsan:

This e-mail is a notification that your account on Aquaculture Research - ScholarOne Manuscripts (previously known as Manuscript Central) site has been modified. Your USER ID for your account at <https://mc.manuscriptcentral.com/are> is as follows:

USER ID: [hafsan.bio@uin-alauddin.ac.id](mailto:hafsan.bio@uin-alauddin.ac.id)

PASSWORD: If you are unsure of your password you can click the link below which will take you directly to the option for setting a new password.

[https://mc.manuscriptcentral.com/are?URL\\_MASK=ca83ae7b5d6a48488b3df8360200c459](https://mc.manuscriptcentral.com/are?URL_MASK=ca83ae7b5d6a48488b3df8360200c459)

[Quoted text hidden]



### My Articles

Articles missing?

Filter my articles

1 article



Aquaculture Research



Dietary *Dracocephalum kotschy* essential oil improved growth, haematology, immunity and resistance to *Aeromonas hydrophila* in rainbow trout (*Oncorhynchus mykiss*)

DOI: <https://doi.org/10.1111/are.15829>

Status: Published

Manage Article

### Find a home for your next article

Wiley partners with the world's best research leaders and organizations to publish hundreds of top-tier journals.

Find a journal

Help

New blog and video content from The Wiley Network



---

**Information: Production Editor Contact ARE - AQUACULTURE RESEARCH |  
Dietary Dracocephalum kotschyi essential oil improved growth, hematology,  
immunity and resistance to Aeromonas hydrophila in rainbow trout  
(Oncorhynchus mykiss)**

1 message

---

**wiley.bookin@sps.co.in** <wiley.bookin@sps.co.in>  
To: hafsan.bio@uin-alauddin.ac.id  
Cc: iselvaraj@wiley.com

Tue, Mar 8, 2022 at 4:04 PM

Dear Author,

We are in the process of preparing "Dietary Dracocephalum kotschyi essential oil improved growth, hematology, immunity and resistance to Aeromonas hydrophila in rainbow trout (Oncorhynchus mykiss)" for publication. Your Production Editor, INDHUMATHI SELVARAJ, will support you and your article throughout the process.

Please get in touch with your Production Editor at [are@wiley.com](mailto:are@wiley.com) if you have any questions.

Sincerely,

Booking-in Team On behalf of Wiley

Article ID: ARE\_15829  
Article DOI: 10.1111/ARE.15829

---

## Action: Proof of ARE\_EV\_ARE15829 for Aquaculture Research ready for review

1 message

---

**Wiley Online Proofing** <onlineproofing@eproofing.in>  
Reply-To: eproofing@wiley.com  
To: hafsan.bio@uin-alauddin.ac.id

Sat, Mar 12, 2022 at 6:14 AM



## Review your proof

ARE\_EV\_ARE15829

Dear **Hafsan Hafsan**,

The proof of your **Aquaculture Research** article **Dietary Dracocephalum kotschy essential oil improved growth, haematology, immunity and resistance to Aeromonas hydrophila in rainbow trout (Oncorhynchus mykiss)** is now available for review:

### Edit Article

To review your article, please complete the following steps, ideally within 48 hours\*, so we can publish your article as quickly as possible.

1. Open your proof in the online proofing system using the button above.
2. Check the article for correctness and respond to all queries. For instructions on using the system, please see the "Help" menu in the upper right corner.
3. Submit your changes by clicking the "Submit" button in the proofing system.
4. If you have not already done so, please log in into Author Services (<https://authorservices.wiley.com>) and click on "My Dashboard". When you locate your article on the Dashboard, please click the button that says "Sign License" you can then complete your license agreement online. **Your article cannot be published until we have received your completed license agreement.**

**NOTE: There are fees associated with the publication of your article. Please download and follow the instructions on the charge form(s) located by clicking the "Forms" button.**

#### Helpful Tips

- Your manuscript has been formatted following the style requirements for the journal.


Any requested changes that go against journal style will not be made.

- Your proof will include queries. These must be replied to using the system before the proof can be submitted.
- The only acceptable changes at this stage are corrections to grammatical errors or data accuracy, or to provide higher resolution figure files (if requested by the typesetter).
- Any changes to scientific content or authorship will require editorial review and approval.
- Once your changes are complete, submit the article after which no additional corrections can be requested.
- Most authors complete their corrections within 48 hours. Returning any corrections promptly will accelerate publication of your article.

If you encounter any problems or have questions, please contact us at ([iselvaraj@wiley.com](mailto:iselvaraj@wiley.com)). For the quickest response, include the journal name and your article ID (found in the subject line) in all correspondence.

Best regards,  
Aquaculture Research

\* We appreciate that the COVID-19 pandemic may create conditions for you that make it difficult for you to review your proof within standard timeframes. If you have any problems keeping to this schedule, please reach out to me at ([iselvaraj@wiley.com](mailto:iselvaraj@wiley.com)) to discuss alternatives.



B I U x<sup>2</sup> x<sub>2</sub> ← → Ω ∞ Zoom Σ Insert

Sections

ORIGINAL ARTICLE

**Dietary *Dracocephalum kotschyi* essential oil improved growth, haematology, immunity and resistance to *Aeromonas hydrophila* in rainbow trout *Oncorhynchus mykiss*)**

**Hafsan Hafsan**<sup>1,\*</sup> | **Dmitry Bokov**<sup>2,3</sup> | **Walid Kamal Abdelbasset**<sup>4,5</sup> | **Mustafa M. Kadhim**<sup>6,7,8</sup> | **Wanich Suksatan**<sup>9</sup> | **Hasan Sh. Majidi**<sup>10</sup> | **Gunawan Widjaja**<sup>11,12</sup> | **Abduladheem Turki Jalili**<sup>13,14</sup> | **Maytham T. Qasim**<sup>15,16</sup> | **Mahboubeh Balvardi**<sup>17</sup>

- <sup>1</sup> Biology Department| Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Gowa, Indonesia
- <sup>2</sup> Institute of Pharmacy, Sechenov First Moscow State Medical University, Moscow, Russian Federation
- <sup>3</sup> Laboratory of Food Chemistry, Federal Research Center of Nutrition, Biotechnology and Food Safety, Moscow, Russian Federation
- <sup>4</sup> Department of Health and Rehabilitation Sciences| College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Al Kharj, Saudi Arabia
- <sup>5</sup> Department of Physical Therapy| Kasr Al-Aini Hospital, Cairo University, Giza, Egypt
- <sup>6</sup> Department of Dentistry, Kut University College, Kut, Iraq
- <sup>7</sup> College of Technical Engineering, The Islamic University, Najaf, Iraq
- <sup>8</sup> Department of Pharmacy, Osol Aldeen University College, Baghdad, Iraq
- <sup>9</sup> Faculty of Nursing| HRH Princess Chulabhorn College of Medical Science| Faculty of Nursing, HRH Princess Chulabhorn College of Medical Science, Chulabhorn Royal Academy, Bangkok, Thailand
- <sup>10</sup> Department of Chemical Engineering and Petroleum Industries, Al-Mustaqbal University College, Babylon, Iraq
- <sup>11</sup> Faculty of Public Health, Universitas Indonesia, Jakarta Depok, Indonesia
- <sup>12</sup> Faculty of Law, Universitas Krisnadwipayana Indonesia, Jakarta atiwaringin, Indonesia
- <sup>13</sup> Faculty of Biology and Ecology, Yanka Kupala State University of Grodno, Grodno, Belarus
- <sup>14</sup> College of Technical Engineering, The Islamic University, Najaf, Iraq
- <sup>15</sup> Ministry of Education, Directorate Thi-Qar Education, Thi-Qar, Iraq
- <sup>16</sup> Department of Anesthesia| College of Health and Medical Technology, Al-Ayen University, Thi-Qar, Iraq
- <sup>17</sup> Department of Biology, Payamnoor University| Payame Noor University, Tehran, Iran

Correspondence

Hafsan Hafsan, Biology Department, Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Sultan Alauddin Street, Gowa 92118, Indonesia.  
Email: hafsan.bio@uin-alauddin.ac.id

Abstract

In the present study, the effect of *Dracocephalum kotschyi* essential oil (DKE) was investigated on growth, haematology, immune and antioxidant defence system and resistance of rainbow trout juveniles to bacterial infection (*Aeromonas hydrophila*). For this purpose, the fish were fed a diet containing different concentrations of DKE including 0 (control), 0.2, 0.25 and 0.3 mg/kg diet in three replicates for 60 days. After feeding trial, the fish were challenged with a pathogenic dose of *A. hydrophila*. Based on the results, immune components in plasma (alternative complement activity [ACH<sub>50</sub>], IgM content, lysozyme activity, total protein and total albumin) and mucus (protease activity, IgM content and lysozyme activity) significantly elevated in fish fed diet containing 0.2 and 0.25 mg DKE/kg diet compared with other groups (*p* < 0.01). The plasma bactericidal activity increased in all DKE-supplemented fish. Supplementation of fish with 0.3 mg DKE/kg diet depressed the mucosal lysozyme activity and IgM content. The liver antioxidant enzymes, glutathione peroxidase (GPx), catalase (CAT) and superoxide dismutase (SOD) showed increased activity in response to 0.25 and 0.3 mg DKE/kg diet compared with other experimental diets

Journal & Article Information

Queries (26)

**AQ5:** "AUTHOR: Reference Hoseinifar e al., 2020 has not been included in the Reference List, please supply full publication details."

it was deleted

**AQ6:** "AUTHOR: The sentence, '*Dracocephalum kotschyi* Boiss ...' has been slightly modified for clarity. Kindly check if the intended meaning of the sentence is retained."

it is ok

Comments (0)

Image Comments (0)

Edits (35)

Supplementary Files (0)

Attachments (0)

# Dietary *Dracocephalum kotschyi* essential oil improved growth, haematology, immunity and resistance to *Aeromonas hydrophila* in rainbow trout (*Oncorhynchus mykiss*)

AQ1

AQ2

**Hafsan Hafsan**<sup>1,\*</sup>  | **Dmitry Bokov**<sup>2,3</sup> | **Walid Kamal Abdelbasset**<sup>4,5</sup> | **Mustafa M. Kadhim**<sup>6,7,8</sup> | **Wanich Suksatan**<sup>9</sup>  | **Hasan Sh. Majdi**<sup>10</sup> | **Gunawan Widjaja**<sup>11,12</sup> | **Abduladheem Turki Jalil**<sup>13,14</sup>  | **Maytham T. Qasim**<sup>15,16</sup> | **Mahboubeh Balvardi**<sup>17</sup>

<sup>1</sup> Biology Department | Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Gowa, Indonesia

<sup>2</sup> Institute of Pharmacy, Sechenov First Moscow State Medical University, Moscow, Russian Federation

<sup>3</sup> Laboratory of Food Chemistry, Federal Research Center of Nutrition, Biotechnology and Food Safety, Moscow, Russian Federation

<sup>4</sup> Department of Health and Rehabilitation Sciences | College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Al Kharj, Saudi Arabia

<sup>5</sup> Department of Physical Therapy | Kasr Al-Aini Hospital, Cairo University, Giza, Egypt

<sup>6</sup> Department of Dentistry, Kut University College, Kut, Iraq

<sup>7</sup> College of Technical Engineering, The Islamic University, Najaf, Iraq

<sup>8</sup> Department of Pharmacy, Osol Aldeen University College, Baghdad, Iraq

<sup>9</sup> ~~Faculty of Nursing | HRH Princess Chulabhorn College of Medical Science~~ Faculty of Nursing, HRH Princess Chulabhorn College of Medical Science, Chulabhorn Royal Academy, Bangkok, Thailand

<sup>10</sup> Department of Chemical Engineering and Petroleum Industries, Al-Mustaqbal University College, Babylon, Iraq

<sup>11</sup> Faculty of Public Health, Universitas Indonesia, JakartaDepok, Indonesia

<sup>12</sup> Faculty of Law, Universitas Krisnadwipayana Indonesia, JakartaJatiwaringin, Indonesia

<sup>13</sup> Faculty of Biology and Ecology, Yanka Kupala State University of Grodno, Grodno, Belarus

<sup>14</sup> College of Technical Engineering, The Islamic University, Najaf, Iraq

<sup>15</sup> Ministry of Education, Directorate Thi-Qar Education, Thi-Qar, Iraq

<sup>16</sup> Department of Anesthesia | College of Health and Medical Technology, Al-Ayen University, Thi-Qar, Iraq

<sup>17</sup> Department of Biology, ~~Payamnoor University~~ Payame Noor University, Tehran, Iran

## Correspondence

Hafsan Hafsan, Biology Department, Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Sultan Alauddin Street, Gowa 92118, Indonesia.

Email: [hafsan.bio@uin-alauddin.ac.id](mailto:hafsan.bio@uin-alauddin.ac.id)

AQ4

## Abstract

In the present study, the effect of *Dracocephalum kotschy* essential oil (DKE) was investigated on growth, haematology, immune and antioxidant defence system and resistance of rainbow trout juveniles to bacterial infection (*Aeromonas hydrophila*). For this purpose, the fish were fed a diet containing different concentrations of DKE including 0 (control), 0.2, 0.25 and 0.3 mg/kg diet in three replicates for 60 days. After feeding trial, the fish were challenged with a pathogenic dose of *A. hydrophila*. Based on the results, immune components in plasma (alternative complement activity [ACH<sub>50</sub>], IgM content, lysozyme activity, total protein and total albumin) and mucus (protease activity, IgM content and lysozyme activity) significantly elevated in fish fed diet containing 0.2 and 0.25 mg DKE/kg diet compared with other groups ( $p < 0.01$ ). The plasma bactericidal activity increased in all DKE-supplemented fish. Supplementation of fish with 0.3 mg DKE/kg diet depressed the mucosal lysozyme activity and IgM content. The liver antioxidant enzymes, glutathione peroxidase (GPx), catalase (CAT) and superoxide dismutase (SOD) showed increased activity in response to 0.25 and 0.3 mg DKE/kg diet compared with other experimental diets ( $p < 0.01$ ). Dietary DKE changed the haematology of the fish. The haematocrit, red blood cell count (RBC), white blood cell (WBC), haemoglobin content and mean erythrocyte cell haemoglobin content (MCHC) increased in response to DKE. In contrast, the mean corpuscular haemoglobin (MCH) decreased in fish fed 0.2 and 0.25 mg DKE/kg diet compared with other diets ( $p < 0.01$ ). The supplementation of fish with 0.2 and 0.25 mg DKE/kg diet decreased the expression of cytokine-related genes, TNF- $\alpha$  and IL-8. In contrast, the DKE up-regulated the expression of the immune-related genes, C3 and lysozyme. DKE at concentration of 0.3 mg/kg diet



depressed the C3 and lysozyme gene expressions. The DKE supplementation decreased the mortality rate of the fish after bacterial challenge compared with non-supplemented ones ( $p < 0.01$ ). In conclusion, the findings of this study revealed the immune-stimulating effects of DKE at optimized dietary concentrations. In addition, DKE decreased the mortality induced by *A. hydrophila* infection.

## Keywords

Bacterial | *Dracocephalum kotschyi* | Essential oil | Fish | Immunity

## 1 | INTRODUCTION

Aquaculture is known as one of the most important ways to provide the protein needs of human (Pradeepkiran, 2019). However, the development of intensive aquaculture and super-intensive aquaculture has increased the risk of fish diseases as well as environmental pollutions (Cao et al., 2007; Cole et al., 2009; Jackson, 1978; Páez-Osuna, 2001; Romero et al., 2012). Today, antibiotics are widely used to treat fish diseases, but there are still some problems with these chemicals that may limit their use (Lulijwa et al., 2020; Romero et al., 2012). Continued use of antibiotics may create resistant bacterial strains, and in addition, the accumulation of these chemicals and their derivatives in fish meat can threaten human health as a consumer (Brown et al., 2021; Lulijwa et al., 2020). Furthermore, the release of antibiotics into natural aquatic environments can have a negative impact on the flora and fauna and ultimately the entire food web of that environment (Liu et al., 2017; Lulijwa et al., 2020). Owing to these problems, it seems necessary to replace antibiotics with safe alternatives. Dietary supplements, including plant-based supplements, are currently used to keep health and to treat many diseases in humans and animals (Fattepur et al., 2018; Gondo, 2021; Othman et al., 2019; Roslan & Shariff, 2021; Shalaby, 2015, 2018; Shalaby, Sakoury, Harthi, et al., 2020; Shalaby, Sakoury, Kholif, et al., 2020; You et al., 2021; Zharif et al., 2018). During the last decade, many studies have reported the medicinal, antioxidant, immunogenic and antibacterial properties of medicinal plants and their derivatives in fish (Ahmadifar et al., 2019; Farsani et al., 2019; Rashidian, Bahrami Gorji, et al., 2020; Van Doan et al., 2020; Bibi et al., 2021; DeiviArunachalam et al., 2021; Ghafarifarsani Kachuei et al., 2021; Ghafarifarsani, Hoseinifar, Adorian, et al., 2021; Ghafarifarsani, Hoseinifar, Talebi, et al., 2021; Ghafarifarsani, Qahir et al., 2021; Rashidian, et al., 2021; ~~Hoseinifar et al., 2020~~; Rashidian et al., 2021; Yousefi, Farsani, et al., 2021; Yousefi, Ghafarifarsani et al., 2021; Raissy et al., 2022; Yousefi et al., 2022). However, the identification and introduction of the new species with immunogenic and antibacterial properties can be useful for

AQ6

aquaculture enhancement. *Dracocephalum* with more than 60 species is native to temperate regions of Asia and Europe, which is consumed in many parts of the world due to its high medicinal properties (Heydari et al., 2019). *Dracocephalum kotschy* Boiss plant is one of the well-known species of *Dracocephalum*, which is used in treatment of infectious diseases due to its medicinal properties and active compounds. The chemical composition of *Dracocephalum kotschy* includes bioactive compounds with anti-cancerous, antioxidant, antibacterial, anti-nociceptive, anti-hyperlipidaemic, anti-spasmodic, cytotoxic and immunostimulatory effects (Heydari et al., 2019; Moradi et al., 2020; Sadraei et al., 2017; Talari et al., 2014). In a study by Ghavam et al. (2021), the antibacterial properties of *D. kotschy* essential oil (DKE) were attributed to some bioactive compounds in the biochemical composition of DKE including  $\alpha$ -pinene, terpinen-4-ol, limonene, neral and geraniol. In addition, the results of Asghari et al. (2015) showed that the methanol extracts of *D. kotschy* inhibit the growth of *Mycobacterium tuberculosis*. The extracts of *D. kotschy* showed antibacterial effects against Gram-positive bacteria (*Staphylococcus aureus*, *Bacillus cereus* and *Listeria Monocytogenes*) and Gram-negative bacteria (*Salmonella enterica*, *Escherichia coli* and *Enterica Aerogenes*) (Kamali et al., 2015). According to a literature review, we found no data regarding the immunogenic, antioxidant and antibacterial effects of *D. kotschy* in fish. Therefore, the present study was conducted to evaluate the effects of DKE on immune and antioxidant defence system and resistance of rainbow trout juveniles to infection induced by *A. hydrophila*. As an opportunistic bacterium, *A. hydrophila* has cosmopolitan distribution in water bodies of the world. *A. hydrophila* causes the bacterial haemorrhagic septicaemia disease in many species of warm and cold water fish including rainbow trout (Constantino Casas et al., 1997; Harikrishnan & Balasundaram, 2005; Nya & Austin, 2010). The results of the present study may suggest a natural way to enhance the immunity and to control infection induced by *A. hydrophila* in rainbow trout.

## 2 | MATERIALS AND METHODS

### 2.1 | Fish and feeding trial

AQ7

Rainbow trout fish with an average weight of  $55 \pm 5.6$  g were purchased from a local cold water fish farm, and then transported using a tank with continuous oxygenation to lab and finally distributed into the 12 polyethylene tanks (containing 400 L disinfected and chlorinated water) for 7 days acclimation. During acclimation, the fish were fed three times a day at a rate of 4% of body weight with a commercial rainbow trout diet (Faradaneh Co., GFT1; diet form: pellet, pellet size:  $5 \pm 0.4$  mm; diet composition: total crude protein: 40%, total crude lipid: 14%, total crude ash: 9% and total crude fibre: 3%).

After the adaptation to lab conditions, fish were distributed into the 12 tanks at density of 30 fish per tank and fed DKE at dietary levels of 0 (control), 0.2, 0.25 and 0.3 mg/kg diet in three replicates. Feeding was carried out daily at a rate of 4% of fish body weight. The dietary concentrations of the essential oil in this study were selected based on the in vitro inhibitory effect (or disk diffusion assay) of the essential oil on the growth of *A. hydrophila* according to the guidelines of CLSI ( [Clinical & Laboratory Standards Institute](#)), formerly known as NCCLS (Wayne, 2007), with some modifications. Briefly, the sterile Petri dishes (9 cm in diameter) were prepared with 20 ml of Mueller–Hinton medium. Then, a 100 µl of the bacterial suspension ( $10^8$  CFU/ml) was spread on the plates. After 6 min, a sterile filter paper disc (6 mm) containing different concentrations of DKE (0, 0.05, 0.1, 0.15, 0.2, 0.25 and 0.3 mg) was placed on the surface of the plates at 4°C for 1 h followed by incubation at 37°C for 30 h. The diameters of the inhibition zones (mm) were measured, including the diameter of the disks. Gentamycin (30 µg/disk) was used as positive control. All tests were done in three replicates. [A disk diffusion assay was conducted to determine the highest antibacterial activity for DKE \(Table 1\). In this regard, ~~Based on the results of disk diffusion assay \(Table 1\),~~ the highest antibacterial activity was observed in treatments, 20, 25 and 30 ;](#) which these concentrations were selected for feeding trial. During feeding experiment, water was daily renewed by 90% and also the water quality indices (temperature:  $15.3 \pm 0.3^\circ\text{C}$  [using thermometer, ZEAL, UK], oxygen:  $6.8 \pm 0.11$  mg/L [using portable oxygen meter: Oxyguard Polaris Dissolved Oxygen Meter, Dynamic Aqua Supply Ltd, Canada], total ammonia:  $0.008 \pm 0.002$  mg/L and pH:  $7.5 \pm 0.13$  [using portable pH meter, Model AE-PH501]) were monitored daily.

**TABLE 1.** Sequences of oligonucleotide primers for real-time PCR

Genes	Primers	Primer sequences	ID number
Lysozyme	Lysozyme Forward	TCCCCAGGTATCCCATGAT	AB027305
	Lysozyme Reverse	GTGTCTGATGTGGCTGTGCT	
Complement	C3 Forward	AGCTTGCTGACTGGCTTTGT	100136951
	C3 Reverse	TCATAAACGGTGACCCCAAC	
TNF- $\alpha$	TNF- $\alpha$ Forward	CATCGGCCCTGCACCTATA	X99303
	TNF- $\alpha$ Reverse	TGTGGAACACAGCAGGGTTGG	
IL-8	IL-8 Forward	AGAATGTCAGCCAGCCTTGT	AJ279069
	IL-8 Reverse	TCTCAGACTCATCCCCTCAGT	
$\beta$ -Actin	$\beta$ -Actin Forward	ACTGCACAGCCAAGAGAGTTCA	AB039726
	$\beta$ -Actin Reverse	GTTATTAAAGCGGCCGATATGC	

T<sub>AQ11</sub> efficacy for target and reference **gensgenes** was 1.93–1.98.

## 2.2 | GC-Mass spectrometry of *Dracocephalum kotschy* essential oil

The gas chromatography–mass spectrometry (Shimadzu QP2010 Ultra series instrument, flow rate 0.7 ml/min, temperature rate: 3°C/min and final temperature: 330°C) was used to analyse the chemical composition of DKE.

## 2.3 | Growth and survival parameters

The growth and survival parameters were measured after feeding trial by the following equations (Ghafariarsani, Hoseinifar, Javahery, et al., 2022):

$$\text{Weight gain (WG) (\%)} = \frac{\text{final weight} - \text{initial weight}}{\text{initial weight}} \times 100$$

$$\text{Specific growth rate (SGR) (\%/d)} = \left( \frac{\ln \text{ final wt(g)} - \ln \text{ initial wt(g)}}{\text{days}} \right) \times 100$$

$$\text{Feed conversion rate (FCR)} = \frac{\text{total feed consumed(g)}}{\text{weight gain(g)}}$$

$$\text{Survival rate (SR) (\%)} = \left( \frac{\text{final fish numbers}}{\text{initial fish numbers}} \right) \times 100$$

## 2.4 | Sampling

The blood, mucus and tissue samples were taken from 10 fish per tank after the feeding trial. For this purpose, fish were starved for 24 h, anesthetized with 20 mg/L clove oil and then few ( $n = 3$ ) were placed in a nylon bag for collection of mucus based on the method of Subramanian et al. (2007). The blood samples were collected from remainders ( $n = 7$ ) from caudal vein using 2-ml syringe. The plasmatic part of blood was separated by centrifuging at 4°C for 5 min. In addition, a part of blood was allocated for measurement of the haematological indices.

For tissue sampling, the fish were first sacrificed with high dose of ethyl 3-aminobenzoate methanesulphonate (100 g/L) to take liver sample. The plasma and liver samples were finally stored in liquid nitrogen until biochemical analysis.

## 2.5 | Biochemical assays

### 2.5.1 | Immune parameters

The activity of lysozyme in plasma and mucus was assayed using lyophilized *Micrococcus luteus* according to Ellis (1990). The immunoglobulin (IgM) content of plasma and mucus was determined using polyethylene glycol according to Siwicki and Anderson (1993). The plasma alternative complement activity (ACH<sub>50</sub>) was assayed using the method of Ortuno et al. (2001) using *haemolysis* of rabbit red blood cells. The plasma bactericidal activity was evaluated by mixing plasma samples with bacterial suspension (*A. hydrophila*, OD: 0.5 at 546 nm) and finally calculating bacterial colony forming units (CFUs) created on nutrient agar plates after 24 h incubation at 36°C (Rao et al., 2006).

The total protein and albumin levels in plasma were spectrophotometrically determined using commercial assay kits (Sigma-Aldrich Co., USA) according to manufacturers' instructions. Protease activity in mucus was determined using the azocasein hydrolysis method, as described by Ross et al. (2000). Mucosal alkaline phosphatase (ALP) activity was measured using a commercial assay kit (Sigma-Aldrich Co., USA) according to manufacturers' instructions.

### 2.5.2 | Liver antioxidant enzymes

All antioxidant enzymes were determined using commercial assay kits, as instructed by manufacturer (*Sigma-Aldrich CO, USA*). Catalase (CAT) activity was assayed at 240 nm using catalysis of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) according to Claiborne (1985).

Superoxide dismutase (SOD) activity was colourimetrically determined at 440 nm through oxidation of xanthine to superoxide radicals (SR) and generation of a red formazan dye upon reaction of WST-1 reagent with the SR (Marklund & Marklund, 1974). Glutathione peroxidase (GPX) activity was measured through oxidation of glutathione by GPx and assaying oxidized glutathione at 340 nm. The lipid peroxidation was spectrophotometrically (at 532 nm) assayed upon reaction of malondialdehyde (MDA) with thiobarbituric acid (Utley et al., 1967).

### 2.5.3 | Haematological parameters

The total count of red and white blood cells was measured using a Neubauer haemocytometer slide at a magnification of 400× (Harikrishnan et al., 2009). Haematocrit (Hct) was measured using the microhaematocrit method (Blaxhall & Daisley, 1973). The haemoglobin (Hb) content was determined using the cyanohaemoglobin method (Oriakpono et al., 2012). The haematological indices including mean corpuscular haemoglobin (MCH), mean corpuscular volume (MCV) and corpuscular haemoglobin concentration (MCHC) were estimated according to the following equations (Blaxhall & Daisley, 1973):

$$\text{MCHC} = \text{Hb} \times \frac{10}{\text{Hct}}; \text{MCV} = \text{Hct} \times \frac{10}{\text{RBC(million)}}; \text{MCH} = \text{Hb} \times \frac{10}{\text{RBC(million)}}$$

## 2.6 | Gene expression assay

### 2.6.1 | RNA extraction

AQ15 The total RNA content was extracted from liver tissues using acid guanidinium thiocyanate–phenol–chloroform based on the method described by Awad et al. (2013) with some modifications. The quantity and quality of the extracted RNA (ER) was estimated by determining the absorbance at 260 nm, whereas the quality of ER was evaluated using electrophoresis according to Kwasek et al. (2014).

### 2.6.2 | RT-PCR

One microgram of ER was allocated to produce first-strand cDNAs using a Fermentas cDNA synthesis kit for reverse transcription polymerase chain reaction (RT-PCR) according to the manufacturer's protocol and a set of oligo-dT as primer. The real-time PCR primers were synthesized based on the sequences of DNA from GenBank (accessions numbers listed in Table 1) and using Gene Runner (version 6) software. The  $\beta$ -actin gene was considered as reference gene to evaluate the gene expressions.



AQ16 The real-time PCR was done using an iCycler (BioRad, USA) and commercial kit (Bio flux-Bioer Technology Co., China) following the manufacturer's protocols. The fold change in TNF- $\alpha$ , IL-8, complement and lysozyme relative to mRNA expression was  
AQ17 estimated using the  $2^{-\Delta\Delta C_t}$  method (Livak & Schmittgen, 2001) and the data were analysed using iQ5 optical system software version 2.1 (Bio-Rad).

## 2.7 | Bacterial challenge

After 60 days of feeding experiment, the remaining fish in the tanks were exposed to *A. hydrophila* bacterium by injection. The stock of the bacterium (Strain number: IBRC-M 10814) was cultured on agar medium for 24 h at 37°C. A sub-lethal dose of bacteria ( $1.3 \times 10^7$  cells/ml phosphate buffered saline [PBS]) was used for bacterial challenge, which was calculated through determining the 7-day LD<sub>50</sub> ( $3 \times 10^7$  cells/ml) in a previous exposure experiment by LaPatra et al. (2010). The cumulative fish mortality was recorded daily for 10 days. In addition, *A. hydrophila*-induced mortality was confirmed through re-isolating the organism from the liver on at least 30% of the  
AQ18 dead fish using conventional methods.

## 2.8 | Statistical analysis

All statistical analysis was done using SPSS software. The normality of data (mean  $\pm$  standard deviation) was evaluated using Kolmogorov–Smirnov test. The *Arcsin* transformation was used on proportional and per cent data before running analysis of variance. One-way analysis of variance was applied to find statistical differences. Finally, the comparison of means was done using Tukey test ( $p < 0.01$ ).

# 3 | RESULTS

## 3.1 | GC-Mass spectrometry of *Dracocephalum kotschyi* essential oil

The chemical composition of DKE is presented in Table 2.  $\delta$ -3-Carene (10.5%), limonene (9.8%), carvacrol (9.2%), 1,8-cineole (7.3%) and p-cymene (4.2%) composed the main compounds in the chemical composition of DKE (Table 2).

**TABLE 2.** Chemical compositions of *Dracocephalum kotschyi* essential oil; the compounds are listed in order of elution time

Compounds	Kovats constant	%
3Z-Hexenol	871	0.3
$\alpha$ -Thujene	935	0.2
$\alpha$ -Pinene	946	2.8
Sabinene	980	2.1
$\beta$ -Pinene	985	2.7
3-Octanol	1000	1.4
$\delta$ -3-Carene	1021	10.5
p-Cymene	1026	4.2
Limonene	1035	9.8
1,8-Cineole	1041	7.3
$\gamma$ -Terpinene	1065	3.1
Terpinolene	1101	3.2

## 3.2 | Growth and mortality parameters

The growth parameters (weight gain per cent, specific growth rate and food conversion ratio) and mortality rate showed no significant differences between the treatments throughout 10 weeks of feeding trial (Table 3,  $p > 0.01$ ).

AQ19

**TABLE 3.** Growth and survival of the fish (rainbow trout) after 70 days of supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschyi* essential oil (DKE)

	Treatments			
	0 mg DKE (control)	0.2 mg DKE	0.25 mg DKE	0.3 mg DKE
FW (g)	252.5 ± 8.5	260.2 ± 11.2	255.8 ± 9.4	264.1 ± 12.3
WG (%)	358.1 ± 9.5	372.5.5 ± 11.2	365.4 ± 8.7	381.8 ± 10.9
SGR (%/d)	2.16 ± 0.05	2.21 ± 0.04	2.19 ± 0.03	2.23 ± 0.06
FCR	1.2 ± 0.04	1.18 ± 0.07	1.16 ± 0.05	1.21 ± 0.04
SR (%)	86.5 ± 2.5 <sup>a</sup>	90.4 ± 2.3 <sup>a</sup>	92.2 ± 1.5 <sup>a</sup>	70.2 ± 3.6 <sup>b</sup>

### Note

Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

Abbreviations: FCR, Feed conversion rate; FW, Final weight; SGR, Specific growth rate (%/d); SR, Survival rate; WG, Weight gain.

## 3.3 | Plasma and mucus immune components

The activity of plasma complement and lysozyme and the levels of IgM, total protein and total albumin significantly increased in response to 0.2 and 0.25 mg DKE/kg diet compared with control and those fed 0.3 mg DKE/kg diet (Table 4,  $p < 0.01$ ).

However, the treatment with 0.3 mg DKE/kg diet showed higher activity in plasma complement, lysozyme and total protein compared with control (Table 4,  $p < 0.01$ ).

The DKE supplementation increased the bactericidal activity, as the number of CFUs decreased in DKE-supplemented fish (Table 4,  $p < 0.01$ ). In mucus, the protease activity, IgM content and lysozyme activity significantly increased in response to 0.2 and 0.25 mg DKE/kg diet compared with control and fish of 0.3 mg DKE/kg diet treatment (Table 4,  $p < 0.01$ ). However, the mucosal lysozyme activity and IgM content significantly decreased in fish fed 0.3 mg DKE/kg diet compared with other experimental groups (Table 4,  $p < 0.01$ ).

**TABLE 4.** The plasma and mucus immune component changes in rainbow trout in response to 70 days supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschy* essential oil (DKE)

	Treatments		
	0 mg DKE (control)	0.2 mg DKE	0.25 mg DKE
Plasma immune components			
Lysozyme activity (IU/ml)	60.6 ± 12.1 <sup>a</sup>	180.8 ± 15.3 <sup>b</sup>	241.5 ± 20.5 <sup>c</sup>
ACH <sub>50</sub> activity (IU/ml)	1.6 ± 0.3 <sup>a</sup>	4.9 ± 0.5 <sup>b</sup>	5.2 ± 0.3 <sup>b</sup>
IgM (mg/ml)	2.33 ± 0.3 <sup>a</sup>	3.84 ± 0.7 <sup>b</sup>	4.46 ± 0.4 <sup>b</sup>
Bactericidal activity (No. of. CFUs)	235.5 ± 22.51 <sup>a</sup>	152.4 ± 19.4 <sup>b</sup>	131.5 ± 12.5 <sup>b</sup>
Total protein (g/dL)	2.2 ± 0.21 <sup>a</sup>	4.3 ± 0.12 <sup>b</sup>	4.5 ± 0.15 <sup>b</sup>
Total albumin (g/dL)	0.73 ± 0.25 <sup>a</sup>	1.42 ± 0.14 <sup>b</sup>	2.65 ± 0.21 <sup>c</sup>
Mucosal immune components			
Lysozyme activity (IU/ml)	21.5 ± 6.3 <sup>a</sup>	42.3 ± 10.2 <sup>b</sup>	50.8 ± 12.7 <sup>b</sup>
IgM (mg/ml)	0.42 ± 0.12 <sup>a</sup>	0.95 ± 0.1 <sup>b</sup>	1.2 ± 0.15 <sup>b</sup>

### Note

Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

## 3.4 | Haematological parameters

After feeding trial, the values of haematocrit and red blood cell count significantly increased in fish fed with 0.2 and 0.25 mg DKE/kg diet (Table 5,  $p < 0.01$ ). Meanwhile, the mean erythrocyte cell volume (MCV), haemoglobin content and mean erythrocyte cell haemoglobin content (MCHC) significantly increased in fish fed 0.25 mg DKE/kg diet than those of fish fed the other diets (Table 5,  $p < 0.01$ ). The mean corpuscular haemoglobin (MCH) showed significant decreases in fish fed 0.2 and 0.25 mg DKE/kg diet than compared with those fed the other diets (Table 5,  $p < 0.01$ ). The values of white blood cell (WBC) significantly increased in response to 0.25 mg DKE/kg diet compared with other diets (Table 5,  $p < 0.01$ ). Also, there were no significant differences in WBC count between control and fish fed 0.3 mg DKE/kg diet (Table 5,  $p > 0.01$ ).

**TABLE 5.** The haematological alternations of rainbow trout after 70 days of supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschy* essential oil (DKE)

	Treatments			
	0 mg DKE (control)	0.2mg DKE	0.25mg DKE	0.3mg DKE
RBC ( $\times 10^6$ /ml)	0.86 $\pm$ 0.07 <sup>a</sup>	1.43 $\pm$ 0.08 <sup>b</sup>	1.58 $\pm$ 0.06 <sup>c</sup>	0.9 $\pm$ 0.04 <sup>a</sup>
WBC ( $\times 10^3$ /ml)	3.34 $\pm$ 0.25 <sup>a</sup>	3.62 $\pm$ 0.15 <sup>a</sup>	5.12 $\pm$ 0.2 <sup>b</sup>	3.7 $\pm$ 0.3 <sup>a</sup>
Hct (%)	26.4 $\pm$ 2.5 <sup>a</sup>	30.2 $\pm$ 1.42 <sup>b</sup>	31.2 $\pm$ 1.4 <sup>b</sup>	28.5 $\pm$ 2.1 <sup>a</sup>
Hb (g/dl)	8.92 $\pm$ 1.81 <sup>a</sup>	9.34 $\pm$ 2.52 <sup>a</sup>	13.61 $\pm$ 1.13 <sup>b</sup>	10.12 $\pm$ 1.21 <sup>a</sup>
MCV (fl)	212.4 $\pm$ 13.5 <sup>a</sup>	221.2 $\pm$ 10.1 <sup>a</sup>	268.3 $\pm$ 12.4 <sup>b</sup>	230.5 $\pm$ 18.1 <sup>a</sup>
MCH (pg)	104.6 $\pm$ 6.2 <sup>a</sup>	65.7 $\pm$ 7.1 <sup>b</sup>	86.1 $\pm$ 5.2 <sup>c</sup>	112.5 $\pm$ 8.8 <sup>a</sup>
MCHC (%)	3.1 $\pm$ 0.06 <sup>a</sup>	3.12 $\pm$ 0.03 <sup>a</sup>	3.24 $\pm$ 0.05 <sup>b</sup>	3.06 $\pm$ 0.11 <sup>a</sup>

### Note

Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

## 3.5 | Liver antioxidant enzymes

The activity of glutathione peroxidase (GPX), catalase (CAT) and superoxide dismutase (SOD) significantly increased in response to 0.25 and 0.3 mg DKE/kg diet compared with control and those fed diet containing 0.2 mg DKE/kg diet (Table 6,  $p < 0.01$ ). In the 0.2 mg DKE/kg diet treatment, only SOD showed significant increases compared with control (Table 6,  $p < 0.01$ ).

**TABLE 6.** The liver antioxidant enzyme changes in the rainbow trout in response to 70 days supplementation with diet containing different levels (mg/kg diet) of *Dracocephalum kotschy* essential oil (DKE)


	Treatments			
	0 mg DKE (control)	0.2 mg DKE	0.25 mg DKE	0.3 mg DKE
SOD (IU/mg)	0.411 $\pm$ 0.08 <sup>a</sup>	0.693 $\pm$ 0.06 <sup>b</sup>	0.752 $\pm$ 0.12 <sup>bc</sup>	0.981 $\pm$ 0.09 <sup>cd</sup>
CAT (IU/mg)	71.23 $\pm$ 13.3 <sup>a</sup>	58.33 $\pm$ 16.45 <sup>a</sup>	112.54 $\pm$ 8.5 <sup>b</sup>	132.5 $\pm$ 11.2 <sup>c</sup>
GPx (IU/mg)	0.121 $\pm$ 0.08 <sup>a</sup>	0.163 $\pm$ 0.1 <sup>a</sup>	0.462 $\pm$ 0.16 <sup>b</sup>	0.714 $\pm$ 0.11 <sup>c</sup>

### Note

Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ ).

### 3.6 | Lipid peroxidation index

As a lipid peroxidation marker, the levels of malondialdehyde (MDA) showed no significant differences between all experimental groups after 60 days feeding trial (Figure 1,  $p > 0.01$ ).




**FIGURE 1.** The changes in lipid peroxidation index (MDA: malondialdehyde levels) in the rainbow trout in response to 70 days of supplementation with diet containing different levels of *Dracocephalum kotschyi* essential oil (DKE). Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ )


### 3.7 | Gene expressions

After feeding trial, the expression of inflammatory genes, TNF- $\alpha$  and IL-8, significantly decreased in response to 0.2 and 0.25 mg DKE/kg diet (Figure 2,  $p < 0.01$ ). In contrast, these treatments exhibited higher expression of the immune-related genes (C3 and lysozyme) in response to 0.2 and 0.25 mg dietary DKE compared with control and fish from 0.3 mg DKE/kg diet treatment (Figure 3,  $p < 0.01$ ). The expression of C3 and lysozyme showed significant decreases in fish fed 0.3 mg DKE/kg diet compared with control (Figure 3,  $p < 0.01$ ).






**FIGURE 2.** The expression of cytokine-related genes, TNF- $\alpha$  and IL-8, in the rainbow trout in response to 70 days of supplementation with diet containing different levels of *Dracocephalum kotschy* essential oil (DKE). Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ )



**FIGURE 3.** The expression of immune-related genes, complement (C3) and lysozyme in the rainbow trout in response to 70 days of supplementation with diet containing different levels of *Dracocephalum kotschy* essential oil (DKE). Significant differences are shown as different superscripted letters (Tukey test,  $p < 0.01$ )

### 3.8 | Bacterial challenge

After 10 days of bacterial injection, the cumulative fish mortality (CFM) showed different values among the treatments (Figure 4,  $p < 0.01$ ). The CFM was significantly lower in fish fed DKE compared with control after 10 days bacterial challenge (Figure 4,  $p < 0.01$ ).



**FIGURE 4.** Cumulative mortality of rainbow trout, *Oncorhynchus mykiss* ( $n = 15/\text{tank}$ ), fed different concentration of DKE in the diet throughout 10-day challenge with *Aeromonas hydrophila*. Significant differences are indicated with different lower-case letters (Tukey test,  $p < 0.01$ ). Data: Mean  $\pm$  SD

## 4 | DISCUSSION

The findings of this study revealed that dietary DKE, especially at concentrations of 0.2 and 0.25 mg DKE/kg diet, can improve the immune and antioxidant system in the rainbow trout. Therefore, DKE can be considered as an immunostimulant, although this role has not yet been reported in fish. The antioxidant properties of *D. kotschy*

are mostly attributed to the flavonoids such as terpinolene, p-Cymene,  $\gamma$ -terpinene, carvone, carvacrol, eugenol and limonene (Ahmad & Beg, 2013; Aydin et al., 2013; Foti & Ingold, 2003; Gülçin, 2011; de Oliveira et al., 2015; Pombal et al., 2017; Yanishlieva et al., 1999).

It is recognized that the compound luteolin plays the main role in the antioxidant activity of the plant (Fattahi et al., 2013; Gohari et al., 2003; Heydari et al., 2019). In this study, the dietary DKE also stimulated the production of antioxidant enzymes (SOD, CAT and GPx) in liver, which ~~this result~~ clearly indicates the enhancing effects of DKE on the enzymatic antioxidant defence defense in the fish. However, we found no changes in the MDA levels as the main indicator of oxidative stress (Del Rio et al., 2005) in response to dietary DKE. In this study, DKE at dietary levels of 0.2 and 0.25 mg DKE/kg elevated immune components in plasma (ACH<sub>50</sub>, IgM content, lysozyme activity, total protein and total albumin) and mucus (protease activity, IgM content and lysozyme activity) and also the expression of some immune-related genes (lysozyme and C3). However, we found no report regarding the immunogenic effects of *D. kotschyi* in the literatures. Some studies have reported a significant decline in secretion of IL-1 $\beta$  from the cells, which may be resulting from the presence of anti-inflammatory compounds in the biochemical composition of *D. kotschyi* (Amirghofran et al., 2011).

The results of the present study support this assumption because we observed down-regulation of the inflammatory genes, TNF- $\alpha$  and IL-8, after supplementation with 0.2 and 0.25 mg DKE/kg diet. The anti-inflammatory effects of *Dracocephalum* sp. has been reported in some studies (Kalantar et al., 2018; Nie et al., 2021; Sadraei et al., 2017; Toshmatov et al., 2019). Some studies have reported the antimicrobial activity of *D. kotschyi* essential oil, which may be related to some compounds such as geranial  $\alpha$ -pinene, geraniol, neral, geraniol acetate and limonene in the biochemical composition of *D. kotschyi* (Dorman & Deans, 2000; Kamali et al., 2015; Singh et al., 2012).

In general, haematology is an important tool to evaluate the fish health in relation to diseases, stress, nutrition and changes in environmental conditions (Burgos-Aceves et al., 2019; Enayat Gholampour et al., 2020; Fazio, 2019; Ghafarifarsani, Hoseinifar, Aftabgard, et al., 2022; Rashidian, Kajbaf, et al., 2020; Sula et al., 2020). In this study, dietary DKE caused significant increases in the values of Hct, RBCs, MCV, Hb content and MCHC. However, MCH decreased in fish fed 0.2 and 0.25mg DKE/kg diet, which may be due to the depressing effect of DKE on the formation of Hb in red blood cells. In addition, the values of WBCs significantly increased in response to 0.25mg DKE/kg diet, which may be attributed to the immunostimulatory role of DKE. Generally, the effects of plants and their derivatives on fish haematology are different, which may be

dependent on plant species, exposure concentration, exposure duration, experimental condition, etc.

In the present study, the supplementation of fish with 0.3mg DKE/kg diet reduced the innate immune responses compared with the 0.2 and 0.25mg DKE/kg diet treatments. Furthermore, the expression of lysozyme and C3 genes in the 0.3mg DKE/kg diet treatment was lower than control. These results may be due to the immunotoxic effects of DKE at high dietary concentrations. The cytotoxic effects of *D. kotschyi* have been previously reported by Ashrafi et al. (2017), Sani et al. (2017) and Shaabani et al. (2020). Ashrafi et al. (2017) indicated that DKE at concentrations of 26.4 µg/ml and 4266.7 µg/ml has toxic effects on Hela cells and human lymphocytes over in vitro condition. Similar results were observed in the study of Sani et al. (2017), when calu-6 and mehr-80 lung cancer cell lines were exposed to 12.5 200 µg/mL of methanolic extract of *D. kotschyi*. In addition, Shaabani et al. (2020) reported the cytotoxic effects of 400 µg/ml *D. kotschyi* extract on glioblastoma U87 cells over in vitro condition.

In the present study, dietary DKE reduced the mortality induced by *A. hydrophila* infection. However, this reduction was not so considerable. Generally, it is found that the antibacterial effect of DKE is stronger on Gram-positive bacteria (Ashrafi et al., 2017), as it is reported for other plant essential oils (Ballester-Costa et al., 2013; Shakeri et al., 2014). Due to the lack of cell wall in Gram-positive bacteria, the phospholipid bilayer of these bacteria is more receptive to hydrophobic compounds of essential oils than Gram-negative bacteria (Nikaido, 2003). The contact of essential oils with cell membrane results in disruptions of permeability and bacterial enzyme systems (Wendakoon & Sakaguchi, 1993). Therefore, in this study, the low antibacterial effect of DKE may be due to the Gram-negative bacteria nature of *A. hydrophila*. However, the antibacterial properties of DKE may be due to some compounds including terpinolene, β-Pinene, p-cymene, carvone, carvacrol, eugenol and limonene in its chemical composition (Devi et al., 2010; Han et al., 2020; Marchese et al., 2017; Porfírio et al., 2017; Rondon et al., 2006; Salehi et al., 2019; Xu et al., 2008).

## 5 | CONCLUSION

The finding of the present study indicated that DKE at dietary may improve the immunity and antioxidant defence in the rainbow trout. However, DKE had no appropriate efficiency in reducing the fish mortality induced by *A. hydrophila*.

## CONFLICT OF INTEREST

## ETHICAL APPROVAL

All applicable international, national and/or institutional guidelines for the care and use of animals were followed.

### Open Research

#### DATA AVAILABILITY STATEMENT

The datasets in this study are available from the corresponding author on reasonable request. All data and materials are available for publication.

## AQ26 REFERENCES

- ✓ Ahmad, S., & Beg, Z. H. (2013). Hypolipidemic and antioxidant activities of thymoquinone and limonene in atherogenic suspension fed rats. *Food Chemistry*, 138(2–3), 1116–1124. <https://doi.org/10.1016/j.foodchem.2012.11.109>
- ✓ Ahmadifar, E., Sheikhzadeh, N., Roshanaei, K., Dargahi, N., & Faggio, C. (2019). Can dietary ginger (*Zingiber officinale*) alter biochemical and immunological parameters and gene expression related to growth, immunity and antioxidant system in zebrafish (*Danio rerio*)? *Aquaculture*, 507, 341–348. <https://doi.org/10.1016/j.aquaculture.2019.04.049>
- ✓ Amirghofran, Z., Malek-Hosseini, S., Golmoghaddam, H., Kalantar, F., & Shabani, M. (2011). Inhibition of nitric oxide production and proinflammatory cytokines by several medicinal plants. *Iranian Journal of Immunology*, 8(3), 159–169.
- ❓ Asghari, G., Nasr Esfahani, B., & Paydar, P. (2015). Evaluating the effect of *Dracocephalum kotschy* methanol extract on *Mycobacterium tuberculosis*. *Research Journal of Pharmacognosy*, 2(3), 31–36.
- ✓ Ashrafi, B., Ramak, P., Ezatpour, B., & Talei, G. R. (2017). Investigation on chemical composition, antimicrobial, antioxidant, and cytotoxic properties of essential oil from *Dracocephalum kotschy* Boiss. *African Journal of Traditional, Complementary and Alternative Medicines*, 14(3), 209–217.

<https://doi.org/10.21010/ajtcam.v14i3.23>

- ❓ Awad, E., Austin, D., & Lyndon, A. R. (2013). Effect of black cumin seed oil (*Nigella sativa*) and nettle extract (Quercetin) on enhancement of immunity in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquaculture*, 388, 193–197.  
<https://doi.org/10.1016/j.aquaculture.2013.01.008>
- ✓ Aydin, E., Türkez, H., & Taşdemir, Ş. (2013). Anticancer and antioxidant properties of terpinolene in rat brain cells. *Archives of Industrial Hygiene and Toxicology*, 64(3), 415–424. <https://doi.org/10.2478/10004-1254-64-2013-2365>
- ✓ Ballester-Costa, C., Sendra, E., Fernández-López, J., Pérez-Álvarez, J. A., & Viuda-Martos, M. (2013). Chemical composition and in vitro antibacterial properties of essential oils of four *Thymus* species from organic growth. *Industrial Crops and Products*, 50, 304–311. <https://doi.org/10.1016/j.indcrop.2013.07.052>
- ❓ Bibi, M., Khan, S., Kakar, A.-U.-R., Khan, N., Tareen, A. H., Bibi, S., & Taj, H. (2021). Determination of essential and non-essential elements in *Xylanthemum macropodum* of Balochistan, Pakistan. *Baghdad Journal of Biochemistry and Applied Biological Sciences*, 2(02), 94–103. <https://doi.org/10.47419/bjbabs.v2i02.39>
- ✓ Blaxhall, P. C., & Daisley, K. W. (1973). Routine haematological methods for use with fish blood. *Journal of Fish Biology*, 5(6), 771–781. <https://doi.org/10.1111/j.1095-8649.1973.tb04510.x>
- ✓ Brown, R., Moore, L., Mani, A., Patel, S., & Salinas, I. (2021). Effects of ploidy and salmonid alphavirus infection on the skin and gill microbiome of Atlantic salmon (*Salmo salar*). *PLoS One*, 16(2), e0243684.  
<https://doi.org/10.1371/journal.pone.0243684>
- ✓ Burgos-Aceves, M. A., Lionetti, L., & Faggio, C. (2019). Multidisciplinary haematology as prognostic device in environmental and xenobiotic stress-induced response in fish. *Science of the Total Environment*, 670, 1170–1183.  
<https://doi.org/10.1016/j.scitotenv.2019.03.275>
- ✓ Cao, L., Wang, W., Yang, Y., Yang, C., Yuan, Z., Xiong, S., & Diana, J. (2007). Environmental impact of aquaculture and countermeasures to aquaculture pollution in China. *Environmental Science and Pollution Research-International*, 14(7), 452–462.  
<https://doi.org/10.1065/espr2007.05.426>
- ❓ Claiborne, A. (1985). Catalase activity. In R. A. Greenwald (Ed.). *CRC handbook of*



*methods in oxygen radical research* (pp. 283–284). CRC Press.

- ✓ Cole, D. W., Cole, R., Gaydos, S. J., Gray, J., Hyland, G., Jacques, M. L., Powell-Dunford, N., Sawhney, C., & Au, W. W. (2009). Aquaculture: Environmental, toxicological, and health issues. *International Journal of Hygiene and Environmental Health*, 212(4), 369–377. <https://doi.org/10.1016/j.ijheh.2008.08.003>
- ✓ Constantino Casas, F., Armijo Ortiz, A., Osorio Sarabia, D., & Chavez Soriano, L. A. (1997). Infection due to *Aeromonas hydrophila* and *Ichthyophthirius multifiliis* in rainbow trout (*Oncorhynchus mykiss*, Walbaum) and tilapia (*Oreochromis aureus*, L) in Morelos, Mexico. Pathological study and review of treatment. *Veterinaria Mexico*, 28(1), 59–62.
- ✓ de Oliveira, T. M., de Carvalho, R. B. F., da Costa, I. H. F., de Oliveira, G. A. L., de Souza, A. A., de Lima, S. G., & de Freitas, R. M. (2015). Evaluation of p-cymene, a natural antioxidant. *Pharmaceutical Biology*, 53(3), 423–428.
- ✓ Deivi Arunachalam, K., Kuruva, J. K., Pradhoshini, K. P., Musthafa, M. S., & Faggio, C. (2021). Antioxidant and antigenotoxic potential of *Morinda tinctoria* Roxb. leaf extract succeeding cadmium exposure in Asian catfish, *Pangasius sutchi*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 249, 109149.
- ✓ Del Rio, D., Stewart, A. J., & Pellegrini, N. (2005). A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutrition, Metabolism and Cardiovascular Diseases*, 15(4), 316–328. <https://doi.org/10.1016/j.numecd.2005.05.003>
- ✓ Devi, K. P., Nisha, S. A., Sakthivel, R., & Pandian, S. K. (2010). Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhi* by disrupting the cellular membrane. *Journal of Ethnopharmacology*, 130(1), 107–115. <https://doi.org/10.1016/j.jep.2010.04.025>
- ✓ Dorman, H. D., & Deans, S. G. (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, 88(2), 308–316. <https://doi.org/10.1046/j.1365-2672.2000.00969.x>
- ❓ Ellis, A. I. (1990). Lysozyme assays. *Techniques in Fish Immunology*, 1, 101–103.
- ✓ Enayat Gholampour, T., Fadaei Raieni, R., Pouladi, M., Larijani, M., Pagano, M., & Faggio, C. (2020). The dietary effect of *Vitex agnus-castus* hydroalcoholic extract

on growth performance, blood biochemical parameters, carcass quality, sex ratio and gonad histology in Zebrafish (*Danio rerio*). *Applied Sciences*, 10(4), 1402.

<https://doi.org/10.3390/app10041402>

- ✓ Farsani, M. N., Hoseinifar, S. H., Rashidian, G., Farsani, H. G., Ashouri, G., & Van Doan, H. (2019). Dietary effects of *Coriandrum sativum* extract on growth performance, physiological and innate immune responses and resistance of rainbow trout (*Oncorhynchus mykiss*) against *Yersinia ruckeri*. *Fish & Shellfish Immunology*, 91, 233–240. <https://doi.org/10.1016/j.fsi.2019.05.031>
- ✓ Fattahi, M., Nazeri, V., Torras-Claveria, L., Sefidkon, F., Cusido, R. M., Zamani, Z., & Palazon, J. (2013). Identification and quantification of leaf surface flavonoids in wild-growing populations of *Dracocephalum kotschyi* by LC–DAD–ESI–MS. *Food Chemistry*, 141(1), 139–146. <https://doi.org/10.1016/j.foodchem.2013.03.019>
- ❓ Fattepur, S., Nilugal, K. C., Rajendran, R., Asmani, F., & Yusuf, E. (2018). Anti-hyperlipidemic activity of methanolic extract of *Boesenbergia pandurata* (finger root) in experimental induced hypercholesterolemic Sprague Dawley rats. *Asian Journal of Pharmaceutical and Clinical Research*, 11(3), 8–12. <https://doi.org/10.22159/ajpcr.2018.v11s3.29962>
- ✓ Fazio, F. (2019). Fish hematology analysis as an important tool of aquaculture: A review. *Aquaculture*, 500, 237–242. <https://doi.org/10.1016/j.aquaculture.2018.10.030>
- ✓ Foti, M. C., & Ingold, K. U. (2003). Mechanism of inhibition of lipid peroxidation by  $\gamma$ -terpinene, an unusual and potentially useful hydrocarbon antioxidant. *Journal of Agricultural and Food Chemistry*, 51(9), 2758–2765. <https://doi.org/10.1021/jf020993f>
- ✓ Ghafarifarsani, H., Hoseinifar, S. H., Adorian, T. J., Ferrigolo, F. R. G., Raissy, M., & Van Doan, H. (2021). The effects of combined inclusion of *Malvae sylvestris*, *Origanum vulgare*, and *Allium hirtifolium* Boiss for common carp (*Cyprinus carpio*) diet: Growth performance, antioxidant defense, and immunological parameters. *Fish & Shellfish Immunology*, 119, 670–677. <https://doi.org/10.1016/j.fsi.2021.10.014>
- ✓ Ghafarifarsani, H., Hoseinifar, S. H., Aftabgard, M., & Van Doan, H. (2022). The improving role of savory (*Satureja hortensis*) essential oil for Caspian roach (*Rutilus caspicus*) fry: Growth, haematological, immunological, and antioxidant parameters and resistance to salinity stress. *Aquaculture*, 548, 737653. <https://doi.org/10.1016/j.aquaculture.2021.737653>
- ✓ Ghafarifarsani, H., Hoseinifar, S. H., Javahery, S., Yazici, M., & Van Doan, H.

(2022). Growth performance, biochemical parameters, and digestive enzymes in common carp (*Cyprinus carpio*) fed experimental diets supplemented with vitamin C, thyme essential oil, and quercetin. *Italian Journal of Animal Science*, 21(1), 291–302.

- ✓ Ghafarifarsani, H., Hoseinifar, S. H., Talebi, M., Yousefi, M., Van Doan, H., Rufchaei, R., & Paolucci, M. (2021). Combined and singular effects of ethanolic extract of persian shallot (*Allium hirtifolium* Boiss) and synbiotic Biomin<sup>®</sup> IMBO on growth performance, serum-and mucus-immune parameters and antioxidant defense in Zebrafish (*Danio rerio*). *Animals*, 11(10), 2995.  
<https://doi.org/10.3390/ani11102995>
- ✓ Ghafarifarsani, H., Kachuei, R., & Imani, A. (2021). Dietary supplementation of garden thyme essential oil ameliorated the deteriorative effects of aflatoxin B<sub>1</sub> on growth performance and intestinal inflammatory status of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 531, 735928.  
<https://doi.org/10.1016/j.aquaculture.2020.735928>
- ❓ Ghafarifarsani, H., Rashidian, G., Sheikhlar, A., Naderi Farsani, M., Hoseinifar, S. H., & Van Doan, H. (2021). The use of dietary oak acorn extract to improve haematological parameters, mucosal and serum immunity, skin mucus bactericidal activity, and disease resistance in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Research*, 52(6), 2518–2527.
- ✓ Ghavam, M., Manconi, M., Manca, M. L., & Bacchetta, G. (2021). Extraction of essential oil from *Dracocephalum kotschy* Boiss. (Lamiaceae), identification of two active compounds and evaluation of the antimicrobial properties. *Journal of Ethnopharmacology*, 267, 113513.
- ❓ Gohari, A. R., Saeidnia, S., Matsuo, K., Uchiyama, N., Yagura, T., Ito, M., Kiuchi, F., & Honda, G. (2003). Flavonoid constituents of *Dracocephalum kotschy* growing in Iran and their trypanocidal activity. *Natural Medicines*, 57(6), 250–252.
- ❓ Gondo, H. K. (2021). The effect of spirulina on apoptosis through the caspase-3 pathway in a Preeclamptic Wistar rat model. *Journal of Natural Science, Biology and Medicine*, 12(3), 280–284.
- ✓ Gülçin, İ. (2011). Antioxidant activity of eugenol: A structure–activity relationship study. *Journal of Medicinal Food*, 14(9), 975–985.  
<https://doi.org/10.1089/jmf.2010.0197>

- ❓ Han, Y., Sun, Z., & Chen, W. (2020). Antimicrobial susceptibility and antibacterial mechanism of limonene against *Listeria monocytogenes*. *Molecules*, 25(1), 33.
- ❓ Harikrishnan, R., & Balasundaram, C. (2005). Modern trends in *Aeromonas hydrophila* disease management with fish. *Reviews in Fisheries Science*, 13(4), 281–320.
- ❓ Harikrishnan, R., Balasundaram, C., Dharaneedharan, S., Moon, Y. G., Kim, M. C., Kim, J. S., & Heo, M. S. (2009). Effect of plant active compounds on immune response and disease resistance in *Cirrhina mrigala* infected with fungal fish pathogen, *Aphanomyces Invadans*. *Aquaculture Research*, 40(10), 1170–1181.
- ✓ Heydari, P., Yavari, M., Adibi, P., Asghari, G., Ghanadian, S. M., Dida, G. O., & Khamesipour, F. (2019). Medicinal properties and active constituents of *Dracocephalum kotschy* and its significance in Iran: A systematic review. *Evidence-Based Complementary and Alternative Medicine*, 2019, 1–14.
- ✓ Jackson, P. B. (1978). Health and disease in intensive aquaculture. *Journal of the South African Veterinary Association*, 49(1), 57–59.
- ✓ Kalantar, K., Gholijani, N., Mousaei, N., Yazdani, M., & Amirghofran, Z. (2018). Investigation of *Dracocephalum kotschy* plant extract on the effective inflammatory transcription factors and mediators in activated macrophages. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Inflammatory and Anti-Allergy Agents)*, 17(1), 39–49.  
<https://doi.org/10.2174/1871523017666180608081656>
- ✓ Kamali, M., Khosroyar, S., & Mohammadi, A. (2015). Antibacterial activity of various extracts from *Dracocephalum kotschy* against food pathogenic microorganisms. *International Journal of PharmTech Research*, 8(9), 158–163.
- ✓ Kwasek, K., Terova, G., Lee, B. J., Bossi, E., Saroglia, M., & Dabrowski, K. (2014). Dietary methionine supplementation alters the expression of genes involved in methionine metabolism in salmonids. *Aquaculture*, 433, 223–228.  
<https://doi.org/10.1016/j.aquaculture.2014.05.031>
- ✓ LaPatra, S. E., Plant, K. P., Alcorn, S., Ostland, V., & Winton, J. (2010). An experimental vaccine against *Aeromonas hydrophila* can induce protection in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 33(2), 143–151.

- ✓ Liu, S., Zhao, H., Lehmler, H. J., Cai, X., & Chen, J. (2017). Antibiotic pollution in marine food webs in Laizhou Bay, North China: Trophodynamics and human exposure implication. *Environmental Science & Technology*, 51(4), 2392–2400. <https://doi.org/10.1021/acs.est.6b04556>
- ✓ Lulijwa, R., Rupia, E. J., & Alfaro, A. C. (2020). Antibiotic use in aquaculture, policies and regulation, health and environmental risks: A review of the top 15 major producers. *Reviews in Aquaculture*, 12(2), 640–663. <https://doi.org/10.1111/raq.12344>
- ✓ Marchese, A., Arciola, C. R., Barbieri, R., Silva, A. S., Nabavi, S. F., Tsetegho Sokeng, A. J., Izadi, M., Jafari, N. J., Suntar, I., Daglia, M., & Nabavi, S. M. (2017). Update on monoterpenes as antimicrobial agents: A particular focus on p-cymene. *Materials*, 10(8), 947. <https://doi.org/10.3390/ma10080947>
- ✓ Marklund, S., & Marklund, G. (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*, 47(3), 469–474. <https://doi.org/10.1111/j.1432-1033.1974.tb03714.x>
- ✓ Moradi, H., Ghavam, M., & Tavili, A. (2020). Study of antioxidant activity and some herbal compounds of *Dracocephalum kotschyi* Boiss. in different ages of growth. *Biotechnology Reports*, 25, e00408.
- ✓ Nie, L., Li, R., Huang, J., Wang, L., Ma, M., Huang, C., Wu, T., Yan, R., & Hu, X. (2021). Abietane diterpenoids from *Dracocephalum moldavica* L. and their anti-inflammatory activities in vitro. *Phytochemistry*, 184, 112680.
- ✓ Nikaido, H. (2003). Molecular basis of bacterial outer membrane permeability revisited. *Microbiology and Molecular Biology Reviews*, 67(4), 593–656. <https://doi.org/10.1128/MMBR.67.4.593-656.2003>
- ✓ Nya, E. J., & Austin, B. (2010). Use of bacterial lipopolysaccharide (LPS) as an immunostimulant for the control of *Aeromonas hydrophila* infections in rainbow trout *Oncorhynchus mykiss* (Walbaum). *Journal of Applied Microbiology*, 108(2), 686–694.
- ❓ Oriakpono, O., Aduabobo, H., & Wokoma, E. (2012). Biomarker response of a Cichlid Fish *Tilapia guineensis* to crude oil exposure. *International Journal of Modern Cellular and Molecular Biology*, 1(1), 10–18.
- ✓ Ortuno, J., Esteban, M. A., & Meseguer, J. (2001). Effects of short-term crowding

stress on the gilthead seabream (*Sparus aurata* L.) innate immune response. *Fish & Shellfish Immunology*, 11(2), 187–197. <https://doi.org/10.1006/fsim.2000.0304>

- ✓ Othman, Z., Khalep, H. R. H., Abidin, A. Z., Hassan, H., & Fattepur, S. (2019). The anti-angiogenic properties of *Morinda citrifolia*. L (Mengkudu) Leaves using chicken chorioallantoic membrane (CAM) assay. *Pharmacognosy Journal*, 11(1), 12–15.
- ✓ Páez-Osuna, F. (2001). The environmental impact of shrimp aquaculture: A global perspective. *Environmental Pollution*, 112(2), 229–231. [https://doi.org/10.1016/S0269-7491\(00\)00111-1](https://doi.org/10.1016/S0269-7491(00)00111-1)
- ✓ Pombal, S., Hernández, Y., Diez, D., Mondolis, E., Mero, A., Moran-Pinzon, J., Guerrero, E. I., & Rodilla, J. M. (2017). Antioxidant activity of carvone and derivatives against superoxide ion. *Natural Product Communications*, 12(5), 1934578X1701200502. <https://doi.org/10.1177/1934578X1701200502>
- ✓ Porfírio, E. M., Melo, H. M., Pereira, A. M. G., Cavalcante, T. T. A., Gomes, G. A., Carvalho, M. G. D., Costa, R. A., & Júnior, F. E. A. C. (2017). In vitro antibacterial and antibiofilm activity of *Lippia alba* essential oil, citral, and carvone against *Staphylococcus aureus*. *The Scientific World Journal*, 2017, 4962707.
- ✓ Pradeepkiran, J. A. (2019). Aquaculture role in global food security with nutritional value: A review. *Translational Animal Science*, 3(2), 903–910. <https://doi.org/10.1093/tas/txz012>
- ❓ Qahir, A., Kakar, A.-R., Khan, N., Khan, S., Hakeem, A., Kamal, R., & Fazal-Ur-Rehman. (2021). The antioxidant, antimicrobial, and clinical effects with elemental contents of pomegranate (*Punica granatum*) peel extracts: A review. *Baghdad Journal of Biochemistry and Applied Biological Sciences*, 2(01), 21–28. <https://doi.org/10.47419/bjbabs.v2i01.33>
- ✓ ~~Quade, M. J., & Roth, J. A. (1997). A rapid, direct assay to measure degranulation of bovine neutrophil primary granules. *Veterinary Immunology and Immunopathology*, 58(3–4), 239–248. [https://doi.org/10.1016/S0165-2427\(97\)00048-2](https://doi.org/10.1016/S0165-2427(97)00048-2)~~
- ❓ Raissy, M., Ghafarifarsani, H., Hoseinifar, S. H., El-Haroun, E. R., Naserabad, S. S., & Van Doan, H. (2022). The effect of dietary combined herbs extracts (oak acorn, coriander, and common mallow) on growth, digestive enzymes, antioxidant and immune response, and resistance against *Aeromonas hydrophila* infection in common carp, *Cyprinus Carpio*. *Aquaculture*, 546, 737287. <https://doi.org/10.1016/j.aquaculture.2021.737287>



- ✓ Rao, Y. V., Das, B. K., Jyotirmayee, P., & Chakrabarti, R. (2006). Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 20(3), 263–273. <https://doi.org/10.1016/j.fsi.2005.04.006>
- ✓ Rashidian, G., Bahrami Gorji, S., Farsani, M. N., Prokić, M. D., & Faggio, C. (2020). The oak (*Quercus brantii*) acorn as a growth promotor for rainbow trout (*Oncorhynchus mykiss*): growth performance, body composition, liver enzymes activity and blood biochemical parameters. *Natural Product Research*, 34(17), 2413–2423.
- ✓ Rashidian, G., Boldaji, J. T., Rainis, S., Prokić, M. D., & Faggio, C. (2021). Oregano (*Origanum vulgare*) extract enhances zebrafish (*Danio rerio*) growth performance, serum and mucus innate immune responses and resistance against *Aeromonas hydrophila* challenge. *Animals*, 11(2), 299. <https://doi.org/10.3390/ani11020299>
- ✓ Rashidian, G., Kajbaf, K., Prokić, M. D., & Faggio, C. (2020). Extract of common mallow (*Malvae sylvestris*) enhances growth, immunity, and resistance of rainbow trout (*Oncorhynchus mykiss*) fingerlings against *Yersinia ruckeri* infection. *Fish & Shellfish Immunology*, 96, 254–261. <https://doi.org/10.1016/j.fsi.2019.12.018>
- ❓ Romero, J., Feijoó, C. G., & Navarrete, P. (2012). Antibiotics in aquaculture—use, abuse and alternatives. In E. D. Carvalho, G. S. David, & R. J. Silva (Eds.), *Health and environment in aquaculture* (p. 159). IntechOpen.
- ❓ Rondon, M., Velasco, J., Hernandez, J., Pecheneda, M., Rojas, J., Morales, A., Carmona, J., & Diaz, T. (2006). Chemical composition and antibacterial activity of the essential oil of *Tagetes patula* L. (Asteraceae) collected from the *Venezuela Andes*. *Revista Latinoamericana De Química*, 34(1/3), 32.
- ✓ Roslan, N., & Shariff, H. (2021). Potential pharmacotherapy of *Olea europaea* (Olive) fruit oil against neurotoxicity induced methamphetamine. *Journal of Cellular and Molecular Anesthesia*, 4(2), 67–68.
- ✓ Ross, N. W., Firth, K. J., Wang, A., Burka, J. F., & Johnson, S. C. (2000). Changes in hydrolytic enzyme activities of naive Atlantic salmon *Salmo salar* skin mucus due to infection with the salmon louse *Lepeophtheirus salmonis* and cortisol implantation. *Diseases of Aquatic Organisms*, 41(1), 43–51.
- ✓ Sadraei, H., Asghari, G., Khanabadi, M., & Minaiyan, M. (2017). Anti-inflammatory effect of apigenin and hydroalcoholic extract of *Dracocephalum kotschyi* on acetic

acid-induced colitis in rats. *Research in Pharmaceutical Sciences*, 12(4), 322.

<https://doi.org/10.4103/1735-5362.212050>

- ✓ Salehi, B., Upadhyay, S., Erdogan Orhan, I., Kumar Jugran, A., L.D. Jayaweera, S., A. Dias, D., Sharopov, F., Taheri, Y., Martins, N., Baghalpour, N., C. Cho, W., & Sharifi-Rad, J. (2019). Therapeutic potential of  $\alpha$ - and  $\beta$ -pinene: A miracle gift of nature. *Biomolecules*, 9(11), 738. <https://doi.org/10.3390/biom9110738>
- ✓ Sani, T. A., Mohammadpour, E., Mohammadi, A., Memariani, T., Yazdi, M. V., Rezaee, R., Calina, D., Docea, A. O., Goumenou, M., Etemad, L., & Shahsavand, S. (2017). Cytotoxic and apoptogenic properties of *Dracocephalum kotschy* aerial part different fractions on calu-6 and mehr-80 lung cancer cell lines. *Farmacia*, 65(2), 189–199.
- ✓ Shaabani, M., Mousavi, S. H., Azizi, M., & Jafari, A. A. (2020). Cytotoxic and apoptogenic effects of *Dracocephalum kotschy* Boiss, extracts against human glioblastoma U87 cells. *Avicenna Journal of Phytomedicine*, 10(6), 594.
- ✓ Shakeri, A., Khakdan, F., Soheili, V., Sahebkar, A., Rassam, G., & Asili, J. (2014). Chemical composition, antibacterial activity, and cytotoxicity of essential oil from *Nepeta ucrainica* L. spp. kopetdaghensis. *Industrial Crops and Products*, 58, 315–321. <https://doi.org/10.1016/j.indcrop.2014.04.009>
- ❓ Shalaby, M. N. (2015). Effects of protein hydrolysates on physical performance and immunity in male soccer players. *The International Scientific Journal of Physical Education and Sport Sciences*, 2(2), 1–8.
- ✓ Shalaby, M. N. (2018). The effect of whey protein (Natural Nanoparticle) on muscle strength, GH, IGF, T. Protein and body composition. *International Journal of Pharmaceutical Research & Allied Sciences*, 7(1), 126–132.
- ❓ Shalaby, M., Sakoury, M. M. A., Harthi, S. M., Alshalawi, F. M., Alhajji, M. M., Alshaikh, Z. H., & Aljaber, A. H. (2020). Vitamin D3 for health and muscle functions of athletes. *Systematic Reviews in Pharmacy*, 11(9), 851–854.
- ❓ Shalaby, M. N., Sakoury, M. M., Kholif, M. A., & Alsayed, N. I. (2020). The role of Amino Acids in improving immunity and growth factors of Volleyball players. *Journal of Advanced Pharmacy Education and Research*, 10(4), 140–144.
- ✓ Singh, S. K., Vishnoi, R., Dhingra, G. K., & Kishor, K. (2012). Antibacterial activity of leaf extracts of some selected traditional medicinal plants of Uttarakhand, North

- ❓ Siwicki, A. K., & Anderson, D. P. (1993). Nonspecific defense mechanisms assay in fish. II. Potential killing activity of neutrophils and macrophages, lysozyme activity in serum and organs and total immunoglobulin level in serum. In A. K. Siwicki, D. P. Anderson, & J. Waluga (Eds.), *Fish disease diagnosis and prevention methods* (pp. 105–112). Wydawnictwo Instytutu Rybactwa Stroladowego.
- ✓ Subramanian, S., MacKinnon, S. L., & Ross, N. W. (2007). A comparative study on innate immune parameters in the epidermal mucus of various fish species. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 148(3), 256–263. <https://doi.org/10.1016/j.cbpb.2007.06.003>
- ✓ Sula, E., Aliko, V., Pagano, M., & Faggio, C. (2020). Digital light microscopy as a tool in toxicological evaluation of fish erythrocyte morphological abnormalities. *Microscopy Research and Technique*, 83(4), 362–369. <https://doi.org/10.1002/jemt.23422>
- ✓ Talari, M., Seydi, E., Salimi, A., Mohsenifar, Z., Kamalinejad, M., & Pourahmad, J. (2014). Dracocephalum: novel anticancer plant acting on liver cancer cell mitochondria. *BioMed Research International*, 1–10. <https://doi.org/10.1155/2014/892170>
- ❓ Toshmatov, Z. O., Li, J., Eshbakova, K. A., Tang, D., Xin, X., & Aisa, H. A. (2019). New monoterpene glucosides from *Dracocephalum komarovi* and their anti-inflammatory activity. *Phytochemistry Letters*, 33, 102–105. <https://doi.org/10.1016/j.phytol.2019.08.005>
- ✓ Utey, H. G., Bernheim, F., & Hochstein, P. (1967). Effect of sulfhydryl reagents on peroxidation in microsomes. *Archives of Biochemistry and Biophysics*, 118(1), 29–32. [https://doi.org/10.1016/0003-9861\(67\)90273-1](https://doi.org/10.1016/0003-9861(67)90273-1)
- ✓ Van Doan, H., Hoseinifar, S. H., Ringø, E., Ángeles Esteban, M., Dadar, M., Dawood, M. A., & Faggio, C. (2020). Host-associated probiotics: A key factor in sustainable aquaculture. *Reviews in Fisheries Science & Aquaculture*, 28(1), 16–42. <https://doi.org/10.1080/23308249.2019.1643288>
- ❓ Wayne, P. (2007). *Implementation guide of POCT for HEALTH care providers* (pp. 1–37). CLSI.

- ✓ Wendakoon, C. N., & Sakaguchi, M. (1993). Combined effect of sodium chloride and clove on growth and biogenic amine formation of *Enterobacter aerogenes* in mackerel muscle extract. *Journal of Food Protection*, 56(5), 410–413. <https://doi.org/10.4315/0362-028X-56.5.410>
- ✓ Xu, J., Zhou, F., Ji, B. P., Pei, R. S., & Xu, N. (2008). The antibacterial mechanism of carvacrol and thymol against *Escherichia coli*. *Letters in Applied Microbiology*, 47(3), 174–179.
- ✓ Yanishlieva, N. V., Marinova, E. M., Gordon, M. H., & Raneva, V. G. (1999). Antioxidant activity and mechanism of action of thymol and carvacrol in two lipid systems. *Food Chemistry*, 64(1), 59–66. [https://doi.org/10.1016/S0308-8146\(98\)00086-7](https://doi.org/10.1016/S0308-8146(98)00086-7)
- ❓ You, S. H., Yoon, M. Y., & Moon, J. S. (2021). Antioxidant and anti-inflammatory activity study of fulvic acid. *Journal of Natural Science, Biology and Medicine*, 12(3), 285–289.
- ❓ Yousefi, M., Farsani, M. N., Ghafarifarsani, H., Hoseinifar, S. H., & Van Doan, H. (2021). The effects of dietary supplementation of mistletoe (*Viscum album*) extract on the growth performance, antioxidant, and innate, immune responses of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 536, 736385. <https://doi.org/10.1016/j.aquaculture.2021.736385>
- ✓ Yousefi, M., Ghafarifarsani, H., Hoseini, S. M., Hoseinifar, S. H., Abtahi, B., Vatnikov, Y. A., Kulikov, E. V., & Van Doan, H. (2022). Effects of dietary thyme essential oil and prebiotic administration on rainbow trout (*Oncorhynchus mykiss*) welfare and performance. *Fish & Shellfish Immunology*, 120, 737–744. <https://doi.org/10.1016/j.fsi.2021.12.023>
- ✓ Yousefi, M., Ghafarifarsani, H., Hoseinifar, S. H., Rashidian, G., & Van Doan, H. (2021). Effects of dietary marjoram, *Origanum majorana* extract on growth performance, hematological, antioxidant, humoral and mucosal immune responses, and resistance of common carp, *Cyprinus carpio* against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 108, 127–133. <https://doi.org/10.1016/j.fsi.2020.11.019>
- ✓ Zharif, N., Santosh, F., Kiran, C. N., Fadli, A., Ibrahim, A., & Nizam, G. (2018). Synergistic effect of ethanolic extract of *Melastoma malabataricum* leaves and antibiotics. *International Journal of Medical Toxicology & Legal Medicine*, 21(3and4), 167–170. <https://doi.org/10.5958/0974-4614.2018.00059.1>

**How to cite this article:** Hafsan, H., Bokov, D., Abdelbasset, W. K., Kadhim, M. M., Suksatan, W., Majdi, H. S., Widjaja, G., Jalil, A. T., Qasim, M. T., & Balvardi, M. (2022). Dietary *Dracocephalum kotschyi* essential oil improved growth, haematology, immunity and resistance to *Aeromonas hydrophila* in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Research*, 00, 000–000. <https://doi.org/10.1111/are.15829>

----- End of Document -----





---

**Corrections successfully submitted for ARE\_EV\_ARE15829, Dietary  
Dracocephalum kotschyi essential oil improved growth, haematology, immunity  
and resistance to Aeromonas hydrophila in rainbow trout (Oncorhynchus mykiss).**

---

Wiley Online Proofing <notifications@eproofing.in>

Sun, Mar 13, 2022 at 11:00 PM

Reply-To: eproofing@wiley.com

To: hafsan.bio@uin-alauddin.ac.id

Cc: iselvaraj@wiley.com



## Corrections successfully submitted

Dear **Hafsan Hafsan**,

Thank you for reviewing the proof of the **Aquaculture Research** article **Dietary  
Dracocephalum kotschyi essential oil improved growth, haematology, immunity and  
resistance to Aeromonas hydrophila in rainbow trout (Oncorhynchus mykiss)**.

[View Article](#)

This is a read-only version of your article with the corrections you have marked up.

If you encounter any problems or have questions please contact us at  
([iselvaraj@wiley.com](mailto:iselvaraj@wiley.com)). For the quickest response include the journal name and your  
article ID (found in the subject line) in all correspondence.

Best regards,  
Aquaculture Research