

The stability of Phytase activity from Burkholderia sp. strain

HF.7

by Hafsan Hafsan

Submission date: 17-Oct-2022 03:11PM (UTC+0700)

Submission ID: 1927513088

File name: ity-of-phytase-activity-from-burkholderia-sp-strain-hf7-7582.pdf (330.67K)

Word count: 2377

Character count: 12941



The stability of Phytase activity from *Burkholderia* sp. strain HF.7

Hafsan ¹, Laily Agustina ², Asamuiddin Natsir ², Ahyar Ahmad ³

¹ Department of Biology, Faculty of Science and Technology, Makassar Islamic State Alauddin University, Gowa, 92113, South Sulawesi, INDONESIA

² Department of Nutrition and Feed Livestock, Faculty of Animal Husbandry, Hasanuddin University, Makassar 90245, INDONESIA

³ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Hasanuddin University, Makassar 90245, INDONESIA

*Corresponding author: hafsan.bio@uin-alauddin.ac.id

Abstract

Objectives: Phytase is an enzyme that hydrolyzes phytic acid but is not produced in the monogastric gastrointestinal tract of poultry. This study aims to discover the stability of extracellular phytase *Burkholderia* sp. strain HF.7 under a range of temperature, pH, and protease activity conditions including those similar to the digestive tract of poultry. **Material and Methods:** Temperature and pH stability tests were carried out at various temperatures (20-65 °C) and pH (1-8) by incubating the phytase in a Na-acetate buffer. The stability of phytase protease was further tested by adding the following levels of proteases (pepsin and/or trypsin): P0: without protease, P1: pepsin (5000 units/mL), P2: trypsin (5000 units/mL) and P3: pepsin + trypsin (2500 units/mL for each). ANOVA was used to analyze the protease stability experiments ($\alpha=0.05$). **Finding:** The results showed that *B. sp.* strain HF.7 phytase had an optimum activity at 37 °C and pH 4; similar to the condition of the poultry digestive tract in general. **Conclusion:** These findings suggest that *B. sp.* strain HF.7 has potential to be used in poultry feed to increase productivity.

Keywords: *Burkholderia* sp., physiology, phytase, poultry digestive tract, stability

Hafsan, Agustina L., Natsir A., Ahmad A (2020) The stability of Phytase activity from *Burkholderia* sp. strain HF.7. Eurasia J Biosci 14: 973-976.

© 2020 Hafsan et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Phytase is an enzyme that hydrolyzes phytic acid but is not produced in the monogastric gastrointestinal tract of poultry. Since it is unable to be processed, phytic acid, as the largest component of phosphorus in poultry feed material cannot be digested and is excreted and thus can become an environmental pollutant. As an anti-nutritional substance, phytic acid also has a negative effect on the absorption of other nutrients by poultry. In natural conditions, phytic acid can bind to proteins and two-divalent minerals needed for growth, causing minerals and proteins cannot be absorbed and digested properly (Widowati et al. 2001). The addition of the phytase enzyme into feed is expected to reduce the activity of phytic acid in the digestive tract so that nutrients can be absorbed optimally and efficiently. Some studies have reported that supplementation of phytase in poultry feed is able to increase its productivity, among Widowati (2016) reported that phytase can improve the digestibility of the protein, amino acids, calcium and phosphorus that has implications for improved performance of poultry.

However, as an enzyme, phytase can be expected to have a certain range of conditions in which it functions,

as well as conditions where it would be expected to be degraded (Anselme 2006). Thus, in practical applications of enzymes for the improvement of poultry feed, the enzymes used must retain protease activity as well as stability at temperatures, pH, found in poultry digestive tracts (Abondano 2009). Production of an extracellular phytase of an endophyte bacteria of corn, *Burkholderia* sp. strain HF.7, has been optimized and is available commercially. However, the stability of the enzyme in the gastrointestinal tract of poultry has not been investigated, so this study aims to discover the stability of extracellular *B. sp.* phytase under a range of temperature, pH, and protease activity conditions including those similar to the digestive tract of poultry. (Hamad et al 2016).

Received: January 2020

Accepted: April 2020

Printed: April 2020

MATERIALS AND METHODS

Materials

A pure bacterial culture of *B. sp.* strain HF.7 (a thermophile bacteria from Sulili hot springs in the district Pinrang of South Sulawesi Indonesia), crude extracellular phytase extract (using phytase production medium as described by El-Toukhy et al. (2013)), pure extracellular phytase extract (extract purification was done by using column chromatography), and Luria Bertani Medium (LB) (Sigma, Catalog P-7012), NaHCO₃, and pancreas (Sigma, Catalog P-3292) were used in this study. Equipment used were dialysis tubes (Sigma, catalog D-9652-100FT), parafilm, NaCl, EDTA, sodium succinate, and sulfuric acid. Therefore, the conducted research is not related to either human or animal use. This research was conducted in the laboratory of Microbiology, Faculty of Science and Technology, Alauddin State Islamic University of Makassar.

In Vitro Analysis of Phytase Stability

Temperature and pH stability was measured by observing the relative activity of crude and purified extracellular phytase extract. The test was carried out by exposing the phytase to various temperatures (20, 30, 37, 40, 45, 50, 55, 60, and 65 °C) for 10 min, followed by a phytase activity trial. The pH stability test was performed by incubating the extracellular phytase extract in Na-acetate buffer at various pH (pH 1, 2, 3, 4, 5, 6, 7, 8).

Analysis of Phytase Stability against Protease

Hydrolytic activity by extracellular phytase was analyzed by digestible P content in vitro on rice bran, based on the method of Wu *et al.* (2002). The experimental treatments were hydrolytic activity test without protease (P0), with pepsin addition (5000 units/mL) (P1), pepsin addition (5000 units/mL), and combination of pepsin + trypsin addition (@ 2500 units/mL) (P3). In vitro digestibility was observed based on P-value measurements using a spectrophotometer ($\lambda=415$ nm). P digestibility value was calculated by observing *p value* of feed after hydrolysis in vitro on each experimental treatment. The increase was compared with the *p value* of feed ingredients prior to hydrolysis by phytase.

Statistical Analysis

Phytase activity at various temperatures and pH was descriptively analyzed by plotting the mean values of relative enzyme activity over the temperature and pH ranges studied. Statistical analysis of protease stability was performed by using Analysis of Variance (ANOVA) ($\alpha=0.05$).

RESULTS AND DISCUSSION

The measurement results of crude and purified extracellular phytase extract is presented in Table 1.

Table 1. Analysis of variance of P concentration phytase-hydrolyzed rice bran

| Variance Source | Sum of Squares | Db | Middle Squares | F Value | Sig. |
|-----------------|----------------|----|----------------|---------|-------|
| Treatment | 1.823 | 3 | 6.076 | 1.156 | 0.384 |
| Error | 4.204 | 8 | 5.255 | | |
| Total | .000 | 12 | | | |

Values of extracellular phytase activity were higher for the pure than the crude extracellular phytase. In production, phytase is obtained by extraction from the culture medium by centrifugation. Then, the extract is purified to certain concentrations based on its purpose (Clarkson et al. 2001). Pre-purification was carried out by fractionation with ammonium sulfate, then dialysis, and then further purification by gel filtration chromatography Sephadex G-100 (Raju et al. 2007). Protein purification is required to eliminate contaminants and various inhibitors, improving the activity of the purified product (Berka et al. 1998).

The results of the phytase stability tested at various temperatures and pH are presented in Figs. 1 and 2. The phytase of *B. sp.* strain HF.7 had activity in all pH. Phytase activity was detected from pH 1 and increased at pH 2 to reach an optimum at pH 4 and pH 5 until its activity decreased at pH 8. The difference in activity at various pH was likely caused by the intramolecular changes from enzymes caused by ionization binding and releasing protons (hydrogen ions) in amino, carboxyl, and other functional groups. If the intramolecular change is too huge it can denature the enzyme and decrease the activity (Alberty 2007). Phytases are a heterologous group of enzymes able to hydrolyze phosphate esters and have optimal at low pH. At neutral and basic pH, catalytic activity decreases. This is due to the instability of structures of the enzyme protein molecules thus causing structural changes under these pH conditions (Berka et al. 1998).

The condition of pH of the chicken gastrointestinal tract is pH 4.5 in the cache, pH 4.4 in the proventriculus, 2.6 in the gizzard, pH 5.7–6.0 in the duodenum, pH 5.8 in jejunum, pH 6.3 in the ileum, pH 6.3 in the colon, and pH 5.7 in ceca (Wyss et al. 1999). Extracellular phytase of *B. sp.* strain HF.7 shows stable activity at pH 4–6. Thus, this phytase may be active at pH found along most of the poultry's digestive tract.

The extracellular phytic stability of *B. sp.* strain HF.7 at various temperatures (Fig. 2) shows that rising temperatures caused an increase in enzyme activity, reaching an optimum point at 37 °C. If the temperature was increased to 40 °C, the activity of the phytase decreased. At 65 °C, there was still active, but the reaction was significantly decreased. At first, with increasing temperature, enzyme reaction also increased rapidly likely due to increased kinetic energy that accelerates vibration, translation, and rotation of the enzyme along with the substrate thereby increasing the chemical reaction. At temperatures above the optimum,




Fig. 1. Phytase activity of *Burkholderia* sp. strain HF.7 at various pH




Fig. 2. Phytase activity of *Burkholderia* sp. strain HF.7 at various temperatures

the conformation of enzyme proteins and substrates likely change, inhibiting the reactive group entering the active site of the reaction and affecting catalysis activity.

The optimum temperature and pH of phytase depends on the type and source of the enzyme. Several pHs and optimum temperatures of the microbial-produced phytases have been reported. Phytase from *B. subtilis* (natto) N-77 with optima at 60 °C and pH 6.0–6.5, and *Enterobacter* sp. 4, with optima at 50 °C, pH 4.5. Phytase is also produced by *Aspergillus niger* (optimal: 58 °C, pH 5.5), *Schwanniomyces castellii* (optimal: 7.7 °C, pH 4.4), and *Klebsiella aerogenes* (optimal: 45 °C and pH 7.0) [Unpublished result].

The results of the P digestibility evaluation to analyze the stability differences of phytase are presented in **Table 1**. Stability tests are important to observe whether enzymes in the animal feed needs specific requirements to avoid being degraded by proteases in the poultry digestive tracts (Kornegay 2000, Wang et al. 2007).

There was no significant difference between treatments at P concentration in rice bran after hydrolysis by phytase in vitro ($p < 0.05$). P concentration remained elevated after hydrolysis by phytase, either

with the addition of pepsin or trypsin and a combination of both. This shows that phytase of *Burkholderia* sp. HF.7 is not degraded by pepsin, trypsin, or combination of two enzymes. Thus, the phytase activity remains stable as evidenced by an increase in P concentration with the addition of phytase. This, according to the study by Monteiro et al. (2015) which shows that phytase demonstrated strong resistance toward pepsin and trypsin, and it retained more than 90% residual activity for both enzymes after 1 h treatment.

CONCLUSION

In conclusion, extracellular phytase enzyme of *Burkholderia* sp. HF.7 strain has a higher activity at higher purity and has good stability at pH, temperature, and is resistant to degradation by proteases pepsin and trypsin, and thus is compatible with general poultry digestive tract conditions.

ACKNOWLEDGEMENT

Authors thank to Alauddin State Islamic University of Makassar for facilitated this research.

REFERENCES

- Abondano E (2009) Enzymatic Functions: Protease, Amylase, and Phytase. North Dakota State University, USA.
- Alberty RA (2007) Changes in binding of hydrogen ions in enzyme-catalyzed reactions. *Biophysical Chemistry* 125: 328-333.
- Anselme P (2006) Considerations on the Use of Microbial Phytase. CEFIC Inorganic Feed Phosphates, Brussels.
- Berka RM, Rey MW, Brown KM, Byun T, Klotz AV (1998) Molecular characterization and expression of a phytase gene from the thermophilic fungus *Thermomyces lanuginosus*. *Applied and Environmental Microbiology* 64: 4423-4427.
- Clarkson K, Jones B, Batt R, Bower B, Chotani G, Becker T (2001) Enzymes: screening, expression, design, and production. In: *Enzyme in Farm Animal Nutrition* (Eds. M. R. Bedford and G. G. Partridge), CABI Publishing, UK: pp. 315.
- El-Toukhy NMK, Youssef AS, Mikhail MGM (2013) Isolation, purification and characterization of phytase from *Bacillus subtilis* MJA. *African Journal of Biotechnology*, 12: 2957-2967.
- Hamad, M. N., Ismail, M. M., & El-Menawy, R. K. (2016). Effect of Jameed Form on the Chemical Composition, Rheological and Microbial Properties. *Journal of Food Technology Research*, 3(2), 72-87.
- Kornegay ET (2000) Digestion of phosphorus and other nutrients: the role of phytases and factors influencing their activity. In: *Enzymes in Farm Animal Nutrition* (Eds. M. R. Bedford and G. G. Partridge), CABI Publishing, UK: pp. 237.
- Monteiro PS, Guimarães VM, de Melo RR, de Rezende ST (2015) Isolation of a thermostable acid phytase from *Aspergillus niger* UFV-1 with strong proteolysis resistance. *Brazilian Journal of Microbiology*, 46: 251–260.
- Raju, S., Hollis, K. and Neglen, P. 2007. Use of compression stockings in chronic venous disease: patient compliance and efficacy. *Annals of Vascular Surgery* 21: 790-795.
- Setiawati D, Sukanto B, Wahyuni HI (2016) Pengimbuhan Enzim Fitase dalam Ransum Ayam Pedaging Meningkatkan Pemanfaatan Kalsium untuk Pertumbuhan Tulang dan Bobot Badan (Adding Phytase Enzymes on Broiler Ration Increasing Calcium Utilization for Bone Growth and Body Weight Gain). *Jurnal Veteriner* 17(3): 468-476.
- Wang Y, Gao X, Su Q, Wu W, An L (2007) Cloning, expression, and enzyme characterization of an acid heat-stable phytase from *Aspergillus fumigatus* WY-2. *Current Microbiology* 55: 65-70.
- Widowati S, Sukamo L, Raharto P, Thontowi A (2001) Studi pengaruh penambahan mineral terhadap aktivitas protease dari *Bacillus circulans* [Study of mineral adding effect toward protease activity from *Bacillus circulans*]. *Proceeding on Research Seminar Pioneer and Plant Biotechnology*, January 30-31, BB Biogen, Indonesia.
- Wu YB, Ravidran V, Pierce J, Hendriks WH (2002) Influence of three preparation in broiler diets based on wheat or corn: In vitro measurements of nutrient release. *The Journal of Poultry Science* 3: 450.
- Wyss M, Pasamontes L, Friedlein A, Rémy R, Tessier M, Kronenberger A, et al. (1999) Biophysical characterization of fungal phytases (myo-inositol hexakisphosphate phosphohydrolases): molecular size, glycosylation pattern and engineering resistance. *Applied and Environmental Microbiology* 65: 359-366.

The stability of Phytase activity from Burkholderia sp. strain HF.7

ORIGINALITY REPORT

7%

SIMILARITY INDEX

4%

INTERNET SOURCES

5%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1 tsukuba.repo.nii.ac.jp 1%
Internet Source

2 www.scirj.org 1%
Internet Source

3 www.innspub.net 1%
Internet Source

4 www.neliti.com 1%
Internet Source

5 www.aensiweb.net 1%
Internet Source

6 Dhurgham Ismael Baqer Al ALNABI, Sarmad Ghazi AL-SHAWI, Zena Kadhim AL-YOUNIS, Wael Ali SWADI et al. "Heavy metal in the soil-grain-food path: an overview of the role of Mycotoxins in potential hazards associated with animal products", Food Science and Technology, 2022 <1%
Publication

7 Elena Ulrikh, Olga Babich, Stanislav Sukhikh. "Use of sweet yellow clover (*Melilotus officinalis*) extract in sheep feeding", E3S Web of Conferences, 2021
Publication <1 %

8 M. V. Chirkova, P. V. Pivovarov, E. G. Litvinova, V. S. Khotimskii. "Chemical modification of poly(vinyltrimethylsilane) and poly(1-trimethylsilyl-1-propyne) using highly reactive metallating systems", Polymer Science Series A, 2006
Publication <1 %

9 Sjamsiah, N. Ramli, R. Daik, M. A. Yarmo, Z. Ajdari. "Nutritional study of Kapparazii powder™ as a food ingredient", Journal of Applied Phycology, 2013
Publication <1 %

10 "Enzyme Technology", Springer Science and Business Media LLC, 2006
Publication <1 %

11 Cho, J.. "Avian multiple inositol polyphosphate phosphatase is an active phytase that can be engineered to help ameliorate the planet's "phosphate crisis"', Journal of Biotechnology, 20061101
Publication <1 %

12 ascidatabase.com
Internet Source <1 %

13

www.locus.ufv.br

Internet Source

<1 %

14

Thermophilic Microbes in Environmental and Industrial Biotechnology, 2013.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On