

Functional Characteristics of Lactobacillus Fermentum Origin of Whey (Waste Processing) Dangke Products as Probiotic Candidate

by Hafsan Hafsan

Submission date: 17-Oct-2022 02:00PM (UTC+0700)

Submission ID: 1927480157

File name: 134-Article_Text-324-1-10-20200831.pdf (730.01K)

Word count: 3910

Character count: 20516



Functional Characteristics of *Lactobacillus Fermentum* Origin of Whey (Waste Processing) Dangke Products as Probiotic Candidate

Authors:

¹Hafsan*, ²Fatmawati Nur, ³Muhammad Halifah Mustami, ⁴Khaerani Kiramang, ⁵Rahmaniah

^{1,2,3,4,5}Faculty of Science and Technology, Universitas Islam Negeri Alauddin, Gowa, 92118, South Sulawesi, Indonesia

*Corresponding Author:

*Hafsan

*Faculty of Science and Technology, Universitas Islam Negeri Alauddin, Gowa, 92118, South Sulawesi, Indonesia

E-mail: hafsan.bio@uin-alauddin.ac.id

Abstract:

This study was aimed to determine the potential of isolates of waste Dangke-production, which is one of the traditional dairy products distinctive of South Sulawesi, i.e. the district Enrekang. Process of his production has left the waste in the form of Whey, which is a liquid or water part of milk that remains after the separation of curd as a coagulation form of milk protein with proteolytic enzymes. Lactic acid bacteria have been isolated from dangke is *Lactobacillus fermentum* is. Furthermore, as local probiotic candidate, conducted many diagnostic examinations, i.e. isolates resistance to acidic pH, resistance to bile salts, the ability of aggregation and adhesion to the surface of the small intestine, and the inhibitory effect on pathogenic bacteria such as *Salmonella thypi* and *Staphylococcus aureus*. Examinations showed that *L. fermentum* isolated from Whey of Dangke has excellent potential to be developed as probiotics. This potency is caused by resistance to acidic pH is pH 2, 3 and 4 as well as able to grow at a concentration of 0,1% bile salts and 0,3 % for 2 x 24-hour incubation with different degrees of growth. *L. fermentum* from Whey of Dangke been able to aggregate with a positive value indicated by the formation of aggregates which forms a precipitate in the base the tube and the supernatant look clear. *L. fermentum* isolates were also able to attach to the surface of the small intestine in vitro, as well as having the ability to inhibit the growth of pathogenic bacteria such as *S. aureus* and *S. thypi*.

Keywords: *Lactobacillus fermentum*, Probiotic, Dangke, Enrekang.

1. Introduction

Dangke is one of the traditional dairy products that is unique to South Sulawesi, namely Enrekang district [1]. Dangke is processed with the essential

ingredients of cow or buffalo milk which is heated to a specific temperature and added with papaya sap to compact and separate curd and Whey [2]. The separation of curd and Whey is related to the effect

of papaya sap which contains papain, a type of proteolytic enzyme. Papain will break the peptide bonds in the residues of asparagine-glutamine, glutamate-alanine, leucine-valine and phenylalanine-tyrosine [3]. The process of making Dangke leaves waste in the form of Whey which is serum or part of the water from the milk that remains after the separation of curd as a result of coagulation of milk protein with proteolytic enzymes. Every 10 litres of milk that are coagulated during the processing process produces about 6 - 9 litres of whey [1]. As a dangke processing waste, Whey still contains about 55% of the total nutrients from milk such as lactose, protein, fat, vitamins, and mineral salts. This condition allows bacteria, especially lactic acid bacteria (LAB) to make it a medium for growth and development, considering that LAB is a natural flora on a substrate containing milk [4].

Lactic acid bacteria are always a concern for researchers because in general, they have the potential to be developed as probiotics for humans, land and aqua²⁰ animal farm. This is based on the fact that LAB is a natural inhabitant o²² the digestive tract of humans and animals with the ability to tolerate environmental conditions with acid and bile salt from the digestive tract. Lactic acid bacteria also function to convert lactose into acetic acid, thereby lowering the pH of the digestive tract and naturally preventing the colonization of various pathogenic bacteria so that in general it can improve the condition of the digestive tract and will improve health [5, 6]. Besides, it can reduce ex²⁵ cholesterol in the blood, thereby reducing the risk of heart disease and other blood vessel disorders [7, 6, 8].

¹³ The variety of lactic acid bacteria that have been isolated from various sources such as from milk and other processed products allows the increase in LAB types that have the potential as probiotics with their respective characteristics [9]. The LAB group used as probiotics must meet several requirements as probiotics, including purity (healthy for the digestive tract), being non-pathogenic⁷ and resistant to antibiotics. Functional aspects, such as the ability to survive and survive in the digestive tract and are antagonistic to pathogenic bacteria [10]. Probiotic strains must also be resistant and survive during the food processing and storage process, be easily

applied to food products and are resistant to psychoche²¹cal processes in food [11]. Based on this, this study aims to determine the potential of isolates from the waste of making dangke, namely *Lactobacillus fermentum* as a probiotic through evaluation.

2. Methods

2.1 Tools and Materials

The tools and materials used in this study were male mice (*Mus musculus*) that had obtained a certificate of ethical eligibility from the Research Ethics Commission with experimental animals at UIN Alauddin, Eppendorf tubes, incubators, ovens, shaker incubators, autoclaves, water bath, Erlenmeyer, Becker glass, hot plates and stirrers, bunsen, ketamine injection, Petri dishes, holders, measuring cups, test tubes and tube racks, micropipettes (0.1, 1.0 and 10 mL), loop needles, pH meter, thermometer, spectrophotometer, analytical balance, oven, laminar airflow, thermometer, digital camera, *L. fermentum* culture from Whey Dangke, *Salmonella thypii* and *Staphylococcus aureus*, NaOH, HCl, glycerol, Aquadest, Alcohol, Sprites, Nutrient Agar (NA) medium, Hilton Muller Agar medium (MHA), medium de Mann, Rogosa and Sharpe (MRS), Bacto agar, Phosphate Buffer Saline (PBS), wrapper, tissue paper, cotton, filter paper, aluminium foil.

2.2 Data collection procedures

Characterization of the functional properties of the isolates was initiated by analyzing the viability against low pH and bile salts [4, 12, 13]. Testing for resistance to pH is b⁶ using liquid MRS media which is pH adjusted with pH 1 treatment; pH 2; pH 3; and pH 4 using a pH-meter of 5 mL. Each treatment was inoculated with the test of bacterial culture as much as 10% of the media volume. The control was liquid MRS medium without the addition of bacteria. After incubation for 1 and 2 x 24 hours at 37°C, observations were made by comparing the turbidity of culture with the control medium. Test of resistance to bile salts with liquid MRS media supplemented with oxgall 0.1%, 0.3% and 0.5%. Each treatment was inoculated with test bacteria as much as 10% of the media volume. The control was liquid MRS medium without the addition of bacteria, after incubating for 1 and 2 x 24 hours at 37 °C under anaerobic conditions.

The ability of *L. fermentum* bacteria that has also been tested is the ability of aggregation and adhesion to the surface of the small intestine [14, 15]. Aggregation test is carried out to determine the ability of a bacterium to form colonies (aggregates). The aggregation test was carried out by growing it on MRSB for 1 x 24 hours at 37°C. Aggregation is considered positive if the MRS shows clear aggregates (particles such as sand) to form a sediment at the bottom of the tube and the supernatant will appear bright. In vitro testing of the attachment of Lactic Acid Bacteria to the surface of the small intestine begins with taking a sample of the intestine on the mice, starting with anaesthesia with the mice, then performing surgery starting from the chest to the abdomen. The small intestine (ileum) is cut into $\pm 1 \text{ cm}^2$. The intestinal sample was further washed with PBS. The washed sample was put into tissue basket, which was then soaked with PBS solution at 4°C for a few minutes and after that was washed three times. The attachment test was carried out by incubating the intestinal sample into a petri dish containing a solution of 1.2×10^9 cell/ml bacterial suspension in PBS at 37°C for 30 minutes then rinsed with PBS. The presence of adhered bacteria was evaluated qualitatively by observing the growth after cultivation on MRSA media from post-incubation intestinal suspension in a solution of 1.2×10^9 cell ml⁻¹ bacterial suspension in PBS.

The antimicrobial ability of *L. fermentum* against pathogenic bacteria such as *S. thypii* and *S. aureus* was each carried out by first making a test pathogenic bacterial suspension which is equivalent to 0.5 Mc. Farland units by taking one ose of the tested bacterial colony from pure culture and Suspended into 5 mL of sterile NaCl. A total of 50 μL of bacterial suspension was pipette and put into a test tube containing 10 mL of MHA medium at 45°C. The medium containing the bacterial suspension was poured into a petri dish containing 20 mL of solidified MHA medium. The Petri dishes were shaken so that the bacterial suspension was evenly distributed, then 3 pencandang fruit were placed on each petri dish. Next, 250 μL of LAB isolate from Dangke whey was added to each pen. Subsequently incubated for 24 hours at 37°C. The clear zone formed around the sample wells indicated the ability of LAB isolates from Dangke Whey to inhibit the growth of each test pathogenic bacteria. [16]

3. Results

As a probiotic candidate, a microbial isolate must fil several functional characteristics, including resistance to low pH, bile salts, aggregation ability, coaggregation and adhesion. Besides, it can inhibit the growth of pathogenic bacteria [17]. Table 1 describes some of the properties of LAB isolates from Dangke Whey as probiotic candidates.

Table 1: The results of examination the functional properties of *L. fermentum* from Whey of Dangke

Functional characteristic test	Test result*
Viability at Low pH:	- - -
• pH 1	+++
• pH 2	+++
• pH 3	+++
• pH 4	+++
Viability at Bile salt:	
• 0.1%	+++
• 0.3%	+++
• 0.5%	+++
Aggregation	- - -
In vitro adhesion	+++
Inhibition against <i>S.typhi</i>	+++
The inhibition against <i>S. aureus</i>	+++

*(-) = negative reaction on each of the replicas

(+) = positive reaction on each of the replicas

Evaluation the resistance of *L. fermentum*, to low pH was carried out by growing it in MRSB media which had the pH set to 1, 2, 3, and 4, then incubated for 1 and 2 x 24 hours. Analysis of the growth was carried out qualitatively by looking at the level of turbidity of the media inoculated with lactic acid bacterial cultures after incubation and comparing it with control media (pH 6.2 according to normal pH of MRSB). Table 1 shows that the isolates identified as *L. fermentum* were able to grow on MRSB with a pH of 2, 3 and 4. Whereas at pH one the isolates did not indicate growth which was marked by no change in the turbidity of the MRSB medium used, as the control.

In addition to testing the ability to grow bacteria on media with low pH, *L. fermentum* was also tested for its ability to grow on media containing oxgall (bile salt). The presence of oxgall in the media is intended

to create conditions such as those in digestion where bile salts are excreted into the digestive tract. In this test, all LAB used were grown in MRSB media containing 0.1%, 0.3% and 0.5% oxgall, then incubated for 1 and 2 x 24 hours. Analysis of growth was carried out qualitatively by looking at the level of turbidity of the media inoculated with lactic acid bacterial cultures after incubation and comparing it with control media (without oxgall).

The results obtained (Table 1) showed that the tested isolates were able to grow at bile salt concentrations of 0.1% and 0.3% for 2 x 24 hours of incubation with different growth degrees (based on turbidity levels). At a bile salt concentration of 0.5% and an incubation time of 2 x 24 hours, from three replications performed, the isolates did not grow (the media was not cloudy). Isolates indicated growth at bile salt concentrations of 0.1% and 0.3% after 2 x 24 hours. This phenomenon occurs because bile salts are bactericidal so that they inhibit bacterial growth, and possibly some of the inoculated bacteria die. Bacteria that still survive require a long adaptation time so that new growth is seen after 48 hours. These results are consistent with research conducted by [7, 18, 19, 20]. Lactic acid bacteria grown on media containing bile salts experienced a delay in growth because it required a longer adaptation than bacteria grown on media that did not contain bile salts.

The aggregation test performed on *L. fermentum* from Dangke whey showed a positive aggregation assessment. This is evidenced by the presence of deposits at the bottom of the tube accompanied by clear liquid on it. According to [21] stated that aggregation is considered positive if the MRSB media contains clear aggregates (particles such as sand), forming sediment at the bottom of the tube and the supernatant looks clear. The formed precipitate indicates the formation of an aggregate or colony in the tested lactic acid bacteria. The precipitate is thought to have formed because the lactic acid bacteria grew and multiplied under media conditions, temperature and incubation time and interacted to form colonies. The aggregation ability of isolates in single and paired forms is influenced by the production of Extra Cellular Polysaccharides (PSE) on the outer cell membrane, in addition to the force of gravity and the motility or movement of bacteria. If PSE production is low and the chemical

bonds of the paired bacterial cell wall components are weak, the aggregation will not occur [22, 23, 24].

The ability of *L. fermentum* to inhibit the growth of pathogenic bacteria is mostly due to the presence of organic acids it produces. Organic acids produced by LAB result in the accumulation of acidic end products and a decrease in pH, which will inhibit the growth of both Gram-positive and Gram-negative bacteria [12, 25]. The activity of lipophilic acids such as lactic and acetic acids in the undissociated form can penetrate microbial cells and at higher intracellular pH, dissociate to produce hydrogen ions and interfere with the function of essential metabolites such as substrate translocation and oxidative phosphorylation, thereby reducing intracellular pH [12]. The inhibitory effect of organic acids comes primarily from the amount of acid that is not dissociated. Lipophilic acids such as lactic acid and acetic acid in their undissociated form can penetrate bacterial cells. [6] states that weak acids can reduce cytoplasmic pH, affect membrane structure and fluidity and chelate (bind) bacterial cell wall ions. The decrease in cytoplasmic pH will affect cell structural proteins, enzymes, nucleic acids and membrane phospholipids. The inhibitory ability of LAB metabolites against the growth of *S. thypi* and *S. aureus* is also caused by the production of bacteriocins [11, 26, 27, 28].

The mechanism of inhibition of *L. fermentum* against the growth of pathogenic bacteria is that in addition to causing interference with cell wall components, it can also inhibit it by reacting with cell membranes [6]. The ability of LAB metabolites to interfere with the components of the cell wall, one of which functions to protect the cytoplasmic membrane, causes bacterial cells to be sensitive to osmotic. The presence of osmotic pressure in bacterial cells will cause lysis. Organic acids produced by LAB will penetrate the cell wall and disrupt membrane permeability. Besides, once in the cytoplasm, it will dissociate to produce protons, and excess protons will cause the balance to be disturbed. This disturbance results in reduced cell energy for growth because it is diverted to balance protons and also disrupts the transport of amino acids and sugars [29].

4. Conclusion

The series of tests on the functional characteristics of *Lactobacillus fermentum* that have been carried out indicate that *L. fermentum* isolated from dangke-production waste has a great potential to be developed as a probiotic because it fulfills several requirements, namely, it has resistance to low pH and bile salts, has the ability of aggregation and attachment. On surface of the small intestine (in vitro), and has the ability to inhibit the growth of pathogenic bacteria, namely *Salmonella thypii* and *Staphylococcus aureus*.

Acknowledgements

This research was carried out based on funding from a competitive grant from the Institute for Research and Service, sourced from DIPA UIN Alauddin. Therefore, the research team would like to thank LP2M UIN Alauddin Makassar, as well as to all the personnel who were involved in the completion of this research.

Reference

- [1.] Fatma, Soeparno, Nurliyani, C. Hidayat, and M. Taufik, 'Karakteristik Whey Limbah Dangke dan Potensinya Sebagai Produk Minuman dengan Menggunakan *Lactobacillus acidophilus* FNCC 0051', *Agritech*, 2012, doi: 10.22146/agritech.9577.
- [2.] A. Syarifuddin, P. Hamsiohan, and M. Bilang, 'Characterization of edible film from dangke whey/pectin, beeswax, and butter aroma', in *AIP Conference Proceedings*, 2019, p. 020021, doi: 10.1063/1.5125525.
- [3.] G. N. Handayani, N. Ida, and A. R. R., 'Pemanfaatan Susu Skim Sebagai Bahan Dasar dalam Pembuatan Produk Olahan Makanan Tradisional Dangke dengan Bantuan Bakteri Asam Laktat', *Jf Fik Uinam*, 2014.
- [4.] M. Thirabunyanon, P. Boonprasom, and P. Niamsup, 'Probiotic potential of lactic acid bacteria isolated from fermented dairy milks on antiproliferation of colon cancer cells', *Biotechnol. Lett.*, vol. 31, no. 4, pp. 571–576, Apr. 2009, doi: 10.1007/s10529-008-9902-3.
- [5.] G. V. de Melo Pereira, B. de Oliveira Coelho, A. I. Magalhães Júnior, V. Thomaz-Soccol, and C. R. Soccol, 'How to select a probiotic? A review and update of methods and criteria', *Biotechnol. Adv.*, vol. 36, no. 8, pp. 2060–2076, Dec. 2018, doi: 10.1016/j.biotechadv.2018.09.003.
- [6.] A. T. Borchers, C. Selmi, F. J. Meyers, C. L. Keen, and M. E. Gershwin, 'Probiotics and immunity', *Journal of Gastroenterology*, 2009, doi: 10.1007/s00535-008-2296-0.
- [7.] Y. Ji *et al.*, 'Functionality and safety of lactic bacterial strains from Korean kimchi', *Food Control*, vol. 31, no. 2, pp. 467–473, Jun. 2013, doi: 10.1016/j.foodcont.2012.10.034.
- [8.] A. Emmawati, B. Sri, L. Suryaatmadja, L. Nuraida, and D. Syah, 'Characterization of Lactic Acid Bacteria Isolates from Mandai Function as Probiotic', *Agritech*, 2015.
- [9.] A. Ljungh and T. Wadström, 'Lactic acid bacteria as probiotics', *Current Issues in Intestinal Microbiology*, 2006.
- [10.] Y. D. Jatmiko, G. S. Howarth, and M. D. Barton, 'Assessment of probiotic properties of lactic acid bacteria isolated from Indonesian naturally fermented milk', in *AIP Conference Proceedings*, 2017, p. 050008, doi: 10.1063/1.5012732.
- [11.] H. Estifanos, 'Isolation and identification of probiotic lactic acid bacteria from curd and in vitro evaluation of its growth inhibition activities against pathogenic bacteria', *African J. Microbiol. Res.*, vol. 8, no. 13, pp. 1419–1425, Mar. 2014, doi: 10.5897/AJMR2014.6639.
- [12.] D. Charalampopoulos, S. S. Pandiella, and C. Webb, 'Growth studies of potentially probiotic lactic acid bacteria in cereal-based substrates', *J. Appl. Microbiol.*, vol. 92, no. 5, pp. 851–859, May 2002, doi: 10.1046/j.1365-2672.2002.01592.x.
- [13.] D. D. Pan, X. Q. Zeng, and Y. T. Yan, 'Characterisation of *Lactobacillus fermentum* SM-7 isolated from koumiss, a potential probiotic bacterium with cholesterol-lowering effects', *J. Sci. Food Agric.*, vol. 91, no. 3, pp. 512–518, Feb. 2011, doi: 10.1002/jsfa.4214.
- [14.] M. S. do Carmo *et al.*, 'Lactobacillus fermentum ATCC 23271 Displays In vitro Inhibitory Activities against *Candida* spp.', *Front. Microbiol.*, vol. 7, Oct. 2016, doi: 10.3389/fmicb.2016.01722.

- [15.] Q. Han, B. Kong, Q. Chen, F. Sun, and H. Zhang, 'In vitro comparison of probiotic properties of lactic acid bacteria isolated from Harbin dry sausages and selected probiotics', *J. Funct. Foods*, vol. 32, pp. 391–400, May 2017, doi: 10.1016/j.jff.2017.03.020.
- [16.] M. Olivares, M. P. Diaz-Ropero, R. Martin, J. M. Rodriguez, and J. Xaus, 'Antimicrobial potential of four Lactobacillus strains isolated from breast milk', *J. Appl. Microbiol.*, vol. 101, no. 1, pp. 72–79, Jul. 2006, doi: 10.1111/j.1365-2672.2006.02981.x.
- [17.] D. Granato, F. Nazzaro, T. C. Pimentelc, E. A. Esmerino, and A. G. da Cruz, 'Probiotic food development: An updated review based on technological advancement', in *Encyclopedia of Food Security and Sustainability*, 2018.
- [18.] D. I. A. Pereira, A. L. McCartney, and G. R. Gibson, 'An In Vitro Study of the Probiotic Potential of a Bile-Salt-Hydrolyzing Lactobacillus fermentum Strain, and Determination of Its Cholesterol-Lowering Properties', *Appl. Environ. Microbiol.*, vol. 69, no. 8, pp. 4743–4752, Aug. 2003, doi: 10.1128/AEM.69.8.4743-4752.2003.
- [19.] C. Tomaro-Duchesneau, M. L. Jones, D. Shah, P. Jain, S. Saha, and S. Prakash, 'Cholesterol Assimilation by Lactobacillus Probiotic Bacteria: An In Vitro Investigation', *Biomed Res. Int.*, vol. 2014, pp. 1–9, 2014, doi: 10.1155/2014/380316.
- [20.] A. Belicová, M. Mikulášová, and R. Dušinský, 'Probiotic Potential and Safety Properties of Lactobacillus plantarum from Slovak Bryndza Cheese', *Biomed Res. Int.*, vol. 2013, pp. 1–8, 2013, doi: 10.1155/2013/760298.
- [21.] A. C. Archer and P. M. Halami, 'Probiotic attributes of Lactobacillus fermentum isolated from human feces and dairy products', *Appl. Microbiol. Biotechnol.*, 2015, doi: 10.1007/s00253-015-6679-x.
- [22.] A. C. Archer and P. M. Halami, 'Probiotic attributes of Lactobacillus fermentum isolated from human feces and dairy products', *Appl. Microbiol. Biotechnol.*, vol. 99, no. 19, pp. 8113–8123, Oct. 2015, doi: 10.1007/s00253-015-6679-x.
- [23.] Y. Bao *et al.*, 'Screening of potential probiotic properties of Lactobacillus fermentum isolated from traditional dairy products', *Food Control*, vol. 21, no. 5, pp. 695–701, May 2010, doi: 10.1016/j.foodcont.2009.10.010.
- [24.] S. N. Casarotti, B. M. Carneiro, S. D. Todorov, L. A. Nero, P. Rahal, and A. L. B. Penna, 'In vitro assessment of safety and probiotic potential characteristics of Lactobacillus strains isolated from water buffalo mozzarella cheese', *Ann. Microbiol.*, vol. 67, no. 4, pp. 289–301, Apr. 2017, doi: 10.1007/s13213-017-1258-2.
- [25.] R. Martín *et al.*, 'Identification and evaluation of the probiotic potential of lactobacilli isolated from canine milk', *Vet. J.*, vol. 185, no. 2, pp. 193–198, Aug. 2010, doi: 10.1016/j.tvjl.2009.04.014.
- [26.] A. C. Gomes, A. A. Bueno, R. G. M. de Souza, and J. F. Mota, 'Gut microbiota, probiotics and diabetes', *Nutr. J.*, vol. 13, no. 1, p. 60, Dec. 2014, doi: 10.1186/1475-2891-13-60.
- [27.] R. Khalil, H. Mahrous, K. El-Halafawy, K. Kamaly, J. Frank, and M. El Soda, 'Evaluation of the probiotic potential of lactic acid bacteria isolated from faeces of breast-fed infants in Egypt', *African J. Biotechnol.*, 2007, doi: 10.5897/AJB2007.000-2114.
- [28.] A. Hutari, W. S. Jaseem, A. A. Hamid, and W. M. W. Yusoff, 'Screening of lactobacillus strains against salmonella both isolated from Malaysian free-range chicken intestine for use as probiotic', *Sains Malaysiana*, 2011.
- [29.] H. König and J. Fröhlich, 'Lactic Acid Bacteria', in *Biology of Microorganisms on Grapes, in Must and in Wine*, Cham: Springer International Publishing, 2017, pp. 3–41.

Functional Characteristics of Lactobacillus Fermentum Origin of Whey (Waste Processing) Dangke Products as Probiotic Candidate

ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

7%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	Jae-yeol Kim, Hyuk-Yoon Kwon. "Threat Classification Model for Security Information Event Management Focusing on Model Efficiency", Computers & Security, 2022 Publication	1%
2	onlinelibrary.wiley.com Internet Source	1%
3	vrf.iranjournals.ir Internet Source	1%
4	www.science.gov Internet Source	1%
5	commons.emich.edu Internet Source	<1%
6	patents.justia.com Internet Source	<1%
7	Dayong Ren, Chang Li, Yanqing Qin, Ronglan Yin et al. "In vitro evaluation of the probiotic and functional potential of Lactobacillus	<1%

strains isolated from fermented food and human intestine", Anaerobe, 2014

Publication

8

Rocío Martín, Mónica Olivares, Manuela Pérez, Jordi Xaus, Celina Torre, Leonides Fernández, Juan M. Rodríguez. "Identification and evaluation of the probiotic potential of lactobacilli isolated from canine milk", The Veterinary Journal, 2010

Publication

<1 %

9

annalsmicrobiology.biomedcentral.com

Internet Source

<1 %

10

ema-rachmawati.blogspot.com

Internet Source

<1 %

11

harvest.usask.ca

Internet Source

<1 %

12

link.springer.com

Internet Source

<1 %

13

ri.conicet.gov.ar

Internet Source

<1 %

14

www.animbiosci.org

Internet Source

<1 %

15

Larissa P. Margalho, Genesy P. Jorge, Deise A.P. Noleto, Christian E. Silva et al.
"Biopreservation and probiotic potential of a large set of lactic acid bacteria isolated from

<1 %

Brazilian artisanal cheeses: from screening to in product approach", Microbiological Research, 2020

Publication

16

Mariángeles Briggiler Marcó, María Florencia Zacarías, Gabriel Vinderola, Jorge A. Reinheimer, Andrea Quiberoni. "Biological and probiotic characterisation of spontaneous phage-resistant mutants of *Lactobacillus plantarum*", International Dairy Journal, 2014

Publication

17

Nurul Farhana Fazilah, Arbakariya B. Ariff, Mohd Ezuan Khayat, Leonardo Rios-Solis, Murni Halim. "Influence of probiotics, prebiotics, synbiotics and bioactive phytochemicals on the formulation of functional yogurt", Journal of Functional Foods, 2018

Publication

18

Wandee Sirichokchatchawan, Puwiya Pupa, Prasert Praechansri, Nutthee Am-in et al. "Autochthonous lactic acid bacteria isolated from pig faeces in Thailand show probiotic properties and antibacterial activity against enteric pathogenic bacteria", Microbial Pathogenesis, 2018

Publication

19

www.hindawi.com

Internet Source

<1 %

<1 %

<1 %

<1 %

20

www.mdpi.com

Internet Source

<1 %

21

www.tandfonline.com

Internet Source

<1 %

22

Martha Montserrat Castorena-Alba, Jesús Alberto Vázquez-Rodríguez, Manuel López-Cabanillas Lomelí, Blanca Edelia González-Martínez et al. "Cholesterol assimilation, acid and bile survival of probiotic bacteria isolated from food and reference strains", CyTA - Journal of Food, 2017

Publication

<1 %

23

Yuanyuan Feng, Lin Qiao, Rui Liu, Hongming Yao, Changbin Gao. "Potential probiotic properties of lactic acid bacteria isolated from the intestinal mucosa of healthy piglets", Annals of Microbiology, 2017

Publication

<1 %

24

"Lactic Acid Bacteria", Springer Nature, 2014

Publication

<1 %

25

"Probiotics and Health Claims", Wiley, 2011

Publication

<1 %

26

Muhamad Firdaus Syahmi Sam-on, Shuhaimi Mustafa, Mohd Termizi Yusof, Amalia Mohd

<1 %

Hashim et al. "Evaluation of three *Bacillus* spp. isolated from the gut of giant freshwater prawn as potential probiotics against pathogens causing Vibriosis and Aeromonosis", Microbial Pathogenesis, 2022

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On