




Contents lists available at BioMedSciDirect Publications

International Journal of Biological & Medical Research

Journal homepage: www.biomedscidirect.com

Original Article

Bioactivity Test and Interpretation of Secondary Metabolites of Sponge *Clathriareinwardhti*

Asri Saleha, Ahyar Ahmadb, Firdausb, HasnahNatsirb, Mashuri Masric

^aHasanuddin University, Graduate Program Department of Chemistry, Jl. Perintis Kemerdekaan Km. 10 Tamalanrea Makassar, South Sulawesi, Indonesia.^bHasanuddin University, Department of Chemistry, Faculty of Mathematic and Natural Science, Jl. Perintis Kemerdekaan Km. 10 Tamalanrea Makassar, South Sulawesi, Indonesia^cIslamic State University of Alauddin Makassar, Department of Biology, Faculty of Science and Technology, Jl. Sultan Alauddin, Samata Campus, Gowa Regency, South Sulawesi, Indonesia.

ARTICLE INFO

Keywords:

Clathriareinwardhti,
Bioactivity,
bacteria patogen,
Triterpenoid,
Steroid.

ABSTRACT

The isolation, structure determination and activity test against bacteria patogen secondary metabolites of sponges *Clathriareinwardhti* has been carried out. Separation techniques used consisted of maceration and fractination, while the structure of compounds were elucidated based on physical, spectroscopie UV and IR data. Two compounds that obtained were predicted as (1) Triterpenoid, and (2) Steroid. Compound (1) showed stronged toxicity against bacteria patogen 9,51 mm (2) showed high toxicity bacteria patogen with 11,41 mm.

© Copyright 2010 BioMedSciDirect Publications IJBMR - ISSN: 0976:6685. All rights reserved.

Introduction

Sponge has the ability to regenerate high. The body of sponge cut or damaged, it will regenerate into a whole again. The ability to regenerate its limits, mialnya pieces of sponge leuconoid must be greater than 0.4 mm and has several choanocyte cells that can regenerate into a new sponge is small. Sponge there are monosious (hermaphrodites) and some that are diosious. Sponge breed do asexually and sexually.

Asexual reproduction occurs with the formation of buds and sexually gemmule. Reproduksi do degan formation of gametes (sperm and ova) produced by koanosit. Sponge produce ova and sperm in the same individuals (hermaphrodites).

Respirators are composed of cells pinakosit (exterior) and koanosit (interior). Oxygen which has been captured by the two types of cells are circulated throughout the body by cells amubosit. As for the process of excretion of substances metabolic waste bins distributed from their internal body by amubosit.

Some researchers reported that the activity of the substance of the sea, among others efficacious antimicrobial, antiviral, anti-HIV, anti-cancer and anti-inflammatory (Satari, 1996). The compound is a secondary metabolite used in self-defense system, which is to preserve life and avoid interference from other organisms in the environment. Because farmakologiknya activity of the compound has the prospect to be isolated and used in medicine (Sardjoko, 1996)

Sponges are multicellular organisms invertebrates potentially be used as a search to discover new compounds for anticancer and antiviral sponge is producing cytotoxic compounds (Garson, 1994). Data from Schmitz 1998 in Setyowati, et al., 2007 states that of the 434 chemical structure of marine life that is cytotoxic sponge was ranked as the largest with 193 compounds, ascidian (57), algae (44), mollusks (46), soft corals (27), gorgonian (20), dinoflagella (8), anemones (8), echinoderms (7), worms (8), bryozoan (5), bacteria (3), and hydroids (3); while according Soest and Braekman (1999) the number of structures of compounds that have been obtained from a sea sponge until May 1998 is a 3500 type of compound, which was taken from the 475 types of two classes Calcarea and Demospongiae. The compounds are mostly taken from the Class Demospongiae especially of the order Dictyoceratida and Dendroceratida (1250 compound of 145 types), Haplosclerida (665 compounds of 85 species), and Halichondrida (650 compounds than 100 types). South Sulawesi waters, have been inventoried four types of sponges, namely: *Halichondria* sp., *Callyspongia* sp., *Callyspongiapseudoreticulatta*, and *Auletta* sp., Each of which has extract bioactive compounds are bactericidal. The test results of bacteria gave the same response to the type of commercial bactericidal, even at lower doses, namely: 20-40 ppm, it can inhibit the growth of bacteria (Ahmad et al., 1995).

Van Soest and Braekman (1999) found some of the family of bioactive compounds such petrosidaepolihidroksilat Oxycetylene, cyclic 3-alkilpiperidin, and siklopropenasterol. In addition, some active compounds have been discovered and reported from the genus Petrossian are alkaloids manzamine-A is cytotoxic (El sayed et al., 2001). At Petrossian sp. polyacetylene compounds found, dideoxypetrosynol A which showed antitumor activity in human

* Corresponding Author : **Asri Saleh**
Hasanuddin University,
Graduate Program Department of Chemistry.
Jl. Perintis Kemerdekaan Km. 10
Tamalanrea Makassar, South Sulawesi, Indonesia.
asri.dosen.uin@gmail.com

skin melanoma cells (Cho et al., 2004). Antibacterial activity was also found in the isolated from the marine sponge *Petrosiancontignata*, namely, Taraxeron and D-homoandrostan (Sutedja et al., 2005). Epidoksi sterol antibacterial compounds from marine sponges *nigransPetrosian* has also been isolated and characterized by a molecular formula C₂₉H₄₈O₃ with the name 5,8-epidioksi-24-etilkolest-6-en-3-ol (Handy et al., 2008).

Test of bacterial resistance to antibacterial agents has been done in Indonesia. From the information obtained that *E. coli* and *Salmonella* sp isolated from chickens, pigs, and other livestock are already many are resistant to antibacterial, not only against the antibacterial but also against some antibacterial (Sri Purnomo et al, 1992). Besides, *Staphylococcus* sp and *Streptococcus* sp, and resistance *Mycoplasma gallisepticum* isolated from cases of mastitis in cows, many multiresistant against some antibacterial.

Actually, the problem of bacterial resistance has been going on for a long time, just the attention to the effects of resistance has not been so great at this time. At this point to know the resistance is not only based on growth inhibition tests in the laboratory, but also marked by a change in the way of treatment, which was originally only with sulfonamides then need to more drug patents. The spread of resistance can occur genetic transfer. On the use of antibacterial which has the effect eradicate very quickly, antibacterial substances that leaves no residue can not create resistant bacteria. Resistance is the result of long-term use in very small amounts. In experiments that the mutant bacteria resistant to antibiotics, especially chloramphenicol, ampicillin, tetracycline, and ciprofloxacin.

Transitional season for most farmers is considered as the most worrying thing, where a decline in the body's physical condition of livestock, especially cattle were minimal antibody, so the response to any treatment on livestock decreased. The pattern of the disease was recorded as the cause of the biggest losses in livestock, such as respiratory and digestive diseases include snoring, defecation lime, cholera, and kolibasilosis (Infovet, 2006). Therefore, in this study will be conducted extraction of antimicrobial compounds such as chloroform fraction and the protein fraction of some types of sponges and analysis of antimicrobial activity against microbial pathogens from poultry.

Searching drug from the sponge in some Indonesian waters have been done, but there are still many locations in Indonesia that has not been touched (Wahyuono, 2003). Based on this potential, so in this study were taken in the waters around the island of Goods Lompo, because it has a diversity of sponge species are abundant (R Rahman, 2004).

RESEARCH METHODS

1. Isolation and purification of secondary metabolites from sponges

The method used in this study is based on a method often used in isolating the chemical compounds of natural materials which include the selection of species sponge, determining the location of sampling, preparation and sampling of animals, maceration,

partitioning, fractionation and spectroscopic analysis of pure compounds were obtained and continued to test the activity of the compound obtained (Soekamto, 2003).

2. Antibacterial Test

Screening is done to the antibacterial activity of *Escherichia coli*, *Staphylococcus aureus* and *Salmonella pullorum*, *PasteurellaMulcotida*. Basic selection because microbes are pathogenic microbes fourth is that for *Escherichia coli* and *Salmonella pullorum* a facultative anaerobic bacteria are pathogenic gram-negative primary cause of chronic diarrhea and typhoid. *Staphylococcus aureus* is a gram positive bacterial catalase positive cocci that are pathogenic cause skin infections and food poisoning, while *PasteurellaMulcotida* is the pathogenic cause of vaginitis or vaginal discharge (Hawley, 2003)

The medium used to grow the microbes are cultured Muller Hiliton Agar (MHA), medium noodles contain meat extract peptone as a source of protein and glucose as a source of carbohydrates which are known to support the growth of bacteria and fungi or also known as a versatile medium. Dilution test is used for solid nutrient agar medium Glucose (GNA) is a solid medium by solidifying such that (Hadioetomo, 1990). Chloramphenicol is selected as the positive control of the test for a broad spectrum of antibacterial activity that is effective for gram-positive bacteria and gram negative bacteria and other microorganisms (Mycek, 2001)

3. Structure Determination of Compound

The structural determination can be done by measuring instruments such as UV and IR against bioactive compounds that have been purified.

RESULT AND DISCUSSION

1. Extraction and Fractionation

Results maceration after filtered chloroform extract is evaporated at low pressure obtained maserat viscous brown residue such as 1280 mL and weight conversion pervolume obtained extract as much as 142.2 g. Results of liquid-liquid extraction in a separating funnel successively with solvent n-hexane, chloroform and ethyl acetate to evaporation using a rotary tool with a low-pressure vapor obtained n-hexane extract (10.45 g), chloroform (11.367 g).

Chloroform extract (11,367g) are then fractionated by using KKV and eluent n-hexane, a mixture of n-Hesan-ethyl acetate with increased polarity obtained 15 fractions. Based on TLC analysis of fractions with the same R_f combined to obtain 8 major fraction (A-I), then evaporated and the severity is determined and monitored by TLC.

Fraction 3, after fractionated by use of the summit eluent n-hexane, a mixture of n-hexane, ethyl acetate with increased polarity obtained 15 fractions.

Fraction 7 in the form of yellowish white powder, crystallized and recrystallized with methanol obtained compound (1) in the form of a white powder of 4.2 mg with a melting point of 118 -119 ° C. The purity of the compound through the TLC analysis showed single spot with three kinds of eluent system.

The main fraction to 3 after crystallized with methanol to produce the compound (2) in the form of white crystals as much as 11.2 mg. The crystals dissolve in the solvent n-hexane. 3 major fraction further fractionated by use of the summit with the eluent ethyl acetate-n-hexane 40% obtained five fractions fraksi. Setelahdikristalisai with acetone then 3 fractions and fractions were analyzed by TLC HX together, because the analysis has Nili TLC Rf together with a melting point of 187-189 ° C. Characters compound does not fluoresce under UV, but by using a stain reagent penampakseriumsulfat show stains blue first thenfaded and dissolved in chloroform. The purity of the compound (2) evidenced by TLC analysis with three kinds of systems eluent showed a single stain.

Table 1. Values of antibacterial activity of 8 major fraction of the chloroform extract fractionation results sponge Clathriareinwardhti

The main fraction	This type of bacteria			
	SA	SP	EC	PM
F1	-	-	-	-
F2	-	12,05	15,5	10,75
F3	-	-	11,15	-
F4	-	12,5	12,5	-
F5	-	18,9		
F6				
F7	12,2	-	11,6	-
F8	-	-	-	10,5
K +	16.6	13,7	22,1	21,9
K -				

Table 2. E. Coli inhibition zone in mm antibacterial compound (single isolates)

No	compounds	weight (Mg)	Zone barriers antibacterial (Mm)
1.	compound 1	4,2	9,51
2.	compound 2	11,2	11,41

3. Measurement Spectroscopy

Compound (1) was obtained as a white powder with a melting point of 118-119 ° C. UV (MeOH) max: 237 nm and 366 nm; the addition of NaOH reagent showed max: 237 nm and 366 nm; IR spectrum (Kbr) Vmax cm-1: > 3000 cm-1 (OH), 2918, 2962, 2850 cm-1 (CH aliphatic) 1705 cm-1 (C=O), 1261, cm-1 (O-CH3), 1097 cm-1 (CO), 1465 cm-1 and 1407 cm-1 (CH2 and CH3) and CH bending out the absorption field 865, 801 and 720 cm-1

Compound (2) dperoleh as white crystals with a melting point of 129-131 ° C. UV (MeOH) max: 229 nm and 274 nm; the addition of NaOH reagent slide show max: 229 nm and 274 nm; IR spectrum (Kbr) Vmax cm-1: 3433 cm-1 (OH), 2924 and 2851 cm-1 (CH aliphatic) 1107 (CO), 1710 cm-1 (C=O), 1464 and 1374 cm-1 (CH2 and CH3) and CH uptake bending out the absorption field 959, 879 and 793 cm-1.

DISCUSSIONS


1. Interpretation compound

Compound 1 was obtained in the form of white powder with a melting point of 118-119 ° C. Qualitative test results with Liebermann Burchard reagent showed positive purple red color indicating the compound triterpenoids.

Of the UV spectrum appears that compound 1 provides an absorption band maximum in the wavelength range λ_{maks} 237 nm (9230) and the absorption at a wavelength of λ_{maks} 366 nm (727), after the addition of reagent sliding NaOH does not cause a wavelength shift that indicates that there is no shift force hydroxyl.

IR spectrum of the data mentioned above, it appears their absorption at maks > 3000 cm-1 indicate the presence of OH groups, absorption at 2918, 2962, 2850 cm-1 were very strong and sharp indicates aliphatic CH group followed by absorption at maks 1463 cm-1 which is an aliphatic CH bending of CH2 and maks absorption at 1385 cm-1 which is an aliphatic CH bending of CH3 typical for the class of triterpenoids (Yoshihiro et al, 2001). Absorption at 1705 cm-1 which shows the strain stretching C=O bond as ketones, cyclic absorption at 1261 cm-1 indicates the presence of the methoxy and absorption at 1097 cm-1 is a strain stalling of CO secondary alcohol and bending out the field of CH on the uptake 865, 801 and 720 cm-1.

Fig. 1. Molecular structure of Triterpenoid Carbocsilat acid



Based on the data above and the results of literature studies triterpenoid compounds it can be concluded that compound 1 is a compound triterpene class.

Compound 2 was obtained in the form of white crystals with a melting point of 129- 131 oC. The character of this compound does not fluoresce under UV, but by using a stain reagent penampakseriumsulfat show stains blue first then fade and dissolve in chloroform. Qualitative test results with Liebermann Burchard reagent produce blue-green color indicating the steroid compound it is also supported by the UV and IR spectral analysis.

Further information about the compound 2 as a steroid compound obtained from infrared spectrum appear to have the maximum wave number in the area maks 3433 cm⁻¹ which is the absorption of the OH group (hydroxyl), an indication of the presence of hydroxyl groups supported by local absorption at 1107 cm maks -1 is stalling strain of CO secondary alcohol which is typical for steroids (Guogiang et al, 2005). In maks wave numbers 2924, 2851 cm⁻¹ absorption are very strong and sharp indicates aliphatic CH maks followed by absorption at 1464 cm⁻¹ which is an aliphatic CH bending of CH₂ and maks absorption at 1374 cm⁻¹ an aliphatic CH bending of CH₃. maks wave number at 1710 cm⁻¹ shows the absorption carbonyl (C = O) as cyclic ketones and the number of wave upon wave maks 1259 cm⁻¹ robust to the methoxy and bending out of plane C-H on the uptake 959.879 and 793 cm⁻¹.




Fig. 2. The molecular structure of steroid compounds (-sitosterol)

Based on the data above and the results of the study of literature steroid compounds it can be concluded that the compound 2 is a steroid compound.

2. Test Compounds bioactivity Secondary Metabolites

Secondary metabolite n-hexane extract, fractions and single isolates obtained from sponge *Clathriareinwardhti*. tested their activity using pathogenic bacteria in poultry. The test results demonstrated toxicity is high enough even no toxicity is considered very high.

Activities extract beginning of chloroform, against pathogenic bacteria in poultry These results indicate that chloroform extract is very active, indicating that fractions of chloroform extract sponge *Clathriareinwardhti* may contain compounds that are bioactive, or there may be some compounds that are inactive are joined and each other strengthen bioactivity causing the active fractions

Fraction 3, Fraction 7 test the bioactivity although not completely soluble in the test solution used in this case is DMSO Then 3 fractions, and fractions 7 whose activity is exceptionally high 10.9 mm after direksistalisasi obtained compound (2) were confirmed as group compound steroid showed that this fraction contained compounds that are bioactive. This is supported by the discovery of the steroid compound in the fraction. Class of steroid compounds that can almost always be found in animals and plants. These compounds are thought to form acetic acid via the mevalonate acid pathway then experience

Some of the condensation reaction, cyclization and forth to form an intermediate / intermediate. The use of pharmacologically active compounds derived from nature such as steroid derivative is very important in terms of health because of its side effects is relatively small compared with synthetic compounds. In addition, raw material compounds can also be updated. Steroid compounds such as -sitosterol has a pharmacological effect that is capable of inhibiting the action of an enzyme that converts testosterone to dehidrotestosteron (DHT) which is a cause of prostate cancer (RenaiSante in Sapar, 2004).

CONCLUSION

The interpretation of physical data and spectra (UV and IR) produces two types of compounds obtained are (1) compounds triterpenoids and (2) steroid compound. Bioactive test results conducted on pathogenic bacteria in poultry shows that the compound (1) is very toxic to pathogenic bacteria *E. coli* in poultry was 9.51 mm compound (2) is quite toxic to pathogenic bacteria in poultry 11.41 mm

SUGGESTION

Clathriareinwardhti potential to be developed as fitofarmaka considering the compounds contained therein are bioactive. It is necessary for further exploration and analysis of more spectrum in order to know for sure secar structure of the compounds contained therein

BIBLIOGRAPHY

1. Achmad, S.A. 2004. Forty years in the organic chemistry of natural materials tropical vegetation Indonesia, recollection and Prospects. Bulletin of the Indonesian Society of Natural Products Chemistry, 4 (2): 5-54.
2. Anderson, J.E., Goetz, C.M, and McLaughlin, J.L. 1990. A Comparison of Simple Banch blind-top Bioassay and Human tumor as Anti-tumor Cell Cytotoxicities prescreen. *Phytochemical Analysis*. 6: 107-111
3. Anonymous, 2003. Foundation Scuba Diver Indonesia. <http://www.Terangi.or.id/Indonesian/reef-Indon>, accessed in 2008.
4. Anonymous 2006, for Panacea Sea. <http://www.Forek.or.id>, accessed in 2008.
5. Amir, I & Bidiyanto, A., 1996, Know Your Marine Sponges (Demosponge) Sec. General. Oseana, Vol 21, No 2, Lipi, Jakarta.
6. Astuti, P., 2003, Invertebrates Marine Sponges Potential as a Source of Raw Materials Natural Medicine, vol 826 October to December (Special Edition). Part-Pharmaceutical Biology, UGM, Yogyakarta.
7. Barnes, R., P. Candidate and P. Olive, 1989. *The Invertebrates*. Blacwell Scientific Pub. Oxford. London. Edinburg. Boston Melborne. Five Pub: 49-53.

8. Barnes, R.S.K., 1999. A New Synthesis. Second Edition. Blacwel Science, UK, 49-52.
9. Na-, G.B., Lazaro, J.E., Concepcio, G.P., 1994, Biological Assays for Screening of Marine Samples, Second Marine Natural Product Workshop, Marine Science Institute and Institute of Chemistry, University of the Philipines.
10. Dini, I. 2005. Secondary Metabolites search Plant Stem Extract Skin Paliasa (*Kleinhoviahospita* Linn.) And bioactivity against Artemiasalina Leach. Unpublished thesis. Makassar: Department of Chemistry, Graduate Program, University of Hasanuddin.
11. EruWibowo, A., et al., 2005, Study of the Secondary Metabolites Compound Exploration of the Sea biota, Center for the Study and application of Pharmacy and Medical Technology. [http: / www. lptek.com](http://www.lptek.com), accessed in 2008.
12. Fusetani, N., J Warabi, Nogata K., Y., Nakao, Y & Matsunaga, S. 1997. Koshikamide A1, a new Cytotoxic Linear Peptide Isolated from a Marine Sponge, *Theonella* sp. *Tetrahedron letters* 40, 4687-4690.
13. Garson, M.J., 1994. The Biosynthesis of Sponge Secondary metabolites: Why it is Important? In: Soest, R. W. M. van, Th. M.G. van Kempen and J. C. Braekman, Sponges in Time and space. Proc. 4 th Int. Sponge congr: Rotterdam: Balkema.
14. Billy, D. 1984. Malignant Tumor Chemotherapy in "Malignant Tumors in Children". Section of Anatomic Pathology, Faculty of Medicine University of Indonesia, Jakarta, 99-105.
15. Gan, S. 1987. Anti-Cancer in "Pharmacology and Therapeutics", Edition III, Section of Pharmacology FK-UI, Jakarta. 625-626, 635 ..
16. Gisela P.C., Gina, C., and Lazaro, J.E. 1994. Biological Essay For Screening Of Marine Samples, "In Natural Product Workshop", Work Book, Marine Science Institute, University Of The Philipines, Philippine, 15-18.
17. Guogiang Li Zhiwei, D., Huasi, G., van Leen, O., Peter, P & Wenhan, L. 2005. *S t e r o i d s f r o m s o f t c o r a l D e n d r o p h y t a s p .* www.elsevier.com/locate/steroids. Accessed 2008.
18. Hadi, S. and John B. Bremner. 2001. Initial Studies on alkaloids from Lombok Medicinal Plants: Molecules. Department of Chemistry; University of Wollongong; 6. 117-129 V. Wollongong, Australia.
19. Harryanto, A.R., Aru, W.S., 1990. Kemoterpi Cancer in Medicine. Volume II, Hall Publishers FK-UI, Jakarta, 524-529.
20. Hooper, J.N, A., 1997. Guide to Sponge Collection and Identification. Version Merch. Queensland Museum South Brisbane, Queensland.
21. Hostettmann, K., Hostettmann, M., Marston, A. 1991. Isolation and Cytotoxic Compounds Test Nature Materials. ITB, Bandung.
22. Jasin, M., 1987. Systematics of Animals Invertebrates and Vertebrates. SinarWijaya. Surabaya
23. Compass, 2004. Digging Drug Raw Materials in Under the Sea. Publication May 12, 2004. Jakarta.
24. Lomis, FY, 1978. Basic Toxicology. Edition III, Translators Imono Argo, Semarang Teachers' Training College Press, 4, 16-21. Manuel J. and Luis M., 2001. Plant Natural Product With Leishmaniacid Activity. J. The Royal Society of Chemistry, 18: 674-688. McLaughlin, J, L., C.J. Chang, and D.L. Smith, 1991. Benctop: Bioassay for The Discovery of Bioactive Natural Products an update; in studies in Natural Products Chemistry. Elsevier, Amsterdam, in Press. 1-10.
25. Meyer, B.N., N.R. Ferrigni, J.E Putnan, L.B. Jacobsen, D.E. Nicholas, J.L. McLaughlin 1982. Brine Shrimp: A Convenient General Bioassays for Active Plant Constituent. Department of Medical Chemistry and Pharmakognocny, School of Pharmacy and Pharmacal science, and Cell Culture Libratory, Perdue Cancer Center. West lavayette. USA.
26. Rafliizar, Adimunca, C., Tuminah, S. 2006. Paliasa leaf decoction (*Kleinhoviahospita* Linn) For Acute Liver Inflammation Medicine. *World Mirror of Medicine*. 50: 10-14.
27. Rahmaniar, 2003. Marine Natural Products as Lead Compound for Pharmaceutical and Agricultural Perspective Presented at one day seminar recently in Drug. Discovery, Makassar 2008.
28. Rahman, R, & Ahmad Abdridhay, 2004. Screening of Antimicrobial compounds of Several Types of Sponges (Sponge) .Tesis not diterbitkan. Makassar, Department of Pharmacy Univeritas Hasanuddin.
29. Romimohtarto, K & Sri Juwana, 2001. Marine Biology, Marine Life Science, Djambatan, Jakarta.
30. Rosmiati & Suryati, E, 2001, Isolation, Identification and Effect of bioactive compounds against Bacterial Pathogens *Spona* shrimp. <http://Pustaka.bogor.net/publ/J/biotech/diakses2008>.
31. Rusli, 2005. Isolation and Identification of Active Compounds Anti Microbial Some of Aquatic Sponges Samalona Island. Unpublished thesis. Makassar: Department of Chemistry, Graduate Program, University of Hasanuddin.
32. Sapar, A., U.S. Kumanireng, N, de Voogd, Alfian N, 2004. Isolation and Structure Determination of Active Secondary Metabolites Sponges *Biemnatriraphis* Origin Kapodasang Island (Islands Spermonde), *Marina Chemica Acta*. J.V. 6 NO.1.
33. Sarjoko, 1996. Quantitative Structure Activity Relationships, Design Rational Bioactive Compound Development, Presented on a one-day seminar in a new perspective Drug. Discovery, Ujung Pandang
34. Satari, RR, 1999. Marine Products Research Natural Products in Indonesia. Directions and prospects: National Seminar on Natural Materials Chemistry. Jakarta.
35. Scheuer, P.J., (Ed), 1978. Marine Natural Products: Chemical and Biological Perspectives. Vol. II. Academic Press, Inc. New York. USA.
36. Schmidt, E.W and Faulkner, D.J. 1998. Mecrosclerodermis C-E, Anrifungal Cyclic, peptide from lithistid Marine Sponges *Theonella* sp and *spMicroscleroderma*, *Tetrahedron* 54, 3043-3056.
37. Soekamto, N.H., 2003. Profile Phytochemicals Some species *Moraceae* Indonesia; Unpublished dissertation. Institut Teknologi Bandung, Bandung.
38. Troter II, R.T., Logan, M.H., Rocha, J.M., and Bonetta, J.L., 1983. Ethnography and Bioassay: Combined Methods for a Preliminary Screen of Home Remedies for Potential Pharmacological Activity. *MFI, J. of Pharm*, vol 6, no 4.
39. Ulfa, M. 2006. Isolation, Characterization and Test bioactivity Secondary Metabolites Plant Stem Extract Skin Paliasa (*Kleinhoviahospita* Linn.). Unpublished thesis. Makassar: Department of Chemistry, Graduate Program, University of Hasanuddin.
40. Wahyuono, S., 2003, for anticancer drug of Sponge Indonesian waters, *Horizon Supplements Mind*. [http: / www. Rakyat.com mind](http://www.Rakyat.com/mind), accessed in 2008.
41. Wiryowidagdo, S & W. Moka, 1995. Identification and Exploration of Marine Organisms as a Source of Raw Material drug in Spermonde Islands Sul-Sel.
42. Yuliani, S. 2001. Prospects for Development of Traditional Medicine Drug Being Fitofarmaka. *Journal of Agricultural Research Research Institute for Spices and Medicinal Plants - Bogor*. 20 (3).
43. Yoshihiro, M., Masoto, F., Akihito, Y., Yutaka, S., Shigenori, and Hiroshi, S. 2001. triterpene glycosides from the roots of *Sanguisorba officinalis*. www.elsevier.com/locate/phytochem. Dieakses 2008.