

Identification, abundance and diversity of forage plants for Anoa (*Bubalus* spp.) in Abdul Latief Forest Park

by Aziz Aziz

Submission date: 04-Jan-2023 08:12PM (UTC+0700)

Submission ID: 1988506831

File name: 1-s2.0-S1872203222000737-main.pdf (1.31M)

Word count: 6191

Character count: 32707



Identification, abundance and diversity of forage plants for Anoa (*Bubalus* spp.) in Abdul Latief Forest Park

Isna Rasdianah Aziz^{*}, Hasyimuddin, Ade Muspa, Masriyan, Zulkarnain, Muhammad Khalifah Mustami

Department of Biology, Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Gowa, South Sulawesi 92113, Indonesia

ARTICLE INFO

Keywords:

Checkered line
Elatostema integrifolium (D. Don) Wedd.
 Forage plant
 In-situ and ex-situ conservation
 South Sulawesi

ABSTRACT

Anoa (*Bubalus* spp.) is a protected endemic species on the Sulawesi Island. An assessment of its habitat is needed to maintain the conservation, especially for the forage plants consume. Therefore, this study aims to determine the types, abundance, and diversity of the available forage plants for Anoa in the Abdul Latief Forest Park. The study was carried out using checkered lines with a descriptive approach in three blocks, including station I as protection block, station II as utilization block, and station III as collection block. The checkered line method contains three repetitions in three blocks with a total area of 1.8 ha. Data analysis was carried out using the Shannon-Wiener variability formula, Margalef wealth index, Evenness abundance, and dominance index. Based on the results, six plants were identified as Anoa feed, including *Ficus racemosa* L., *Elatostema integrifolium* (D. Don) Wedd., *Nephrolepis* sp., *Begonia* sp., *Scleria multifoliata* Boeck., and *Alpinia* sp. Furthermore, the leaf is the most consumed part of the plant by Anoa, and the *E. integrifolium* (D. Don) Wedd. is the forage plant with the highest abundance value of 49,167 (ind ha⁻¹) in the protection block. Meanwhile, for the diversity value, each block was in a low category at 0.88, 1.28, and 1.44, respectively. Based on the index value of species wealth, the highest value was observed in the collection block; hence more species were discovered in this block than the other two blocks. However, *Elatostema integrifolium* (D. Don) Wedd. showed a dominance index in the protection block. This study affects the conservation of Anoa in the area to be continued by observing the biodiversity utilization, which is too burdensome without regard to sustainability.

1. Introduction

Anoa (*Bubalus* spp.) is an endemic ungulate mammal rare on the Sulawesi Island, including animals protected by wildlife protection laws in Indonesia [1,2], also classified as endangered species according to IUCN red list [3,4]. However, their population and habitat are decreasing with limited distribution. This decrease in population is evidenced by the difficulty locating these animals randomly in their habitat [5,6]. The second threat to Anoa after poaching is the degradation and loss of habitat [7] because most of these animals are isolated when forests in protected areas are cut down or converted to plantation land [8,9].

Anoa habitat's existence affects feed availability, which is a crucial component in animal management both in captivity and naturally [10]. The quality or quantity of feed, including nutrient content, preferences, appetite, and feeding behavior of Anoa [11,12] is an essential limiting

factor in captivity, survival, and reproduction [10,12]. A primary feed for Anoa consists of coarse forage particles [13,14] rich in crude protein, fiber, energy, and minerals, making their internal rumen structure more efficient in the digesting process phylogenetic and anatomic affiliation to either *Bubalus bubalis* L. [12,15]. In wild habitat, Anoa commonly consumes herbs (*Physalis minima* L.) [16], *Elatostema* sp., *Impatiens* sp. [17]), flowering plant (*Schizostachyum* spp., *Trema orientalis* (L.) Blume [12,16]), liana (*Merremia peltata* (L.) Merr. [12,16]), fern (*Microlepia* spp. [12], *Angiopteris evecta* (G.Forst.) Hoffm. [17], *Scleria purpurascens* Steud. [18]), grass (*Scleria* spp. [12], *Brachiaria mutica* (Forssk.) Stapf, *Panicum* sp. [18]), reed (*Imperata cylindrica* (L.) P.Beauv. [18]), leaves of shrubs (*Acalypha boehmerioides* Miq. [16]), young leaves/shoots (*Micromelum minutum* (G.Forst.) Wight & Arn., *Sonneratia alba* Sm. [16], *Athyrium nigripes* (Blume) T.Moore, *Elatostema* sp., *Pinanga* sp., *Zyzygium* sp., *Calamus* sp., *Crypteronia griffithii* C.B.Clarke, *Cratogeomys celebicum* Blume, *Areca vestiaria* Giseke, *Impatiens* sp. [17], *Ficus vasculosa* Wall. ex

^{*} Corresponding author at: Department of Biology, Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar, Jl. H.M. Yasin Limpo 36 Romangpolong, Gowa, South Sulawesi 92113, Indonesia.

E-mail address: isna-rasdianah@uin-alauddin.ac.id (I.R. Aziz).

<https://doi.org/10.1016/j.chnaes.2022.09.008>

Received 13 March 2021; Received in revised form 25 September 2022; Accepted 26 September 2022

Available online 4 October 2022

1872-2032/© 2022 Ecological Society of China. Published by Elsevier B.V. All rights reserved.

Miq. [18]), and fruit (*Diospyros pilosanthera* Blanco, *Eugenia* sp., *Ficus vasculosa* Wall. ex Miq. [16,18], *Dillenia serrata* Thunb., *Zyzygium* sp., *Crypteronia griffithii* C.B.Clarke, *Cratogeomys celebicum* Blume [17], and *Musa* sp. [18]).

Primary and secondary forest in Sulawesi Island supply various fruit-producing trees which provided food for animals throughout the year [19]. However, the Ministry of Environment and Forestry Republic Indonesia stated that deforestation in Sulawesi reached 13.9% from 2017 to 2018 [20]. The limited existence of natural forest as Anoa habitat has an impact on the Anoa population pressure [21]. An area in South Sulawesi which acts as Anoa habitat is the Abdul Latief Forest Park at Sinjai Regency, which has not been widely studied.

Abdul Latief Forest Park is included in the conservation area with an area \pm 720 ha and an altitude of 1200–2000 masl. It is divided into several blocks including utilization, collection, protection, rehabilitation and traditional blocks [22]. Furthermore, in this area, Anoa has become an icon of conserving unique fauna [23]. Several efforts which were previously carried out in order to preserve these animals need to be improved by looking at various factors, especially in their habitat. In in-situ and ex-situ, which each have benefits and drawbacks or integration of both component, a solution may be by paying attention to the addition or regulation of diet and feed availability [5,24–26]. This proves that knowledge on the food for Anoa is crucial in preserving these animals and ensuring that they adapt successfully to the conservation process. This study aimed to determine the types, abundance, and diversity of Anoa forage plants. It provides support for in-situ and ex-situ Anoa as well as conservation efforts in South Sulawesi, particularly in Abdul Latief Forest Park.

2. Materials and methods

2.1. Study area

This study was carried out in August–September 2020 (dry season), in Abdul Latief Forest Park with an altitude between 1200 and 2000 masl. In addition, it was divided into five blocks, such as utilization, collection, protection, rehabilitation and traditional blocks. Based on direct observations and information from the local government officials, three blocks were suspected to be locations frequently visited by Anoa (*Bubalus* spp.) include utilization (1346–1430 masl), collection (1514–1570 masl) and protection (1535–1598 masl) (Fig. 1). Furthermore, sample identification and analysis were conducted at the Laboratory of Ecology and Laboratory of Genetics and Molecular, Department of Biology, Faculty of Science and Technology, Universitas Islam Negeri Alauddin Makassar.

2.2. Sample collection

Sampling was conducted using the line plot method with a stub line length of 100 m after determining the station. Observation of the feed type for Anoa was carried out on square plots with various sizes such as respectively 20 m \times 20 m, 10 m \times 10 m, 5 m \times 5 m and 2 m \times 2 m for tree, poles, saplings and seedlings placed on the right and left of the stub line, respectively. Each block was drawn 3 times on the 100 m stub line [27]. Data such as plant species, number of individuals for each species, diameter for the tree and pole levels, including plant parts consumed by Anoa were recorded. Furthermore, several samples were identified on




Fig. 1. Map of the observation station in the Abdul Latief Forest Park: Station I protection block (red dot); station II utilization block (yellow dot); station III collection block (blue dot).

the spot and in the Laboratory. Labeled herbarium specimens were made entirely for taxonomic purposes.

2.3. Data analysis

Data obtained were analyzed descriptively and displayed in tables and graphs. The name of plant species was identified based on taxonomic literature [28–30] and results of some previous study related to the type of feed consumed by Anoa [17,31–34]. Furthermore, data obtained in the field is analyzed to calculate the importance value index (IVI) obtained from the sum of relative density (RDe), relative frequency (RF) and relative dominance (RD). In addition, it is used to calculate diversity index (H'), wealth index (DMg), evenness index (J'), dominance index (C) and abundance index. Each value is obtained using the following formula [34]:

$$IVI = RDe + RF + RD$$

RDe states as the ratio of forage plant density in the plot, RF shows the proportion of total forage plants appears, and RD states as the proportion of the total basal area of all forage plants present. Species diversity index (H') was calculated as follows [35]:

$$H' = - \sum P_i \ln P_i$$

The Shannon-Wiener diversity index has indicators such as low, moderate, and high diversities in the range of $H' < 1.5$, $1.5 \leq H' \leq 3.5$ and $H' > 3.5$, respectively. Margalef wealth index was calculated as follows [35]:

$$DMg = \frac{S - 1}{\ln N}$$

The margalef wealth index has the following indicators: $DMg < 3.5$, $3.5 < DMg < 5$ and $DMg > 5$ classified as low, medium, and high types, respectively. Calculation of evenness index is shown as follows [35]:

$$J' = \frac{H'}{D_{max}}$$

The evenness index has the following indicators, namely at values of 0.00–0.25, 0.26–0.50, 0.51–0.75, 0.76–0.95 and 0.96–1.00, classified as uneven, less evenly, quite evenly, almost even and evenly distributed, respectively. The dominance index has the following indicators namely at a value of $0 < C < 0.5$, no species dominate and at $0.5 < C < 1$, there is a dominating species. The calculation as follows [35]:

$$C = \sum \left(\frac{n_i}{N} \right)^2$$

According to Pujaningsih [36], qualitative abundance of forage plants has indicators at values of 1–4000, 4001–16,000, 16,001–30,000 and $> 30,000$ (ind ha⁻¹) are classified into rare, occasional, frequent, and abundant categories, respectively.

3. Results

3.1. The forage plants and fruit for Anoa (Bubalus spp.)

Protection, utilization, and collection block, which are observation locations in Abdul Latif Forest Park, has their own characteristics. In this study, six plants were discovered to be included in Anoa feed. These species consist of six different families, presented in Table 1.

Elatostema integrifolium (D. Don) Wedd. identified in three blocks was having the largest number, consisting of 493 individuals. Otherwise, *Begonia* sp. was only found in collection block. In line with Broto [37] that identified the same species as abundance and dominant cover crop for Anoa habitat in Mekongga mountains protected forest, about 260 km from Abdul Latief forest park. *Elatostema* sp. also has higher abundance among 28 plants feed for Anoa in Bogani Nani Wartabone National Park [27], about 771 km from Abdul Latief forest park.

Based on the parts most consumed in the plant, Anoa choose the preferred three part of six plants such leaves, shoot and fruit. But the leaves dominate it. Abdul Latief Forest Park is covered by various types of mountainous/upland plants which are suitable with the Anoa habitat dominated by woody, medicinal plants and some lower-level vegetation [23]. This habitat provides plants with abundant leaves that yearly availability in large amounts. Compared to *Ficus racemosa* L. which produces fruit, Anoa prefers to consume herbs which provide more leaves and shoots. Although *Ficus* sp. will bear fruit throughout the year in the rainy and dry season, the species at three blocks is entirely 19 individuals, which is mostly scattered in station II.

Furthermore, some signs of Anoa were discovered in Abdul Latief Forest Park with ripped or torn surfaces marks in several feed plants including *Alpinia* sp. and *E. integrifolium* (D. Don) Wedd., also feeds from Anoa as shown in Fig. 2.

The importance value index (IVI) is a reference to discover and determine the dominant species in a community. Therefore, the species with a high IVI value indicates that these feed plants have wider power areas compared to other plants as presented in Table 2.

Ficus racemosa L. is the only tree-level vegetation found in the whole plant community, showing the highest dominance guided IVI of 1.2. The herb and seedling community with five species have the values for dominance directed IVI of 0.9, 0.4, 0.31, 0.2, and 0.04, respectively. The lowest value of IVI is recorded by *Begonia* sp. among all species, representing the lowest heterogeneity and species diversity because it only grows in one block.

3.2. The abundance of forage plants

The availability of a feed is seen from its abundance which is influenced by environmental conditions and the growth characteristics. Graph 1 shows the abundance of Anoa feed from each block.

E. integrifolium (D. Don) Wedd. has the highest abundance among the six plant species in protection, utilization, and collection blocks with values of 49,167, 16,833, and 16,167 ind ha⁻¹, respectively. Three species in St.I, St.II, and St.III are categorized as rare species, including *Begonia* sp., *Ficus racemosa* L., and *Alpinia* sp. Then *Scleria multifoliata* Boeck. and *Nephrolepis* sp. are categorized as occasional species, while

Table 1
Plant species for Anoa feed in Abdul Latief Forest Park, Sinjai Regency.

Family	Species	Local Name (Sulawesi)	N	The consumed part	Habitus	Location
Moraceae	<i>Ficus racemosa</i> L.	Karoti; bilajeng; lambere	19	F	T	St.I, St.II, St.III
Begoniaceae	<i>Begonia</i> sp.	Lariang	18	L	H	St.III
Poaceae	<i>Scleria multifoliata</i> Boeck.	Rumput piso	134	L	H	St.I, St.II, St.III
Utriacaeae	<i>Elatostema integrifolium</i> (D. Don) Wedd.	Katiloporo	493	L	H	St.I, St.II, St.III
Zingiberacea	<i>Alpinia</i> sp.	Kunyi' hutan; unnyi	44	S, L	H	St.I, St.II, St.III
Lomariopsidaceae	<i>Nephrolepis</i> sp.	Paku yanre; paku pabbura	155	L	H	St.I, St.II, St.III

Note: N = total of individu; F = fruits; L = leaves; S = shoots; T = tree; H = herbs; St.I = station I protection block; St.II = station II utilization block; St.III = station III collection block.



Fig. 2. The traces of Anoa (*Bubalus* spp.), A. Feces, B. Ripped leaves are shown in *Elatostema integrifolium* (D. Don) Wedd., C. Ripped/torn leaves are shown in *Alpinia* sp.

Table 2
Importance value index of Anoa (*Bubalus* spp.) feed plant vegetation in Abdul Latief Forest Park.

Family	Species	Level of Growth	RDe (%)	RF (%)	RD (%)	IVI (%)
Moraceae	<i>Ficus racemosa</i> L.	Tree	0.0002	0.2	1	1.2
Utricaceae	<i>Elatostema integrifolium</i> (D. Don) Wedd.	Herb & Seedling	0.6	0.4	–	0.9
Lomariopsidaceae	<i>Nephrolepis</i> sp.	Herb & Seedling	0.2	0.2	–	0.4
Poaceae	<i>Scleria multifoliata</i> Boeck.	Herb & Seedling	0.16	0.15	–	0.31
Zingiberacea	<i>Alpinia</i> sp.	Herb & Seedling	0.1	0.1	–	0.2
Begoniaceae	<i>Begonia</i> sp.	Herb & Seedling	0.02	0.01	–	0.04

Note: RDe = relative density; RF = relative frequency; RD = relative dominance; Level of growth [38].

E. integrifolium (D. Don) Wedd. is categorized as frequent species in St. II and St.III. Lastly, only *E. integrifolium* (D. Don) Wedd. among all blocks is categorized as abundant species.


3.3. The diversity of forage plants

The diversity index shows the diversity of species in a particular area. Values in each category may affect Anoa's survival because this feed will be a necessity to survive. Based on Table 3, the three blocks are classified as a low category in diversity index value, which indicates that the community is only composed of a few species. However, station III as collection block has a higher value among the three blocks, which is influenced by a more open land cover by looking at the light intensity value compared to each block.

According to the Margalef wealth index, the forage plant richness indicates that the utilization and protection block is in the medium category. In contrast, the collection block is classified as a high type due to the highest number of species, where bioecological patterns and conditions play a role in allowing species to proliferate. The evenness index indicates species relatively evenly distributed in the protection block, while the utilization and collection blocks are classified as almost evenly species distributed categories, which illustrates the evenness

Table 3
Index value of Anoa forage plants in Abdul Latief Forest Park.

Location	Shannon-Wiener diversity index (H')	Margalef wealth index (DMg)	Evenness index (J)	Dominance index (C)
St.I	0.88	4.83	0.55	0.6
St.II	1.28	4.82	0.79	0.3
St.III	1.44	5.82	0.81	0.3



Graph 1. The abundance of Anoa forage plants in three stations of Abdul Latief Forest Park. St.I = station I protection block; St.II = station II utilization block; St.III = station III collection block.

balance of the six species between species communities. Furthermore, the protection block's dominance index is classified into the high dominant species category by *E. integrifolium* (D. Don) Wedd., due to the topography structure in accordance with species reproduction.

4. Discussion

In general, Abdul Latief Forest Park, located in Sinjai district, about 134 km from Makassar, the capital of South Sulawesi Island, is categorized as a conservation area with endemic flora and fauna. Consisting of several blocks includes utilization, collection, protection, rehabilitation, and traditional blocks as the government's prototype conservation initiative. We limited ourselves to researching the three blocks due to the pandemic condition.

The forage plants as Anoa (*Bubalus* spp.) feed that we identified in three blocks consisted of flowering plants (*Ficus racemosa* L., *Elatostema integrifolium* (D. Don) Wedd., *Scleria multifoliata* Boeck., *Alpinia* sp., *Begonia* sp.) and a sword fern (*Nephrolepis* sp.) (Table 1). *Nephrolepis* sp. is identified as a forage plant with the second-largest individual after *E. integrifolium* (D. Don) Wedd., propagate on the detritus, covered with a kidney-shaped sorus [39], and attach as epiphytes to the trees. Anoa consumes this species leaves as the preferred and staple food for survival adaptation in Abdul Latief Forest Park. This study is in line with previous study by Pujaningsih et al. [40] who found *Nephrolepis* sp. as delicate feed for Anoa in National Lore Lindu Park, although it was not abundant. The presence of this tree fern was also investigated by Bowling et al. [41] who found *N. biserrata* (Sw.) Schott and *N. hirsutula* (G. Forst.) C. Presl growth limited to ultrabasic soils Lambusango forest Buton, which indicate the Anoa abundance in that site. *N. acutifolia* (Desv.) Christ was also identified by Tangkoro et al. [42] as one of the forage plants for Anoa feed with high IVI value among the herb and seedling community in Tadulako University Education Forest. The same species found in the three blocks are the key and main forage plants as Anoa feed.

Anoa classified as a ruminant animal, selects and prefers to consume leaves, shoots, fruit, herbs, marsh, grass, and aquatic plants. Young leaves and shoots contain crude fiber at 8.4–62.68% [16,43], protein content at 0.6–21.60 [16,43], tenderness texture, and easily digested. Therefore, they are more popular with Anoa over the tough, old and woody feed. Of the six forage plants consumed by Anoa at the study sites, the most selected part was the young leaf. Apart from the leaf morphology's thin and soft texture, Anoa is considered to choose their food containing low tannin composition, including in the leaves. It will facilitate the digestion process of crude fiber, assisted by bacteria's activity in the Anoa's rumen. According to the phytochemical screening result of previous studies, tannins were not strongly present in leaves of *N. biserrata* (Sw.) Schott [43–46] and *N. hirsutula* (G. Forst.) C. Presl [47]. *A. galanga* (L.) Willd., *A. nigra* (Gaertn.) Burt, and *A. eremochlamys* K. Schum. leaves revealed the presence of tannins, but not strongly present as other secondary metabolites [47–49]. *A. rubricaulis* K. Schum. showed the absence of tannin in leaves [50]. *S. biflora* Roxb., *S. laevis* Willd., *S. polycarpa* Boeckeler, *S. pubescens* Steud. [51] also showed the absence of tannin in leaves. *F. racemosa* L. becomes the only plant in the three blocks whose fruit is favored by Anoa. This fruit consists of protein, crude fiber, various chemical constituents, and 82.3% water content, which can be an alternative to meet the requirement of the water needs. Besides, *Ficus* sp. does not contain tannins [51–53], except for the leaf and stem bark. Or it has a low tannin content ranging from 0.34% [54], 1.9–2.1% [55]. In order for Anoa to acquire the optimal level of food consumption to obtain the required nutrients. Anoa ate falling fruit directly from the tree in this forest park, or more fruit drop may occur by stabbing the *Ficus* stem. Furthermore, we found a *Ficus* sp. which has a shorter height than *F. racemosa* L., thus, the fruit is reachable close to Anoa. We have particularly limited our research area in this study due to the pandemic situation; hence it will be our next concern.

The presence of feces and ripped leave of the forage plant indicate the signs of Anoa in this forest park (Fig. 2). Ripped or torn leaves and

falling fruit were found not too far from the Anoa feces, <1 m. The Anoa feeding behavior, which is different from deer and squirrels in the site, conducts through smelling the food, sticking out the tongue to the edge of the leaf, wraps around the leaf, then stomps the head to pull and grab the leaf. These results are indicative of the evidence of Anoa abundance with was reported by Wardah et al. [21], who found Anoa feces in Pangi Binangga Nature Reserve, Central Sulawesi. With an area slope > 25%, it provides quite a lot of forage plants.

F. racemosa L. on tree-level vegetation has an advantage in the relative cover, which causes this plant to dominate and have extensive territorial power in all three blocks (Table 2). At the herb and seedling level, the high IVI value in *E. integrifolium* (D. Don) Wedd. indicates that it has a wide distribution area to sustain life, develop and expand its territory to obtain energy on the forest floor in Abdul Latief Forest Park. The protection block which is supported by highest humidity rate of 83–93% compared to the other three blocks had the highest abundance value in *E. integrifolium* (D. Don) Wedd. (Graph 1). It thrived in moist conditions, commonly near water bodies and traversed by streams of tributaries as a permanent water source. Hence it can spread rapidly on the forest floor of Abdul Latief Forest Park. The humidity support plant growth through cell extension. In addition, with a temperature range of 16–23 °C, protection block is a habitat that far from human reach and activities. Therefore, it is chosen by Anoa as an effort to defend themselves. According to local government [23], the protection block in Abdul Latief Forest Park is a site for protecting animal or plant species allocated to maintain tropical biodiversity and its natural ecosystem's integrity. Local government [23] claims that the distribution area centre of unique and endemic flora and fauna is generally located in the protection block.

The collection block shows the highest diversity value compared to the other two blocks, even though the community is composed of a few species and belongs to the low category (Table 3). The collection block obtains a light intensity at 900–2000 lx, higher than in the protection block at 100–800 lx. The light spectrum within a forest canopy on the collection block can directly affect the photosynthesis and morphogenesis of the herbs and seedling on the forest floor. This study's finding was similar to a study conducted by Satyanti [56], which investigated several species of *Begonia* sp. in Reboisation Forest Bali were received more abundant light at 2246 lx and relatively warmer as its microclimate preference. However, Undaharta et al. [57] have reported contradictory findings on three species of *Begonia* sp. that reproduce through the low light intensity at 200 lx in Manusela National Park, Maluku. In fact, *Begonia* sp. has the ability to cope with the spectra light of the environment as an ecophysiological adaptation by modified the chloroplast structure to tolerate conditions in a wide range of habitat type, [55,57–59]. These findings together imply that the pre-conservation of Anoa in Abdul Latief Forest Park needs an extra effort by utilizing the forage plants abundance as a facility to design an efficient production system to support these animals' survival and welfare.

5. Conclusion

In summary, there were six forage plants in Abdul Latief Forest Park, including *Ficus racemosa* L., *Elatostema integrifolium* (D. Don) Wedd., *Nephrolepis* sp., *Scleria multifoliata* Boeck., *Alpinia* sp., and *Begonia* sp. as Anoa food. *E. integrifolium* (D. Don) Wedd. has the highest abundance among the six forage plants in the protection block. In contrast, the collection block is classified as a high diversity of forage plants due to the highest number of species. This study emphasizes the suitable forage plants for Anoa as our acting for support in-situ and ex-situ conservation activities in Abdul Latief Forest Park. Future studies should address the other blocks observation and microclimate factors that distinguish the abundance and establishment of forage plants. We recommended the local government should carry out integrated conservation action to protecting the endemic species.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgment

We thank Arifuddin, S.Sos., M.Sc., the head of the environment and forestry service of Sinjai Regency, particularly to Ir. Makmur Tasbih, the head of the Abdul Latief Forest Park management division, Syamsul Qamar Ahmad, S.Hut., MM., Yusuf Palulla, S.Hut., Sommeng, and Sultan, S.Hut., M.Sc. as the section heads, for their guidance and cooperation for this research. We also thank the TAHURA team for their contribution to the research sites. This work was supported by Universitas Islam Negeri Alauddin Makassar.

References

- [1] Sekretariat of the Cabinet of the Republic of Indonesia, Undang Undang No. 5 tentang konservasi sumberdaya alam hayati dan ekosistemnya, Peraturan Presiden Republik Indonesia, Jakarta, 1990, pp. 1–12.
- [2] The Ministry of Environment and Forestry Republic of Indonesia, Strategi dan rencana aksi konservasi Anoa (*Bubalus depressicornis* dan *Bubalus quarlesi*) Tahun 2013–2022, Peraturan Menteri Kehutanan Republik Indonesia Nomor P.54/ Menhut-II/2013, Jakarta, 2013, pp. 1–31.
- [3] J. Burton, P. Wheeler, A.H. Mustari, *Bubalus quarlesi*, The IUCN Red List of Threatened Species 2016, e.T3128A46364433, 2016, p. 1.
- [4] J. Burton, P. Wheeler, A.H. Mustari, *Bubalus depressicornis*, The IUCN Red List of Threatened Species 2016, e.T3126A46364222, 2016, p. 1.
- [5] Y. Rusiyantono, M.I. Mumu, Y. Duma, The ability to adapt for ex situ conservation of Anoa (*Bubalus* sp.) through regulation of feeding pattern, *J. Phys. Conf. Ser.* 1242 (2019) 1–5.
- [6] R.P.A. Allo, A. Sudhartono, E. Labiro, Karakteristik komponen abiotik Anoa (*Bubalus* sp.) habitat dalam Hutan Pendidikan Tadulako Universitas Kecamatan Bolano Lambunu Kabupaten Parigi Moutong, *J. War. Rimba J. Ilm. Kehutan.* 6 (1) (2018) 25–31.
- [7] S.N. Ranuntu, R.A. Mallombasang, Studi populasi dan habitat Anoa (*Bubalus* sp.) di kawasan hutan lindung Desa Sanggirona Kabupaten Poso, *Mitra Sains* 3 (2) (2015) 81–94.
- [8] J.A. Burton, S. Hedges, A.H. Mustari, The taxonomic status, distribution and conservation of the lowland Anoa (*Bubalus depressicornis*) and mountain Anoa (*Bubalus quarlesi*), *Mammal Rev.* 35 (1) (2005) 25–50.
- [9] D.I.D. Arini, A. Nugroho, Preferensi habitat Anoa (*Bubalus* spp.) di Taman Nasional Bogani Nani Wartabone, *Pros. Sem. Nas. Masy. Biodiv. Indon.* (2016) 103–108.
- [10] D.I.D. Arini, A. Mayasari, Produktivitas rumput pakan Anoa (*Bubalus* sp.) sekitar penangkaran pada kondisi pra budidaya, *Pastura* 6 (2) (2017) 85–89.
- [11] M. Basri, Selera makan Anoa gunung (*Bubalus quarlesi*) pada sistem kafetaria (Studi prabudidaya untuk penangkaran Anoa di Palu, Sulawesi Tengah), *Agrol. J. Ilmu-ilmu Pertanian.* 16 (3) (2009) 283–289.
- [12] A.H. Mustari, A.U. Prilianti, B. Masyud, Pakan dan perilaku makan Anoa (*Bubalus* sp.) di Taman Margasatwa Ragunan, Jakarta Selatan, *Media Konserv.* 20 (2) (2015) 261–268.
- [13] D.I.D. Arini, Y. Kaffar, Preferensi pakan Anoa (*Bubalus* sp.) di Penangkaran Balai Penelitian Kehutanan Manado, *J. Wasian* 1 (2) (2014) 83–90.
- [14] M. Basri, R. Rukmi, Jenis dan kandungan tanin pakan satwa Anoa (*Bubalus* sp.), *Media Peternak.* 34 (1) (2011) 30–34.
- [15] M. Clauss, S. Reese, K. Eulenberger, A. Fidgett, J.M. Hatt, T. Huisman, J. Hummel, G. Janssen, J. Nijboer, A. Plovman, Macroscopic digestive anatomy of a captive lowland Anoa (*Bubalus depressicornis*), *Zoo Anim. Nutr.* 4 (2009) 255–263.
- [16] A.H. Mustari, Nutritional analysis of Anoa (*Bubalus depressicornis* and *Bubalus quarlesi*) food plants in Tanjung Peropa wildlife reserve, Southeast Sulawesi, *Media Konserv.* 16 (2) (2011) 92–94.
- [17] D.I.D. Arini, Anoa dan habitatnya di Sulawesi Utara, Balai Penelitian Kehutanan Manado, Manado, 2013, pp. 1–71.
- [18] M. Basri, S. Suryahadi, T. Toharmat, H.S. Alikodra, Preferensi pakan dan kebutuhan nutrisi Anoa gunung (*Bubalus quarlesi* Ouwens 1910) pada kondisi prabudidaya, *Media Peternak.* 31 (1) (2008) 53–62.
- [19] The Ministry of Environment and Forestry Republic of Indonesia, National Forest Reference Emission Level for Deforestation and Forest Degradation, Jakarta, 2015, pp. 1–82.
- [20] The Ministry of Environment and Forestry Republic of Indonesia, Deforestasi Indonesia Tahun 2017–2018, Jakarta, 2019, pp. 1–67.
- [21] W. Wardah, E. Labiro, S.D. Massiri, S. Sustru, M. Mursidin, Vegetasi kunci habitat Anoa di Cagar Alam Pangli Binangga, Sulawesi Tengah, *J. Penelit. Kehutan. Wallacea* 1 (1) (2012) 1–12.
- [22] The Ministry of Environment and Forestry Republic of Indonesia, Perubahan fungsi sebagian kawasan hutan lindung Bulu Pattiroang pada kelompok hutan Lompobattang-Apparang seluas +720 hektar yang menjadi kawasan pelestarian alam dengan fungsi taman hutan raya, Keputusan Menteri Kehutanan No. SK.267/ Menhut-II/2008 Tanggal 1 Agustus 2008, Jakarta, 2008, pp. 1–10.
- [23] Environment and Forestry Service Sinjai Regency, Profil Taman Hutan Raya Abdul Latief Sinjai, Government of Sinjai Regency, Sinjai, 2018, p. 128.
- [24] K.R. Schwartz, E.C.M. Parsons, L. Rockwood, T.C. Wood, Integrating *in-situ* and *ex-situ* data management processes for biodiversity conservation, *Front. Ecol. Evol.* 5 (120) (2017) 1–7.
- [25] D.I.D. Arini, M. Christita, A. Mayasari, R. Suryaningsih, A.T.A.J. Simamora, A review of Anoa conservation efforts in Sulawesi, Indonesia, *IOP Conf. Ser.: Earth Environ. Sci.* 533 (2020) 1–14.
- [26] R.I. Pujaningsih, C.I. Permana, I.G. Sutrisno, Study of diet preference and feeding behaviour of Anoa (*Bubalus* sp.) in wildlife Park Ragunan for ex situ conservation and cultivation, in: The 3rd International Conference on Sustainable Animal Agriculture for Developing Countries, Nakhon Ratchasima, Thailand vol. 3, 2011, pp. 317–321.
- [27] D.I.D. Arini, N.I. Wahyuni, Kelimpahan tumbuhan pakan Anoa (*Bubalus* sp.) di Taman Nasional Bogani Nani Wartabone, *J. Penelit. Kehutan. Wallacea* 5 (1) (2013) 91–102.
- [28] C.C. Berg, E.J.H. Corner, Flora Malesiana (Moraceae-Ficus), *National Herbarium Nederland, Univ.t Leiden Branch* 17 (2) (2005) 730.
- [29] T.C. Whitmore, I.G.M. Tantra, U. Sutisna, Tree Flora of Indonesia Check List for Sulawesi [1989], Ministry of Forestry, Agency for Forestry Research and Development, Forest Research and Development Centre, Jakarta, 1989, p. 204.
- [30] N. Kurnia, O. Jumadi, S.F. Hiola, Atlas tumbuhan Sulawesi Selatan, Jurusan Biologi Fakultas Matematika dan Ilmu Pengetahuan Alam Universitas Negeri Makassar, 2014, p. 205.
- [31] R.I. Pujaningsih, Anoa (*Bubalus* sp.). Badan Penerbit Universitas Diponegoro, Semarang, 2012, pp. 1–102.
- [32] W.H. Ardi, T. Chikmawati, J.R. Witono, D.C. Thomas, A synopsis of Begonia (Begoniaceae) of southeastern Sulawesi including four new species, *Phytotaxa* 381 (1) (2018) 27–50.
- [33] I.T.P. Tangkoro, E. Labiro, I.N. Korja, Komposisi jenis-jenis pakan Anoa di kawasan Hutan Pendidikan Universitas Tadulako Kecamatan Bolano Lambunu Kabupaten Parigi Moutong, *J. War. Rimba J. Ilm. Kehutan* 6 (2) (2018) 1–9.
- [34] J.R. Bray, J.T. Curtis, An ordination of the upland Forest communities of southern Wisconsin, *Ecol. Monogr.* 27 (4) (1957) 325–349.
- [35] E.P. Odum, Dasar-dasar ekologi, Gadjah Mada University Press, Yogyakarta, 1993, p. 697.
- [36] R.I. Pujaningsih, Identification of natural feed of Anoa (*Bubalus* spp.) in advance of wildlife conservation, in: Proceedings of the XXVII Congress of The International Union of Game Biologist, Hannover, 2005.
- [37] B.W. Broto, Struktur dan komposisi vegetasi habitat Anoa (*Bubalus* spp.) di Hutan Lindung Pegunungan Mekongga, Kolaka, Sulawesi Tenggara, *Biodiversitas, J. Biol. Divers.* 1 (3) (2015) 615–620.
- [38] A.K. Bhadra, S.K. Pattanayak, Abundance or dominance: which is more justified to calculate importance value index (IVI) of plant species, *Asian J. Sci. Technol.* 7 (9) (2016) 3577–3601.
- [39] I. Fatahillah, I.F. Lestari, K. Salsabila, R. Pratiwi, T. Amalia, A. Septyaningsih, U. Kulsun, R.H. Ristanto, A. Sedayu, Inventarisasi tumbuhan paku di jalur Ciwalean Taman Nasional Gunung Gede Pangrango, Jawa Barat, *Biog. J. Ilm. Biol.* 6 (1) (2018) 43–45.
- [40] R.I. Pujaningsih, B. Sukanto, E. Labiro, Identifikasi dan teknologi pengolahan pakan hijauan dalam upaya penangkaran Anoa, Fakultas Peternakan Universitas Diponegoro, Semarang, 2004, pp. 1–44.
- [41] A. Powling, A. Phillips, R. Pritchett, S.T. Segar, R. Wheeler, A. Mardiatuti, The vegetation of Lambusango Forest, Buton, Indonesia, *Reinwardtia* 14 (2) (2016) 265–286.
- [42] I.T.P. Tangkoro, E. Labiro, I.N. Korja, Komposisi jenis-jenis pakan Anoa di Kawasan Hutan Pendidikan Universitas Tadulako Kecamatan Bolano Lambunu Kabupaten Parigi Moutong, *J. War. Rimba* 6 (2) (2018) 1–9.
- [43] R.I. Pujaningsih, C.I. Sutrisno, Y. Supriandho, A. Malik, S. Pudyatmoko, M.A. Amir, S. Aryanto, Diet composition of Anoa (*Bubalus* sp.) studied using direct observation and dung analysis method in their habitat, *J. Indones. Trop. Anim. Agric.* 34 (3) (2009) 223–228.
- [44] M.D. Shah, Y.S. Yong, M. Iqbal, Phytochemical investigation and free radical scavenging activities of essential oil, methanol extract and methanol fractions of *Nephrolepis biserrata*, *Int J Pharm Pharm Sci* 6 (9) (2014) 269–277.
- [45] A.O. Adebiyi, Secondary metabolites and bioactivities of two species of *Nephrolepis* in Ekiti state, Nigeria, *Int. J. Life Sci.* 8 (3) (2019) 95–99.
- [46] M.E. Bassey, I.I. Johnny, O.T. Umoh, F.T. Douglas, Phytomedicinal potentials of species of *Nephrolepis* (Schott.), world, *J. Pharm. Res.* 9 (4) (2020) 1400–1410.
- [47] S. Sajeev, P. Raj, D.B. Adarsh, S. Hegde, Phytoconstituents of *Nephrolepis hirsutula* and *Pityrogramma calomelanos*, two medicinal ferns of the Western Ghats, *Indian Fern J.* 32 (2015) 244–256.
- [48] D. Iyer, B.K. Sharma, U.K. Patil, Isolation of bioactive phytoconstituent from *Alpinia galanga* L. with anti-hyperlipidemic activity, *J. Diet. Suppl.* 10 (4) (2013) 309–317.
- [49] A.A. Ahmed, F. Sharmen, A. Mannan, M.A. Rahman, Phytochemical, analgesic, antibacterial, and cytotoxic effects of *Alpinia nigra* (Gaertn.) Burtt leaf extract, *J. Tradit. Complement. Med.* 5 (4) (2015) 248–252.
- [50] R. Ramadani, D. Damry, R. Rusdy, B. Hamzah, M.S. Zubair, Traditional usages and phytochemical screenings of selected Zingiberaceae from Central Sulawesi Indonesia, *Pharmacogn.* 11 (3) (2019) 505–510.
- [51] T. Uji, Keenekaragaman dan potensi flora di Cagar Alam Muara Kendawangan, Kalimantan Barat, *Biodiv. J. Biol. Divers.* 4 (1) (2003) 112–117.

- [52] R.K. Yadav, B.C. Nandy, S. Maity, S. Sarkar, S. Saha, Phytochemistry, pharmacology, toxicology, and clinical trial of *Ficus racemosa*, *Pharmacogn. Rev.* 9 (17) (2015) 73–80.
- [53] P. Sivakumar, I. Manimekalai, K. Sivakumari, K. Ashok, Phytochemical profiling of fig fruit (*Ficus racemosa*) extracts, *Int. J. Res. Anal. Rev.* 6 (1) (2019) 784–822.
- [54] M. Basri, R. Rukmi, Jenis dan kandungan tanin pakan satwa Anoa (*Bubalus sp.*), *Media Peternak.* 34 (1) (2011) 30–34.
- [55] S.A. Sumi, A. Siraj, A. Hossain, S. Mia, S. Afrin, M. Rahman, Investigation of the key pharmacological activities of *Ficus racemosa* and analysis of its major bioactive polyphenols by HPLC-DAD, evidence-based complement, *Altem. Med.* (2016) 1–10.
- [56] A. Satyanti, The natural habitat of *Begonia* on the island of Bali: microclimate preference in two forest sites in Bedugul, Biotropia Southeast Asian *J. Trop. Biol.* 19 (2) (2012) 80–91.
- [57] N.K.E. Undaharta, S. Sutomo, M. Ardaka, I.G. Tirta, Autekologi *Begonia* di sebagian Kawasan Taman Nasional Manusele, Maluku, *J. Penelit. Hutan dan Konserv. Alam* 9 (1) (2012) 1–11.
- [58] M. Théry, Forest light and its influence on habitat selection, in: K.E. Linsenmair, A. J. Davis, B. Fiala, M.R. Speight (Eds.), *Tropical Forest Canopies: Ecology and Management*, 2001, pp. 251–261.
- [59] M. Jacobs, M. Lopez-Garcia, O.P. Phrathep, T. Lawson, R. Oulton, H.M. Whitney, Photonic multilayer structure of *Begonia* chloroplasts enhances photosynthetic efficiency, *Nat. Plants* 2 (11) (2016) 1–6.

Identification, abundance and diversity of forage plants for Anoa (*Bubalus spp.*) in Abdul Latief Forest Park

ORIGINALITY REPORT

6%

SIMILARITY INDEX

4%

INTERNET SOURCES

4%

PUBLICATIONS

%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

1%

★ repository.futminna.edu.ng:8080

Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On