

Short Communication

Feed Nutrition Evaluation and Metagenomic Analysis

Hafsan*

Department of Biology, Universitas Islam Negeri Alauddin, Makassar, South Sulawesi Indonesia

*Correspondence to: Hafsan, E-mail: hafsan.bio@uin-alauddin.ac.id, Tel: +62-8114448781

ABSTRACT

Feed is a source of nutrients needed by livestock and is one of the main determining factors in increasing productivity. Evaluating the quality and absorption of feed nutrients with a metagenomic approach is promising to detect and overcome livestock production constraints accurately and more quickly. Identifying the whole microbiota in the digestive tract of livestock is essential because the involvement of microbes in nutrient metabolism in the gastrointestinal tract directly or indirectly affects the productivity and quality of livestock products. This technique serves as feed quality control to determine the accuracy of the most suitable feed formulation to ensure the absorption of feed nutrients for the metabolic processes of livestock.

KEYWORDS: metagenomics, microbiota, NGS, Nutrition Evaluation

INTRODUCTION

Microbes that live in a free environment have differences with microbiota that live in the bodies of animals, including livestock. Microbes in the free environment have extreme nature because they are exposed to a dynamic external environment. In contrast to the microbiota environment inside the animal body, it is considered an exceptional environment. After all, the conditions are relatively warm, ectotrophic, and stable. The microbiota that lives and settles in the body consist primarily of bacteria, eukaryotic, archaea, and a small portion is the viruses.

Microbiota is an essential factor for the development process and immunity of the host. Except for pathogenic bacteria, bacteria in the host's body are harmless; even colonization of microbiota in some body regions plays a beneficial role. The role of the gut microbiota is to assist in the digestion and metabolism of nutrients, regulate the immune system, and protect against pathogens. Each individual has a different response to the metabolism of the microbiota, which determines the health status of the organism as a critical and a consequential factor for the host (Gabriel et al., 2006).

Metagenomics is an appropriate technique to understand the structure of the microbiota community in livestock without going through cultivation first. The principle of diversity analysis in microbiota communities with metagenomics is based on DNA analysis collected directly from the spot of a community or ecosystem. The DNA from the community can be directly analyzed for diversity, and the species composition of the community can be identified, using phylogenetic markers such as the gene encoding 16s rRNA (ribosome-Ribonucleic Acid), which is the carrier of the hereditary properties of living things, including microbes (Hugenholtz & Tyson, 2008). In detail, this metagenomic technique attempts to read DNA from the entire microbial composition contained in the sample.

With metagenomic analysis, an organism or group of microbes in detail and accurately can be known for its function and role in livestock. This metagenomic technique was developed to solve the weakness in analyzing unculturable microbes from an ecosystem that cannot grow in artificial growth media. The metagenomic analysis develops conventional molecular techniques to identify a particular

microbial isolate. The fundamental difference is in the scope of reading DNA sequences as genetic information can be carried out more thoroughly, in detail and accurately and simultaneously (Singh et al., 2012). Therefore, metagenomics can be called NGS (Next Generation Sequencing) technology. For example, microbes with a gene region of 16s rDNA (DNA commonly found in bacteria to encode 16s rRNA, which is conserved) will be read accurately and entirely with this NGS technology. It is not just a profile of bacterial DNA sequences; its functional activity in the metabolism of nutrients in the livestock body can be predicted with accuracy. In other words, the results of the sequences are no longer partial but in the form of complete data that provides an overview of the composition of the microbiota that inhabits the environment in the body of livestock.

Metagenomic applications in poultry farming can be used to determine the nature and role of microbes in the digestive tract of livestock. The involvement of microbes in nutrient metabolism in the gastrointestinal tract directly or indirectly affects the productivity and quality of livestock products (meat, eggs and milk). The metagenomic analysis will quickly read the complexity and composition of microbes in the digestive tract. It helps diagnose the cause of disease due to pathogenic microbes in poultry which can consist of more than one species, which is challenging to identify with conventional techniques. Through this metagenomic approach, identifying groups of pathogenic microbes that cause disease or trigger stunting of livestock can be known accurately and simultaneously, which is very difficult if done by conventional methods.

Technically, the application of metagenomics in evaluating animal feed nutrition is beneficial to know the microbes involved in the digestion of food substances, how the metabolic processes occur and what enzymes are involved through specific metabolic pathways. This parameter will help determine feed formulation strategies to improve livestock performance by increasing feed efficiency value, making it more economical and productive. In other words, we can ensure the availability of nutrients available in the feed to be absorbed in the digestive tract and then involved in metabolic and biosynthetic processes until finally converted into body mass (Stanley et al., 2012).

The stages of metagenomic analysis of broiler digestive tract microbiota were carried out by collecting broiler digesta samples at the end of rearing. Furthermore, DNA extraction from the microbiota that inhabits the digesta was carried out and continued with amplification (multiplication of sequences) of DNA in specific target genes. This process uses specific primers in the PCR (polymerase chain reaction) cycle. Furthermore, the PCR results on specific target genes were sequenced with an NGS tool (e.g. MiSeq Sequencer). The result is a DNA sequence profile which is then analyzed using the QIIME (Quantitatively Insight into Microbial Ecology) computational program.

As it is known that the number of microbiota in each individual has a significant difference compared to the variation of the genome. Each genome in the same species has 99% similarity. However, the microbiota in each individual has a difference of approximately 80-90%. The gut microbiota of chickens is very complex. In the ileum area of broilers fed with phytase supplementation, more than 900 different bacterial species were found. The species diversity index of Shannon-Wiener bacteria inhabiting the broiler ileum was 5.95. There is variability in the composition of broiler microbiota on different diets accompanied by different growth performance values. Sequencing genetic bands from metagenomics allow the identification of several bacterial species that are thought to contribute to changes in the microbiota in broilers with different performance levels. Information on the diversity of microbiota communities in some areas of the digestive tract helps make decisions related to the handling of nutrient absorption constraints due to the balance of nutrients in the feed that has not been maximized.

Feed supplemented with bacterial phytase as additives to increase nutritional utility has improved growth performance. Several factors can contribute to this performance variability, including differences in the expression of host adaptive genes affected by nutrient uptake. Broilers that are reared hold similar genetics and are raised together in a carefully controlled environment with access to the same feed and water. Information on the variability of the composition of the digestive tract microbiota can be correlated with the variability of the broiler's ability to convert nutrients into

body mass. Identifying microbiota that can affect the efficiency of nutrient absorption by broilers will be very useful because it will help direct efforts to develop ways to manipulate the digestive tract microbiota to achieve maximum productivity. More specifically, this information will serve as supporting data related to increasing the efficiency of the utilization of phosphorus and other minerals bound by phytate and reducing its negative effect on nutrient utilization through breaking its bonds with the hydrolysis process by the phytase enzyme (Hafsan et al., 2021).

REFERENCES

- Gabriel, I., Lessire, M., Mallet, S., and Guillot, J. F. 2006. Microflora of the digestive tract: critical factors and consequences for poultry. *World's Poultry Science Journal*, 62(03), 499–511.
- Hafsan, Thaha, A. H., Natsir, A., and Ahmad, A. 2021. Performance, Carcass and Internal Organs Characterizes of Broiler Chickens with Phytase Supplementation from *Burkholderia* sp. Strain Hf.7. *Online Journal of Animal and Feed Research*, 11(2), 52–56.
- Hugenholtz, P., and Tyson, G. W. 2008. Metagenomics. *Nature*, 445(25 September), 481–483.
- Singh, K. M., Ahir, V. B., Tripathi, A. K., Ramani, U. V., Sajjani, M., Koringa, P. G., Jakhesara, S., Pandya, P. R., Rank, D. N., Murty, D. S., Kothari, R. K., and Joshi, C. G. 2012. Metagenomic analysis of Surti buffalo (*Bubalus bubalis*) rumen: A preliminary study. *Molecular Biology Reports*, 39(4), 4841–4848.
- Stanley, D., Denman, S. E., Hughes, R. J., Geier, M. S., Crowley, T. M., Chen, H., Haring, V. R., and Moore, R. J. 2012. Intestinal microbiota associated with differential feed conversion efficiency in chickens. *Applied Microbiology and Biotechnology*, 96(5), 1361–1369.