

Sarfaini Suyuti

by Sarfaini Suyuti

Submission date: 30-Jan-2023 11:55AM (UTC+0700)

Submission ID: 2002213826

File name: 111_REVISION.docx (36.78K)

Word count: 2549

Character count: 13495

1

The effect of substituting carrots (*Daucus Carota L.*) into tempeh brownies on the intake of protein energy deficiency in Indonesia

Syarfaini*, Sukfitrianty Syahrir, Nildawati, Fatmawaty Mallapiang, Surahmawati, Uswatun Hasanah

Department Of Public Health, Faculty Of Medicine And Health Sciences, Universitas Islam Negeri Alauddin, Makassar 92113, Indonesia

*Corresponding author.

E-mail Address: syarfaini.suyuti@uin-alauddin.ac.id; pmc@agri.unhas.ac.id (Syarfaini).

Abstract 1

Objective: This research investigates the use of carrot (*Daucus carota L.*) to substitute tempeh as the main ingredient of tempeh brownies and its efficacy in changing the nutritional status of students with PEU.

Methods: Quasi-experimental method, non randomised control group pretest research design. The samples consist of 22 students who were selected using purposive sampling technique (case and control).

Results: The energy intake of case group is 91.94 kilocalories ($p=0.005$) while the one in control group is 80.6 kilocalories ($p=0.005$). The protein intakes of case group and control group are 7.55 grams ($p=0.010$) and 6.95 grams ($p=0.038$) respectively. The findings also reveal that the body weight in case groups is 0.91 kg ($p=0.000$) in the control group is 0.57 kg ($p=0.000$).

Conclusions: There is an effect of giving carrot tempeh brownies (*Daucus carota L.*) to changes in energy intake and protein intake in students lacking in protein energy.

KEYWORDS: Protein Energy Undernutrition; Carrot Substituted (*Daucus Carota L.*); Tempeh Brownies.

Introduction

Protein Energy Deficiency (PEM) is one of the nutritional disorders that is important for Indonesia and for developing countries in Asia, Africa, Central America and South America.¹ Children's protein needs include maintenance of tissues, changes in body composition and formation of new tissue. Protein requirements for growth are estimated to range from 1-4 g / kg in addition to body tissue.²⁻⁵ World Health Organization (WHO) in 2015 reported that the prevalence of carelessness in children around the world was around 14.3% with the number of children experiencing carping as much as 95.2 million children.⁶ Trends in nutritional problems in Indonesia based on the results of monitoring nutritional status from 2015 to 2017 tend to decrease.⁷ In the category of malnutrition (BB / U) there were 18.8% (2015), 17.8% (2016) and 17.9% (2017).⁸ Optimal child growth and development depends on the provision of nutrients with good and correct quality and quantity.⁹

Based on the results of Riskesdas in 2010, the nutritional status of children aged 6-12 years (BMI / U) in Indonesia with a thin prevalence of 12.2% consisted of 4.6% very thin and 7.6% thin.¹⁰⁻¹³ While the results of Riskesdas in 2013, it is known that the prevalence of malnutrition in the national thin category (BMI / U) aged 5-12 years is 11.2% consisting of 4% very thin and 7.2% thin.^{14,15} South Sulawesi is one of the provinces in Indonesia which has nutritional problems in children aged 5-12 years. Based on the 2016 Nutrition Status Monitoring (PSG) results in South Sulawesi Province for the category of school

children aged 5-12 years (BMI / U) it was found to be very thin 2% and 9.4% thin while in 2017 it decreased for the category of very thin school children to 1.7% and thin 7.1%.^{8,16,17} Based on the results of monitoring the nutritional status of the Ministry of Health in 2017 in Gowa, it is known that the nutritional status of children aged 5-18 (BMI / U) years is the prevalence of very thin and thin 10.1%.¹⁴ Nutritional status of children aged 5-18 years according to BMI / U for the prevalence of very thin and thin in Gowa Regency is ranked eighth.¹⁸

This is consistent with the results of monitoring the nutritional status of the Ministry of Health in 2017 the highest is Palopo (15.3%), Maros (13.4%), Barru (12.6%), Bulukumba (11.9%), Sidrap (11.9%), Jeneponto (10.9%), Sinjai (10.6%), Gowa (10.1%), Selayar (9.9%) and Pangkep (9.2%).¹⁵ The basic ingredients used during the intervention are tempeh made from grains and carrots from vegetables. Tempe is one of the foods with good nutrition and is fairly cheap but has many health benefits. Tempe is a perishable food ingredient, so to improve the durability, diversity and attractiveness of consumers, modifications are needed to the food ingredients. Besides tempeh has a distinctive taste because of the fermentation process so to modify the food ingredients need additions such as carrots that have attractive colors and sweet taste.^{16,19,20}

Methods

Research location

This research was conducted on a number of students in grades I-III aged 7-9 years who experienced a lack of protein nutritional status in SD Inpres Bakung, Gowa Regency. This study used a Quasi Experiment study, a non randomized control group pretest design design with a sampling method that is purposive sampling, aimed to determine the effect of giving carrot tempeh brownies (*Daucus carota L.*) to the intake of students lacking in protein energy in SD Inpres Bakung, Gowa Regency. Samples that met the inclusion and exclusion criteria were 22 students who were divided into 2 groups namely cases and controls.

Types and sources of data

Primary data obtained through direct observation on the research object that is doing anthropometric measurements based on BB / U include weight measurements of the research object and interviews to determine food intake in the sample using 24-hour recall. Secondary Data Obtained through library research, research journals, relevant literature books, reports and related agencies.

This research has been evaluated and given recommendations from the Health Research Ethics Committee of the Faculty of Medicine and Health Sciences, Alauddin State Islamic University, Makassar with Number: A.075 / KEPK / FKIK / II / 2019.

Data collection techniques

This method starts with a 24-hour food recall from the respondent waking up yesterday morning until the night's sleep rest or can also be started from the time when the interview was carried out backwards to 24 hours full with the help of photographs of food based on household size (URT) to facilitate the respondent considering the size / portion of food that has been used for the past 24 hours. Then use the 24-hour food recall form to find out energy and protein consumption. The recall results were then calculated with the help of the 2007 Nutry Survey computer software and then calculated with the

Nutrition Adequacy Rate (RDA) of children aged 7-9 years and the level of respondents' consumption of the RDA. Furthermore, to monitor the consumption of tempeh brownies and carrot tempeh brownies (*Daucus carota* L.) on respondents. Monitoring brownie consumption is done by weighing the remaining brownies consumed by respondents using digital food scales. Then do a digital weight weighing to measure the weight of the sample (with a capacity of 150 kg with an accuracy of 0.1 kg).

Result

Table 1 Characteristics of Respondents The Effect of Giving Carrot Substitute Brownies (*Daucus carota* L.) Brownies Against Student Intake of Protein-less Energy

Characteristics of Respondents	Case		Control	
	N	%	n	%
Age				
7 Years	8	72.7	7	63.6
8 Years	2	18.2	2	18.2
9 Years	1	9.1	2	18
Gender				
Women	3	27.3	7	63.6
Men	8	72.3	4	36.4
Grade Level				
Class I	0	0	5	45.5
Class II	8	72.7	2	18.2
Class III	3	27.3	4	36.3
Fathers Occupation				
Laborers	5	45.5	7	63.6
General Employees	1	9.1	2	18.2
Furniture Work	1	9.1	0	0
TV Service	1	9.1	0	0
PLN	1	9.1	0	0
Driver	1	9.1	0	0
Entrepreneur	1	9.1	2	18.2
Mothers Job				
Housewife	11	100	10	90.9
Entrepreneur	0	0	1	9.1
Total Product Consumption				
Consumed	3870.7	70.9	3655.5	67
Not Consumed	1589.3	29.1	1804.5	33
Average Product Consumption				
Consumed	148.9	70.9	140.6	67
Not Consumed	61.1	29.1	69.4	33

Source : Primary Data, 2019

Table 1 shows the highest age group in the case group is 7 years old by 8 respondents (72.7%) and the lowest is the age of 9 years with 1 respondent (9.1%) while the highest control group was 7 years old with 7 respondents (63.6%) and the lowest was 8 and 9 years old with 2 respondents (18.2%)

respectively. The sex in the highest case group was 8 respondents (72.7%) and the lowest was 3 respondents (27.3%) while the highest control group was 7 respondents (63.6%) women and the lowest was male with 4 respondents (36.4%). Class levels in the case group were 8 respondents (72.7%) class II and 3 respondents (27.3%) class III while the highest control group was class I as many as 5 respondents (45.5%) and the lowest was class II as much 2 respondents (18.2%).

Table 2 Univariate Analysis Based on Average Energy Intake and Protein Intake of Respondents Before and After Intervention

Average Intake	Case Group	Control Group	<i>Independent t-test</i>
Before Intervention			
Energi Intake (kkal)	833.93	687.32	0.027
Protein Intake (g)	34.06	25.45	0.044
After Intervention			
Energi Intake (kkal)	925.88	767.92	0.026
Protein Intake (g)	41.61	32.4	0.020

Source : Primary Data, 2019

Table 2 shows the average energy intake and protein intake of the respondents before the intervention obtained Independent Test t-test results on each variable. In the variable energy intake ($p = 0.027$) and protein intake ($p = 0.044$) p values < 0.05 were concluded "statistically there are differences in the mean of energy intake and protein intake between the case group and the control group before the intervention". Based on the average energy intake and protein intake of the respondents after the intervention the Independent Test t-test results were obtained for each variable. On the variable energy intake ($p = 0.026$) and protein intake ($p = 0.020$).

Table 3 Bivariate Analysis Based on Average Changes in Energy Intake and Protein Intake Case Group and Control Group Before and After Intervention

Intake	Intake mean		Mean (difference)	Paired t-test
	Before	After		
Energi				
Case Group	833.92	925.88	91.94	0.005
Control Group	687.32	767.92	80.6	0.005
Protein				
Case Group	34.06	41.61	7.55	0.010
Control Group	25.45	32.4	6.95	0.038

Source : Primary Data, 2019

Table 3 shows on the average change in energy intake and protein intake in the case group and the control group before and after the intervention. Energy intake in the case group obtained p value = 0.005 and the control group obtained p value = 0.005 which is smaller than the value of $\alpha = 0.05$ the results obtained that there is the effect of intervention on increasing energy intake in each group. In the case group, the average increase in energy intake was 91.94 kkal and in the control group the average increase in energy intake was 80.6 kkal. Protein intake in the case group obtained p value = 0.010 and the control group obtained p value = 0.038 which is smaller than the value of $\alpha = 0.05$, the results obtained that there is an intervention effect on increasing protein intake in each group. In the case

group, the average increase in protein intake was 7.55 grams and in the control group the average increase in protein intake was 6.95 kcal (Primary Data, 2019).

Discussion

Both interventional groups experienced an increase in energy intake but a significant increase in energy intake occurred in the case group with an average difference in energy intake of 91.94 kcal while in the control group showed an average difference in energy intake of 80.6 kcal. The difference in respondent's energy intake between the case and control groups was due to the different nutrients contained in both of them that affected the metabolic process and the nutritional content of tempe brownies and carrot tempeh brownies (*Daucus carota* L.) given to each different group. The lack of product intake in the control group was due to intake based on organoleptic test results conducted by Isma (2016), tempeh brownies (1: 0) had lower organoleptic values than carrot tempeh brownies in terms of color, aroma, texture and taste. Although there was an increase in energy and protein intake in the case and control groups, this amount was still less than the energy requirements in accordance with the 2013 AKG.¹⁶ This can be seen in the energy consumption of case group respondents and the control group in the average energy intake before and after the intervention. but not significant increase because both the case group and the control group prefer outside snacks and consume less rice, some respondents did not spend the brownies given because of the taste of the brownies so that the recommended energy intake based on the RDA was insufficient even still far from the required RDA. However, students prefer to consume carrot tempeh brownies given instead of tempe brownies. This is what makes the students' body weight lacking in protein energy both in the case group and the control group only slightly increases and the nutritional status of malnutrition into good nutrition is also slightly.

Conclusion

The results showed that: 1) There were differences in energy intake and protein intake in students lacking protein energy before and after administration of carrot tempeh brownies (*Daucus carota* L.). 2) There is an effect of giving carrot substitution tempeh brownies (*Daucus carota* L.) to changes in energy intake and protein intake in protein-deficient students before and after giving carrot substitution tempeh brownies (*Daucus carota* L.).

References

1. Kementerian Kesehatan Republik Indonesia. Buku Saku Pemantauan Status Gizi (PSG) Tahun 2016. Jakarta: Dirjen Kesehatan Masyarakat; 2017.
2. Purnamasari DU. Panduan Gizi & Kesehatan Anak Sekolah. Yogyakarta: Andi; s. f.
3. Almatsier S. Prinsip Dasar Ilmu Gizi. 1.^a edición. Jakarta: PT. Gramedia Pustaka Utama; 2001.
4. Almatsier S. Prinsip Dasar Ilmu Gizi. 2.^a edición. Jakarta: PT. Gramedia Pustaka Utama; 2009.
5. Adriani M., Wirjatmadi B. Peranan Gizi dalam Siklus Kehidupan. Jakarta: Kencana Pranada Media Group; 2012.
6. World Health Organization. World Health Statistics 2015.
7. Briawan D. Ilmu Gizi Teori dan Aplikasi. Jakarta: Penerbit Buku Kedokteran EGC; 2017.
8. Kementerian Kesehatan Republik Indonesia. Bahan Ajar Penilaian Status Gizi.

9. Arsyad NN. Pengaruh Pemberian Otak-Otak Ikan Kembung (*Rastrelliger brachysoma*) Substitusi Buah Lamun (*Enhalus acoroides*) Terhadap Status Gizi Pada Siswa Gizi Kurang Di SDN Cambaya Kecamatan Ujung Tanah Kota Makassar 2016. UIN Alauddin Makassar, 2017.
10. Kementerian Kesehatan Republik Indonesia. Angka Kecukupan Gizi Energi, Protein Yang Dianjurkan Bagi Bangsa Indonesia. Lampiran Peraturan Menteri Kesehatan Republik Indonesia Nomor 75/2013.
11. Badan Penelitian dan Pengembangan Kesehatan. Laporan Riset Kesehatan Dasar. Jakarta; 2007.
12. Badan Penelitian dan Pengembangan Kesehatan. Laporan Riset Kesehatan Dasar. Jakarta; 2013.
13. Badan Pusat Statistik Jakarta Pusat. Statistik Indonesia Tahun 2018. Jakarta Pusat: Badan Pusat Statistik; 2018.
14. Dinas Kesehatan Sulawesi Selatan. Dinas Kesehatan Sulsel Fokus Delapan Program Prioritas.
15. Dinas Kesehatan Sulawesi Selatan. Data Bina Kesehatan Masyarakat.
16. Isma R. Analisis Kandungan Zat Gizi Brownies Tempe Substitusi Wortel (*Daucus Carota L.*) Sebagai Alternatif Perbaikan Gizi Terhadap Masyarakat. *J Media Gizi Pangan*. 2016;24(2):76-82.
17. Lestari RA., Syarfaini., Satrianegara F. Pengaruh Pemberian Nugget Tempe Dengan Substitusi Ikan Gabus Terhadap Status Gizi Anak Sekolah Dasar di MIS DDI Ainus Syamsi Kel. Lette, Kota Makassar 2014. *Al-sihah Public Heal Sci J*. 2016;8(2):151-60.
18. Anggraeny O., Dianovita C., Putri EN., Sastrina M., Dewi RS. Korelasi Pemberian Diet Rendah Protein Terhadap Status Protein, Imunitas, Hemoglobin dan Nafsu Makan Tikus Wistar Jantan. *IJHN Indones J Hum Nutr*. 2016;3(2):105-22.
19. Pratiwi CE. Pengaruh Proporsi Kedelai (*Glycine max*) dan Lamtoro Gung (*Leucaena leucephala*) dengan Penambahan Angkak Terhadap Karakteristik Tempe. Universitas Sanata Dharma, 2018.
20. Kusumawati HN., Ratna Noer E. Perbedaan Asupan Zat Gizi Makro sebelum dan Setelah Pemberian Makanan Pendamping ASI (MP-ASI) Bubur Instan Berbasis Ikan Gabus dan Labung Kuning Pada Balita Gizi Kurang. Universitas Diponegoro, 2016.

Sarfaini Suyuti

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

5%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

minio.umkt.ac.id

Internet Source

11%

2

B. Suriani, Jumrah Sudirman, Sitti Mukarramah, Syamsyuriyana Sabar, Hasriani Saleng. "Fermented soybean cake nugget (tempeh) as an alternative for increasing weight of little children aged 36–60 months", Gaceta Sanitaria, 2021

Publication

3%

3

Fangfei Ji, Yuexin Yang, Lei Xu, Jing Cai, Mingde Ni, Qiuzhen Wang, Shanliang Zhao, Aiguo Ma. "Poor diet quality evaluated with the China Healthy Diet Index in Chinese tuberculosis patients", Journal of Human Nutrition and Dietetics, 2021

Publication

2%

Exclude quotes Off

Exclude bibliography On

Exclude matches < 2%