

Dietary thymol improved growth, body composition, digestive enzyme activities, 2 hematology, immunity, antioxidant defense, and resistance to Streptococcus iniae in the 3 rainbow trout (Oncorhynchus)

Submission date: 21-Mar-2023 02:32PM (UTC+0700)

Submission ID: 2042509917

File name: Manuscript_Hafsan_et_al.pdf (869.53K)

Word count: 9886

Character count: 49424

by Hafsan Hafsan

1 **Dietary thymol improved growth, body composition, digestive enzyme activities,**
2 **hematology, immunity, antioxidant defense, and resistance to *Streptococcus iniae* in the**
3 **rainbow trout (*Oncorhynchus mykiss*)**

4
5 **Abstract**

6 In this study, thymol (TYM) at dietary levels of 0, 1, 1.5, 2, 2.5 g/kg diet was used to evaluate
7 its effects on growth, digestive performance, immunity and resistances to the infection induced
8 by *Streptococcus iniae* in the rainbow trout, *Oncorhynchus mykiss*. A number of 450 fish (35.8
9 ± 4.4 g; Mean ± SD) were distributed to 15 tanks (30 fish/tank) in three replicates and fed TYM
10 for 60 days. After feeding period, Fish fed 1.5-2.5 g TYM showed better growth, higher
11 digestive enzyme activity and body protein content compared to other diets (P<0.05).
12 Regression analysis indicated a polynomial relationship between growth parameters and
13 dietary TYM levels. Based upon the varied growth parameters, the optimum dietary TYM
14 level, was 1.89 % for FCR. TYM at dietary levels of 1.5-2.5 g significantly enhanced liver
15 antioxidant enzyme activity [superoxide dismutase (SOD), glutathione peroxidase (GPx),
16 catalase (CAT)], immune components in blood [alternative complement activity (C3), total
17 immunoglobulin (Ig), lysozyme activity, bactericidal activity and total protein)] and in mucus
18 [alkaline phosphatase (ALP), protease activity, lysozyme activity, bactericidal activity and
19 total protein)] compared to other diets (P<0.05). TYM at dietary levels of 2-2.5 g significantly
20 decreased malondialdehyde (MDA) levels compared to other experimental groups (P<0.05).
21 In addition, use of TYM at dietary levels of 1.5-2.5 g up-regulated the expression of the
22 immune-related genes (*C3*, *Lyz* and *Ig*) (P<0.05). In contrast, the expression of inflammatory
23 genes, tumor necrosis factor (TNF- α) and Interleukin-8 (IL-8) significantly were down-
24 regulated in response to 2-2.5 g TYM (P<0.05). The hematology of the fish also altered in
25 response to dietary TYM, where the values of corpuscular hemoglobin concentration (MCHC),
26 hemoglobin (Hb), red blood cell (RBC), hematocrit (Hct) and white blood cell (WBC)
27 significantly increased in fish fed 2-2.5 g TYM compared to other diets (P<0.05). In addition,
28 MCV significantly decreased in response to 2-2.5 g TYM (P<0.05). After challenge with
29 *Streptococcus iniae*, the survival rate was significantly higher in fish fed 2-2.5 g TYM
30 compared to other diets (P<0.05). The results of the present study concluded that TYM in the
31 diet of rainbow trout can improve the fish growth and immunity and increase the resistance of
32 the fish to *Streptococcus iniae* infection. The results of this study recommend an optimized
33 dietary level of 2-2.5 g TYM for the fish.

34 **Keywords:** Thymol, Growth, Immunity, Disease resistance, Fish.

35
36 **1. Introduction**

37 Prevalence and spread of diseases is one of the main problems in aquaculture causing great
38 economic losses in fish farms. Today, although a wide range of antibiotics are used to treat
39 diseases in fish, the use of these chemicals poses health and environmental related problems.
40 Release of antibiotics in natural water bodies can make antibiotic-resistant strains, change the
41 natural flora and fauna and also be biomagnified throughout food chain. In addition, antibiotics
42 and their derivatives can accumulate in the tissues of aquatic organisms, which can be
43 dangerous to human health as a consumer (Cabello, 2006; Romero et al., 2012; Manage, 2018).
44 Therefore, use of natural materials as alternative to chemicals can be an efficient way to
45 enhance fish immunity and to treat diseases (Dawood, 2021; Vijayaram et al., 2022). Over the
46 last decade, the use of plant-based materials and their derivatives has increased to enhance the

47 fish immune system (Harikrishnan et al., 2011; Bulfon et al., 2015; Ghafarifarsani et al.,
48 2022a). Thymol (2-isopropyl-5-ethylphenol) (TYM), is the main phenolic compound of
49 *Thymus vulgaris* essential oil, with antimicrobial properties (Kachur and Suntres, 2020), anti-
50 inflammatory (Braga et al., 2006) and antioxidant (Mastelic et al., 2008) functions. Some
51 studies have studied the effect of thymol as a dietary supplement on fish growth, hematology
52 and immunity (Ezzat Abd El-Hack et al., 2016; Alagawany et al., 2021; Kong et al., 2021;
53 Ghafarifarsani et al., 2022b,c). However, there is limited data on commercial fishes such as
54 rainbow trout with thymol.

55 The antioxidant system is a vital part of fish immunity since it makes the first line of enzymatic
56 and non-enzymatic defense against free radicals (Tan et al., 2017; Hoseinifar et al., 2020).
57 Although the enhancing effects of TYM on the antioxidant defense of vertebrates have been
58 reported in many studies (Nagoor Meeran et al., 2012; Hashemipour et al., 2013; Ezzat Abd
59 El-Hack et al., 2016; Güvenç et al., 2019; Javed et al., 2019), little is known with fish
60 (Giannenas et al., 2012; Amer et al., 2018; Abd El-Naby et al., 2020; Kong et al., 2021). TYM
61 appears to stimulate the activity of the enzymes such as catalase (CAT), superoxide dismutase
62 (SOD) glutathione (GSH) and glutathione peroxidase (GPx) (Ghafarifarsani et al., 2022b,c;
63 Yousefi et al., 2022).

64 Cytokines are signaling metabolites that regulate immune responses in fish and other
65 vertebrates by activating inflammatory and anti-inflammatory reactions (Sakai et al., 2021).
66 Inflammatory responses in fish are activated by cytokines such as interleukin-8 (IL-8) and
67 tumor necrosis factor- α (TNF- α), while other cytokines, such as transforming growth factor- β
68 (TGF- β) and IL-10, act as anti-inflammatory molecules and prevent extra activation of the
69 inflammatory responses and prompt tissue recovery processes (Wang and Secombes, 2013).
70 TYM has been shown to moderate inflammatory reactions (Braga et al., 2006; Riella et al.,
71 2012; Liang et al., 2014; Fouad et al., 2022), however, this is somewhat unknown in fish (Kong
72 et al., 2021). Apart from immune components, hematological indices are also used to evaluate
73 overall health of fish. The hematological components reflect the health status of fish with
74 nutritional and environmental factors and is usually assessed in nutrition studies (Fazio, 2019).
75 According to literatures, TYM also affects the hematology profile (Toghyani et al., 2010;
76 Saadat Shad et al., 2016), however, this effect need to be studied more in fish (Ahmadifar et
77 al., 2011; Mirzargar et al., 2021).

78 *Streptococcus iniae* is one of the most bacterial diseases of fish throughout the world, causing
79 streptococcosis disease in the rainbow trout. This disease may result in high mortality if not
80 controlled in the fish farms (Eldar and Ghittino, 1999; Lau et al., 2003; Soltani et al., 2013).

81 Considering the immunogenic properties of thymol, in the present study, the hypothesis was
82 designed that whether thymol can have immunogenic effects in rainbow trout and protect the
83 fish against bacterial infection caused by *Streptococcus iniae*? In this study, this hypothesis
84 was evaluated by feeding fish with dietary levels of thymol, determining growth and immune
85 components and resistance of fish against the bacteria.

86 **2. Materials and methods**

87 **2.1. Fish and feeding experiment**

88 Rainbow trout (n= 450) with an average weight of 35.8 ± 4.4 g (Mean \pm SD) were provided
89 from a local trout farm, Karaj, Iran, transported to lab and distributed in fifteen 500L tanks (30
90 fish/tank) with continuous aeration. After 14 days adaptation period, fish were fed diet
91 containing various levels of thymol (TYM) including 0 (control), 1, 1.5, 2 and 2.5 g/kg diet in
92 three replicates for 60 days (Amer et al., 2018). The thymol crystals was provided from Oxford

93 Laboratory Company, Mumbai India with purity of 99%. Feeding was done four times a day
94 (6:00 AM, 12:00 noon, 6:00 PM, 12:00 midnight) to satisfaction. The experimental diets were
95 prepared by mixing TYM with a basal diet (Table 1; crude protein: 40%, crude lipid: 15%,
96 Ash: 9%, crude fiber: 3%, moisture: 11%, phosphorus: 1.2%; energy content: 21.2 MJ/kg)
97 according to (Amer et al., 2018). The chemical composition of the basal diet in this regard,
98 TYM was mixed with fish oil and then mixed with previously powdered basal diet. The diets
99 were prepared in pelleted form (4 mm) and stored in plastic bags at 0 °C until usage (Amer et
100 al., 2018). During the feeding period, Fish were under natural photoperiod, continuous flow of
101 water was kept in tanks and water quality parameters (Mean ± SD) checked for temperature:
102 15.6 ± 0.3 °C (Zeal thermometer, UK), oxygen: > 6 mg/l (Portable Oxygen Meter, Hanna,
103 HI9146, UK), total ammonia: <0.01 mg/l (Hi-700 Ammonia Low Range Colorimeter –
104 Checker, Hanna instruments CO., UK), pH: 7.3 ± 0.1 (pH meter, Hanna-HI 98128, Hanna
105 instruments CO., UK).

106 **2.2. Growth and survival parameters**

107 After feeding period, the growth indices were estimated as follow:

108 Weight gain percent (%) = [final weight (g)–initial weight (g)] / [initial weight (g)] × 100

109 Specific growth rate (SGR) (%/d) = ([ln final wt(g) – ln initial wt(g)] / days) × 100

110 Feed conversion rate (FCR) = total feed given(g) / weight gain(g)

111 Survival rate (SR) (%) = (final numbers / initial numbers) × 100

112 **2.3. Body composition analysis**

113 The fish body composition of fish were determined based on the methods recommended by
114 Association of Official Analytical Chemists (AOAC, 2000). Crude protein (CP), lipid,
115 moisture and ash content were determined following specific procedure identification numbers
116 of AOAC 923.03 for ash, AOAC 920.87 for protein, AOAC 945.38F; 920.39C for fat and
117 AOAC 925.09 for moisture. Six fish per treatment were randomly taken and then grounded by
118 a grinder (MFW68640, Bosch, Germany). The moisture content was measured after oven-
119 drying at 105 °C to reach a constant weight. The ash content was measured through incineration
120 of fish carcass in a muffle furnace at 550 °C for 24 h. The crude protein (N × 6.25) and lipid
121 content were determined by the Kjeldahl and Soxhlet methods respectively.

122 **2.4. Blood and tissue sampling**

123 The samples of blood, mucus and tissue were collected after the feeding period. Fish were first
124 starved for 24 h and then 15 fish randomly captured from each tank. The mucus collection was
125 conducted by putting fish in nylon bags according to Ghafarifarsani et al. (2021a). The blood
126 samples were taken from caudal vein by 2.5 ml heparinized syringe after sedation of the fish
127 with 200 mg/l clove powder (Tukmechi and Bandboni, 2014). Then, the plasma was obtained
128 through centrifugation at 13000 g for 10 min. Also, the blood slides were prepared for
129 hematological examinations according to Kokou et al. (2015). Liver samples were collected
130 after dissecting the fish. Finally, the plasma, mucus and liver samples were kept in liquid
131 nitrogen (- 196 °C) for further biochemical analysis.

132 **2.5. Immune parameters**

133 The plasma and mucosal lysozyme activity was measured using lyophilized *Micrococcus*
134 *luteus* according to Ellis (1990). The total immunoglobulin (Ig) concentration in plasma and
135 mucus was assayed by polyethylene glycol method (Siwicki and Anderson, 1993). The plasma
136 alternative complement activity (C3) was determined by measuring the *haemolysis* rate of
137 rabbit red blood cells (Ortuno et al. 2001). The blood bactericidal activity was determined
138 against *Streptococcus iniae* (OD: 0.5 at 546 nm) inside a bacterial suspension and following

139 calculating bacterial colony forming unites (CFUs) on nutrient agar plates after 24 h incubation
140 at 35 °C (Rao et al. 2006).

141 The total protein was determined according to Bradford (1976) using a Sigma-Aldrich Protein
142 Assay Kit. Protease activity in mucus was assayed by the Azocasein hydrolysis procedure, as
143 described by Ross et al. (2000). Alkaline phosphatase (ALP) activity in mucus was assayed
144 colorimetrically at 405 nm by assay kit (Sigma-Aldrich CO, USA) based on the hydrolysis of
145 p-nitrophenol phosphate to p-nitrophenol (Ross et al. 2000).

146 **2.6. Liver antioxidant enzymes and lipid peroxidation**

147 The activity of antioxidant enzymes in liver were assayed using commercial assay kits,
148 according to manufacturer's instructions (Sigma-Aldrich CO, USA). Catalase activity was
149 assayed through the reduction of hydrogen peroxide at 240 nm (Tocher et al., 2002).
150 Superoxide dismutase (SOD) activity was assayed through inhibiting the oxygen-dependent
151 oxidation of adrenaline (epinephrine) to adrenochrome using xanthine oxidase plus xanthine
152 (Panchenko et al., 1975). Glutathione peroxidase (GPx) activity was determined through
153 measuring the rate of NADPH oxidation at 340 nm under glutathione reductase action (Bell et
154 al. 1985). The lipid peroxidation was measured spectrophotometrically at 532 nm upon reaction
155 of malondialdehyde (MDA) with thiobarbituric acid (Utley et al. 1967).

156 **2.7. Digestive enzymes**

157 The amylase activity was measured colorimetrically at 600 nm using a 2% starch solution (as
158 substrate) in 0.1 M citrate phosphate buffer (Robyt and Whelan, 1968). Lipase enzyme activity
159 was determined at 405 nm upon hydrolysis of polyphenol myristate (as substrate) according to
160 Gawlicka et al. (2000). Protease activity was determined at 440 nm using azocasein as substrate
161 (Garcia-Carreno, 1992).

162 **2.8. Haematological parameters**

163 The total number of red and white blood cells was done using a haemocytometer slide under
164 optical microscope at 400x magnification (Rothmann et al., 2000). Haematocrit (Hct) was
165 determined by the microhematocrit method (Blaxhall and Daisley, 1973). The haemoglobin
166 (Hb) content was measured by the cyanohaemoglobin method (Rawling et al., 2012). The
167 haematological indices [mean corpuscular volume (MCV), and corpuscular hemoglobin
168 concentration (MCHC), mean corpuscular hemoglobin (MCH)] were estimated according to
169 following equations (Blaxhall and Daisley, 1973):

170 $MCHC = Hb \times 10 / Hct$ $MCV = Hct \times 10 / RBC$ (million) $MCH = Hb \times 10 / RBC$ (million)

171 **2.9. Gene expression assay**

172 **2.9.1. RNA extraction**

173 The total RNA content of liver tissue was extracted by acid guanidinium thiocyanate-
174 phenolchloroform procedure according to Chomczynski and Sacchi, (2006) with some
175 modifications. The quantity and quality of the extracted RNA was estimated by determining
176 the absorbance at 260 nm with a Nanodrop spectrophotometer (NanoDrop technologies;
177 Wilmington, DE, USA). Also, the quality of RNA was evaluated by electrophoresis (Kwasek
178 et al. 2014).

179 **2.9.2. Reverse transcription polymerase chain reaction (RT-PCR)**

180 The extracted RNA (1 µg) was used for making first-strand cDNAs by a Fermentas cDNA
181 synthesis Kit for RT-PCR (Reverse transcription polymerase chain reaction). All procedures

182 were conducted according to manufacturer's instructions by an iCycler (BioRad). The RT-PCR
183 primers were synthesized based on the sequences of DNA from Gen Bank (Table 2) using Gene
184 Runner (version 6.) software. The β -actine gene was used as reference gene to estimate the
185 gene expressions. The fold changes in the gene [C3 (complement), lysozyme, and Ig
186 (immunoglobulin), IL-8 (Interlukin-8), TNF- α (Tumor necrosis factor- α) was measured by the
187 $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen 2001). Finally, the RT-PCR data were analyzed using
188 iQ5 optical system software version 2.1 (Bio-Rad).

189 **2.10. Bacterial challenge**

190 After 60 days feeding period, fish (10 fish/tank) were exposed to *Streptococcus iniae* by
191 injection. The bacterium, originally isolated from diseased rainbow trout, and cultivated on
192 agar medium for 24 h at 36 °C. The dose of injection dose was selected in a previous experiment
193 by calculation of seven day LD₅₀ (lethal dose 50), which it was 1.6×10^7 cells/ml phosphate
194 buffered saline (PBS).

195 The cumulative mortality of the fish was recorded daily over ten days challenge. I dead fish,
196 the bacterial infection was confirmed by cultivation of liver tissue extract growth medium using
197 conventional methods.

198 **2.11. Statistical analysis**

199 The analysis of data was done using SPSS software (version 16). After evaluation of data
200 normality by Kolmogorov-Smirnov test, data were subjected to One-way analysis of variance
201 to investigate the significants. Finally, the comparison of means was conducted using Tukey
202 test ($P < 0.05$). Also, the growth responses to dietary TYM were evaluated through polynomial
203 regression.

204 **3. Results**

205 **3.1. Growth parameters**

206 The supplementation of fish with 1.5-2 g TYM significantly improved the growth parameters
207 (final weight, WG (%) and SGR) in the fish compared to other diets (Table 3, $P < 0.05$). The
208 FCR values significantly decreased in the treatments 2-2.5 g TYM compared to other
209 experimental groups (Table 3, $P < 0.05$). There were no significant differences in growth
210 parameters between control and the treatment, 1 g TYM (Table 3, $P > 0.05$). Also, the dose-
211 response model (Fig. 1) showed the potential response of the fish to graded levels of dietary
212 TYM (Fig. 1). In this regard, the lowest FCR was observed at 1.89 % of TYM.

213 **3.2. Body composition analysis**

214 The protein (%) content of the fish body significantly increased in fish fed 2-2.5 g TYM
215 compared to compared to other diets (Table 4, $P < 0.05$). However, the lipid, ash and moisture
216 content had no significant differences between the experimental groups after feeding
217 experiment (Table 4, $P > 0.05$).

218 **3.3. Digestive enzymes**

219 The protease and lipase activities showed significant increases in fish fed 1.5-2.5 g TYM
220 compared to control (Table 5, $P < 0.05$). The highest activity of these enzymes were observed
221 in the group fed 2.5 g TYM (Table 5, $P < 0.05$). The amylase activity showed no significant
222 differences between all groups after feeding period (Table 5, $P > 0.05$).

223 **3.4. Plasma and mucus Immune parameters**

224 The immune components in plasma (C3, total Ig, lysozyme activity, bactericidal activity and
225 total protein) and in mucus (ALP), protease activity, lysozyme activity, bactericidal activity
226 and total protein) significantly increased in response to 1.5-2.5 g TYM compared to other diets
227 (Table 6, $P < 0.05$). The highest values of immune components was observed mostly in fish fed

228 2-2.5 g TYM (Table 6, P<0.05). There were no significant differences in immune components
229 of plasma and mucus between control and fish fed 1 g TYM (Table 6, P>0.05).

230 **3.5. Hematology**

231 The hematology of the fish altered in response to dietary TYM (Table 7, P<0.05). The values
232 of corpuscular hemoglobin concentration (MCHC), hemoglobin (Hb), red blood cell (RBC),
233 hematocrit (Hct) and white blood cell (WBC) significantly increased in fish fed 2-2.5 g TYM
234 compared to other diets (Table 7, P<0.05). The values of MCV significantly increased in
235 response to 1.5 g TYM compared to other diets (Table 7, P<0.05). In addition, MCV
236 significantly decreased in fish fed 2-2.5 g TYM (Table 7, P<0.05). There were no significant
237 differences in MCH between all groups (Table 7, P>0.05).

238 **3.6. Liver antioxidant enzymes**

239 The liver antioxidant enzymes showed significant changes in response to TYM (Table 8,
240 P<0.05). In comparison with control group, the CAT activity with maximum activity in the
241 treatments, 2 and 2.5 g TYM significantly increased in all TYM supplemented fish, while GPx
242 and SOD elevated only in fish fed 1.5-2.5 g TYM (Table 8, P<0.05). There were no significant
243 differences in GPx and SOD activity between control and fish supplemented with 1 g TYM
244 (Table 8, P>0.05).

245 **3.7. Lipid peroxidation index**

246 The levels of MDA significantly decreased in fish fed 2-2.5 g TYM compared to other diets
247 (Fig. 2, P<0.05). There were no significant differences in MDA levels between the control with
248 fish fed 1-1.5 g TYM (Fig. 2, P>0.05).

249 **3.8. Gene expression**

250 The supplementation of fish with 1.5-2.5 g TYM significantly increased the expression of C3,
251 lysozyme and Ig genes compared to control (Fig. 3, P<0.05). The expression of inflammatory-
252 related genes (TNF- α and IL-8) significantly decreased in fish supplemented with 1.5-2.5 g
253 TYM compared to other groups (Fig. 4, P<0.05).

254 **3.9. Bacterial challenge**

255 After 10 days bacterial challenge, the fish supplemented with 2-2.5 g TYM significantly
256 showed lower mortality rate compared to other experimental groups (Fig. 5, P<0.05). The
257 highest mortality rate were observed in control and fish fed 1 g TYM (Fig. 5, P<0.05).

258 **4. Discussion**

259 Antibiotics are widely used to control bacterial pathogens in aquaculture. Although the
260 bacterial infections can be controlled by antibiotics, their continued use weakens the immune
261 system and may creates bacterial resistance strains (Manage, 2018). Introducing an
262 environmentally and human health friendly alternative to antibiotics could be important. In this
263 way, herbal products and their derivatives have been shown to have high performance
264 (Syahidah et al., 2015). In the present study, we used thymol (TYM) as dietary supplement for
265 rainbow trout. The growth performance improved in response to TYM, as the FW, WG % and
266 SGR increased in fish fed 1.5-2.5 ml TYM, and FCR decreased in those fed 2-2.5 g TYM. In
267 addition, the body protein content and the activity of digestive enzymes, lipase and protease
268 increased in response to 1.5-2.5 g TYM. In line with our results, Morselli et al. (2020b)
269 observed better growth in grass carp, *Ctenopharyngodon idella* supplemented with 100 mg
270 thymol/kg diet. In the rainbow trout, the growth performance improved in fish fed diets
271 containing thymol-carvacrol (Ahmadifar et al., 2011). Similar results were found in same fish,
272 where the fish fed 6 g/kg thymol showed better feed efficiency compared to non-supplemented
273 individuals (Giannenas et al., 2012). In the study of Abd El-Naby et al. (2020), thymol alone
274 or in combination with chitosan nanoparticles significantly improved growth performance,

275 feed and protein utilization in the Nile tilapia, *Oreochromis niloticus*. However, Hoseini and
276 Yousefi (2019) did not observe any changes in the growth performance of rainbow trout after
277 a 60-day feeding with a thymol-containing diet (5, 10, 20 g/kg diet) between experimental
278 treatments. Similar results were observed in the Nile tilapia, where 500 ppm thymol had no
279 effect on the fish growth (Aanyu et al., 2018). Therefore, the effect of thymol on growth may
280 be different depending on dietary levels, fish species and experiment duration and conditions.
281 In this study, the improved growth performance in the supplemented fish may be due to
282 prompting effects TYM on digestive enzymes, feed utilization and health status of the fish, as
283 the digestive enzyme activity, the immune and antioxidant components elevated in the fish fed
284 1.5-2.5 g TYM. According to results, dietary TYM (mostly 1.5-2.5 g/kg diet), prompted the
285 antioxidant and immune system by stimulating the liver antioxidant enzymes (GPx, SOD,
286 CAT) and immune components of blood (C3, total Ig, protein content, lysozyme and
287 bactericidal activities) and mucus (Ig, protein content, lysozyme, protease, ALP and
288 bactericidal activities). Additionally, TYM at dietary levels of 1.5-2.5 g/kg diet up-regulated
289 the expression of immune-related genes (C3, lysozyme and Ig). As a component of humoral
290 defense, Ig immunizes fish through opsonization of pathogens and also neutralization of toxic
291 molecules (Salinas et al., 2011; Mashoof and Criscitiello, 2016). Furthermore, the proteins of
292 complement system are involved in both non-specific and specific immunity by opsonization
293 of pathogens and activating inflammatory reactions (Boshra et al., 2006). The blood protein
294 content can also be an indicator of the immune status of fish, because it includes antibodies and
295 albumin (Corbel, 1975). Alkaline phosphatase is a mucosal enzyme with antibacterial and
296 hydrolytic activities. Therefore, the elevated levels of ALP in the TYM-supplemented fish may
297 indicate an improvement in immune status (Hoseinifar et al., 2018). Proteases are a group of
298 fish mucus enzymes with catalytic function on the peptidoglycan layer of bacteria (Dash et al.,
299 2018). Enzymatic and non-enzymatic components of antioxidant system are involved in fish
300 immunity through neutralizing free radicals, reducing oxidative stress (Hoseinifar et al., 2020).
301 SOD catalyzes the dismutation of the superoxide anion (O_2^-) to molecular oxygen and H_2O_2 .
302 Finally, the generated H_2O_2 is eliminated by CAT and GPx action (Gupta et al., 2018).
303 The results of the present study were in consist with other studies. The supplementation of
304 seabream, *Sparus aurata*, with TYM decreased the growth capacity of pathogenic bacteria in
305 the mucus (Firmino et al., 2021). Dietary TYM enhanced the immune system of Nile tilapia by
306 reducing oxidative stress and by increasing lysozyme activity and Ig levels in the serum (Amer
307 et al., 2018). Giannenas et al. (2012) indicated that dietary thymol alone or in combination with
308 carvacrol is capable to ameliorate oxidative stress in the rainbow trout and stimulate liver CAT
309 activity and lysozyme and complement activities in serum. The supplementation of Northern
310 snakehead, *Channa argus*, with 300 and 450 mg/kg thymol significantly ameliorated oxidative
311 stress and stimulated immune components (Ig levels, serum acid phosphatase, ALP,
312 complement and lysozyme activities) in serum, enhanced the activity of antioxidant enzymes
313 (GPx, SOD, CAT) and increased the resistance to *Aeromonas veronii* infection (Kong et al.,
314 2021). In the Nile tilapia, a dietary combination of thymol and chitosan nanoparticles stimulated
315 the CAT activity in liver and kidney tissues (Abd El-Naby et al., 2020). In grass carp of
316 100-300 mg/kg feed thymol enhanced the activity of SOD and GPx in liver and ameliorated
317 oxidative stress (Morselli et al., 2020). The results of the present study were in line with studies,
318 suggesting a potential immune-promoting function for TYM, which may be associated with its
319 stimulating effects on humoral innate immune components, immune-related gene expressions
320 and antioxidant defense system.

321 Fish hematology profile is known as an indicator of fish overall health status, which may
322 change in relation to environmental, nutritional and physiological conditions (Fazio, 2019).

323 In the present study, dietary TYM increased the values of RBC, Hct, and Hb, which may
324 enhance the capacity of oxygen transportation in blood and thus improve the fish health and
325 welfare. In addition, the WBC levels elevated in TYM supplemented fish, suggesting an
326 immunogenic function for TYM, as previously reported by Ahmadifar et al. (2011). In
327 agreement with our results, Abd El-Naby et al. (2020) reported increases in lymphocytes count,
328 Hct, MCV, and MCHC in the Nile tilapia supplemented with TYM.

329 Cytokines are the signaling and regulating molecules of both innate and fish immune system
330 involved in inflammatory reactions and phagocytic activities (Wang and Secombes, 2013;
331 Chauhan et al., 2021; Ghafarifarsani et al., 2021b). In this study, TYM at dietary levels of 2-
332 2.5 g/kg diet down-regulated the expression of inflammatory-related genes (*TNF- α* and *IL-8*),
333 which this result may be associated with the anti-inflammatory function of TYM, as
334 demonstrated previously in other studies (Braga et al., 2006; Riella et al., 2012). Kong et al.
335 (2021) observed the up-regulation of anti-inflammatory (*IL-10* and *TGF- β*) and down-
336 regulation of inflammatory (*HSP70*, *TNF- α* , *IL-1 β* and *IL-8*) genes, following supplementation
337 of the northern snakehead with 300-400 mg thymol/kg feed, which attributed to anti-
338 inflammatory function of thymol.

339 In this study, the mortality rate was lower in fish fed 2-2.5 g TYM compared to other groups
340 after the bacterial challenge, which clearly indicates the antibacterial properties of TYM. The
341 antibacterial properties of thymol have been previously reported in an *in vitro* study by Heo et
342 al. (2012), where TYM efficiently inhibited the growth of the fish pathogens including *Vibrio*
343 *vulnificus*, *V. parahaemolyticus* and *V. Anguillarum*, *Aeromonas salmonicida*, *A. Hydrophila*
344 and *Edwardsiella tarda*. Also, in the study of Morselli et al. (2020), the supplementation of the
345 grass carp with thymol significantly increased the survival rate of the fish by 62.5 % following
346 challenge with *Aeromonas hydrophila*.

347 **5. Conclusion**

348 In conclusion, the results of the present study suggest an optimum dietary level of 1.5-2.5 g/kg
349 diet for TYM, improving the growth performance, immunity and resistance against
350 *Streptococcus iniae* infection in the rainbow trout.

351 **Ethical Approval**

352 In this study, all stages of sampling and manipulation of animals have been performed in
353 accordance with ethical standards.

354 **Data Availability Statements**

355 The datasets generated during and/or analysed during the current study are available from the
356 corresponding author on reasonable request.

357 **Funding**

358 There is no government or organizational fund for this work.

359 **Author's contributions**

360 All the authors of this article have made important contributions to testing, collecting data,
361 analyzing results, and writing the article.

362 **Conflict of interest**

363 The authors have no conflict of interest

364 **Acknowledgments**

365 The authors would like to thank all those who help us to do the research.

367 **References**

- 368 Aanyu, M., Betancor, M. B., & Monroig, O. (2018). Effects of dietary limonene and
369 thymol on the growth and nutritional physiology of Nile tilapia (*Oreochromis*
370 *niloticus*). *Aquaculture*, *488*, 217-226.
- 371 Rawling, M. D., Merrifield, D. L., Snellgrove, D. L., Kühlwein, H., Adams, A., &
372 Davies, S. J. (2012). Haemato-immunological and growth response of mirror carp (*Cyprinus*
373 *carpio*) fed a tropical earthworm meal in experimental diets. *Fish & Shellfish*
374 *Immunology*, *32*(6), 1002-1007.
- 375 Alagawany, M., Farag, M. R., Abdelnour, S. A., & Elnesr, S. S. (2021). A review on
376 the beneficial effect of thymol on health and production of fish. *Reviews in Aquaculture*, *13*(1),
377 632-641.
- 378 Kong, Y. D., Li, M., Xia, C. G., Zhao, J., Niu, X. T., Shan, X. F., & Wang, G. Q. (2021).
379 The optimum thymol requirement in diets of *Channa argus*: effects on growth, antioxidant
380 capability, immune response and disease resistance. *Aquaculture Nutrition*, *27*(3), 712-722.
- 381 Abd El-Naby, A. S., Al-Sagheer, A. A., Negm, S. S., & Naiel, M. A. (2020). Dietary
382 combination of chitosan nanoparticle and thymol affects feed utilization, digestive enzymes,
383 antioxidant status, and intestinal morphology of *Oreochromis niloticus*. *Aquaculture*, *515*,
384 734577.
- 385 Ahmadifar, E., Falahatkar, B., & Akrami, R. (2011). Effects of dietary thymol-
386 carvacrol on growth performance, hematological parameters and tissue composition of juvenile
387 rainbow trout, *Oncorhynchus mykiss*. *Journal of Applied Ichthyology*, *27*(4), 1057-1060.
- 388 Amer, S. A., Metwally, A. E., & Ahmed, S. A. (2018). The influence of dietary
389 supplementation of cinnamaldehyde and thymol on the growth performance, immunity and
390 antioxidant status of monosex Nile tilapia fingerlings (*Oreochromis niloticus*). *The Egyptian*
391 *Journal of Aquatic Research*, *44*(3), 251-256.
- 392 AOAC 2000. Official Methods of Analysis, 17th Edition. Washington: Association of
393 Official Analytical Chemists
- 394 Bell, J. G., Cowey, C. B., Adron, J. W., & Shanks, A. M. (1985). Some effects of
395 vitamin E and selenium deprivation on tissue enzyme levels and indices of tissue peroxidation
396 in rainbow trout (*Salmo gairdneri*). *British Journal of Nutrition*, *53*(1), 149-157.
- 397 Blaxhall, P. C., & Daisley, K. W. (1973). Routine haematological methods for use with
398 fish blood. *Journal of fish biology*, *5*(6), 771-781.
- 399 Boshra, H., Li, J., & Sunyer, J. O. (2006). Recent advances on the complement system
400 of teleost fish. *Fish & shellfish immunology*, *20*(2), 239-262.
- 401 Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram
402 quantities of protein utilizing the principle of protein-dye binding. *Analytical*
403 *biochemistry*, *72*(1-2), 248-254.
- 404 Braga, P. C., Dal Sasso, M., Culici, M., Bianchi, T., Bordoni, L., & Marabini, L. (2006).
405 Anti-inflammatory activity of thymol: inhibitory effect on the release of human neutrophil
406 elastase. *Pharmacology*, *77*(3), 130-136.
- 407 Bulfon, C., Volpatti, D., & Galeotti, M. (2015). Current research on the use of plant-
408 derived products in farmed fish. *Aquaculture Research*, *46*(3), 513-551.

- 409 Cabello, F. C. (2006). Heavy use of prophylactic antibiotics in aquaculture: a growing
410 problem for human and animal health and for the environment. *Environmental*
411 *microbiology*, 8(7), 1137-1144.
- 412 Chauhan, P., Nair, A., Patidar, A., Dandapat, J., Sarkar, A., & Saha, B. (2021). A primer
413 on cytokines. *Cytokine*, 145, 155458.
- 414 Chomczynski, P., & Sacchi, N. (2006). The single-step method of RNA isolation by
415 acid guanidinium thiocyanate–phenol–chloroform extraction: twenty-something years
416 on. *Nature protocols*, 1(2), 581-585.
- 417 Corbel, M. J. (1975). The immune response in fish: a review. *Journal of Fish*
418 *Biology*, 7(4), 539-563.
- 419 Dash, S., Das, S. K., Samal, J., & Thatoi, H. N. (2018). Epidermal mucus, a major
420 determinant in fish health: a review. *Iranian Journal of Veterinary Research*, 19(2), 72.
- 421 Dawood, M. A. (2021). Nutritional immunity of fish intestines: Important insights for
422 sustainable aquaculture. *Reviews in Aquaculture*, 13(1), 642-663.
- 423 Eldar, A. A., & Ghittino, C. (1999). Lactococcus garvieae and Streptococcus iniae
424 infections in rainbow trout Oncorhynchus mykiss: similar, but different diseases. *Diseases of*
425 *aquatic organisms*, 36(3), 227-231.
- 426 Ellis, AE I. "Lysozyme assays." *Techniques in fish immunology* 1 (1990): 101-103.
- 427 Ezzat Abd El-Hack, M., Alagawany, M., Ragab Farag, M., Tiwari, R., Karthik, K.,
428 Dhama, K., ... & Adel, M. (2016). Beneficial impacts of thymol essential oil on health and
429 production of animals, fish and poultry: a review. *Journal of Essential Oil Research*, 28(5),
430 365-382.
- 431 Fazio, F. (2019). Fish hematology analysis as an important tool of aquaculture: a
432 review. *Aquaculture*, 500, 237-242.
- 433 Firmino, J. P., Fernández-Alacid, L., Vallejos-Vidal, E., Salomón, R., Sanahuja, I.,
434 Tort, L., ... & Gisbert, E. (2021). Carvacrol, Thymol, and Garlic Essential Oil Promote Skin
435 Innate Immunity in Gilthead Seabream (*Sparus aurata*) Through the Multifactorial Modulation
436 of the Secretory Pathway and Enhancement of Mucus Protective Capacity. *Frontiers in*
437 *Immunology*, 12, 559.
- 438 Fouad, A. A., Moussa, N. A., Kareem, M. M. A., Akl, U. I., Abdelghany, M. I., &
439 Abdel-Aziz, A. M. (2022). Thymol exerts antioxidant, anti-inflammatory, and anti-apoptotic
440 protective effects against gentamicin nephrotoxicity in rats. *Pharmacia*, 69(1), 181-186.
- 441 García-Carreño, F. L. (1992). Protease inhibition in theory and practice. *Biotechnology*
442 *Education*, 3(4), 145-150.
- 443 Gawlicka, A., Parent, B., Horn, M. H., Ross, N., Opstad, I., & Torrissen, O. J. (2000).
444 Activity of digestive enzymes in yolk-sac larvae of Atlantic halibut (*Hippoglossus*
445 *hippoglossus*): indication of readiness for first feeding. *Aquaculture*, 184(3-4), 303-314.
- 446 Ghafarifarsani, H., Rashidian, G., Bagheri, T., Hoseinifar, S. H., & Van Doan, H.
447 (2021a). Study on growth enhancement and the protective effects of dietary prebiotic inulin on
448 immunity responses of rainbow trout fry infected with. *Annals of Animal Science*, 21(2), 543-
449 559.
- 450 Ghafarifarsani, H., Kachuei, R., & Imani, A. (2021b). Dietary supplementation of
451 garden thyme essential oil ameliorated the deteriorative effects of aflatoxin B₁ on growth

452 performance and intestinal inflammatory status of rainbow trout (*Oncorhynchus*
453 *mykiss*). *Aquaculture*, 531, 735928.

454 Ghafarifarsani, H., Hoseinifar, S. H., Aftabgard, M., & Van Doan, H. (2022a). The
455 improving role of savory (*Satureja hortensis*) essential oil for Caspian roach (*Rutilus caspicus*)
456 fry: Growth, haematological, immunological, and antioxidant parameters and resistance to
457 salinity stress. *Aquaculture*, 548, 737653.

458 Ghafarifarsani, H., Hoseinifar, S. H., Javahery, S., & Van Doan, H. (2022b). Effects of
459 dietary vitamin C, thyme essential oil, and quercetin on the immunological and antioxidant
460 status of common carp (*Cyprinus carpio*). *Aquaculture*, 738053.

461 Ghafarifarsani, H., Hoseinifar, S. H., Sheikhlar, A., Raissy, M., Chaharmahali, F. H.,
462 Maneepitaksanti, W., ... & Van Doan, H. (2022c). The Effects of Dietary Thyme Oil (*Thymus*
463 *vulgaris*) Essential Oils for Common Carp (*Cyprinus carpio*): Growth Performance, Digestive
464 Enzyme Activity, Antioxidant Defense, Tissue and Mucus Immune Parameters, and Resistance
465 against *Aeromonas hydrophila*. *Aquaculture Nutrition*, 2022, Article ID 7942506.
466 <https://doi.org/10.1155/2022/7942506>

467 Giannenas, I., Triantafyllou, E., Stavrakakis, S., Margaroni, M., Mavridis, S., Steiner,
468 T., & Karagouni, E. (2012). Assessment of dietary supplementation with carvacrol or thymol
469 containing feed additives on performance, intestinal microbiota and antioxidant status of
470 rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 350, 26-32.

471 Gupta, D. K., Palma, J. M., & Corpas, F. J. (Eds.). (2018). *Antioxidants and antioxidant*
472 *enzymes in higher plants*. Springer.

473 Güvenç, M., Cellat, M., Gökçek, İ., Yavaş, İ., & Yurdagül Özsoy, Ş. (2019). Effects of
474 thymol and carvacrol on sperm quality and oxidant/antioxidant balance in rats. *Archives of*
475 *Physiology and Biochemistry*, 125(5), 396-403.

476 Harikrishnan, R., Balasundaram, C., & Heo, M. S. (2011). Impact of plant products on
477 innate and adaptive immune system of cultured finfish and shellfish. *Aquaculture*, 317(1-4), 1-
478 15.

479 Hashemipour, H., Kermanshahi, H., Golian, A., & Veldkamp, T. (2013). Effect of
480 thymol and carvacrol feed supplementation on performance, antioxidant enzyme activities,
481 fatty acid composition, digestive enzyme activities, and immune response in broiler
482 chickens. *Poultry science*, 92(8), 2059-2069.

483 Heo, G., Kim, C., Park, S., Zoysa, M. D., & Shin, G. (2012). Antimicrobial activity of
484 thymol against pathogenic Gram-negative bacteria of fishes. *Philippine Journal of Veterinary*
485 *Medicine*, 49(2).

486 Hoseini, S. M., & Yousefi, M. (2019). Beneficial effects of thyme (*Thymus vulgaris*)
487 extract on oxytetracycline-induced stress response, immunosuppression, oxidative stress and
488 enzymatic changes in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture nutrition*, 25(2),
489 298-309.

490 Hoseinifar, S. H., Yousefi, S., Capillo, G., Paknejad, H., Khalili, M., Tabarraei, A., ...
491 & Faggio, C. (2018). Mucosal immune parameters, immune and antioxidant defence related
492 genes expression and growth performance of zebrafish (*Danio rerio*) fed on *Gracilaria gracilis*
493 powder. *Fish & shellfish immunology*, 83, 232-237.

494 Hoseinifar, S. H., Yousefi, S., Van Doan, H., Ashouri, G., Gioacchini, G., Maradonna,
495 F., & Carnevali, O. (2020). Oxidative stress and antioxidant defense in fish: the implications
496 of probiotic, prebiotic, and synbiotics. *Reviews in Fisheries Science & Aquaculture*, 29(2),
497 198-217.

- 498 Javed, H., Azimullah, S., Meeran, M. F., Ansari, S. A., & Ojha, S. (2019).
 499 Neuroprotective effects of thymol, a dietary monoterpene against dopaminergic
 500 neurodegeneration in rotenone-induced rat model of Parkinson's disease. *International journal*
 501 *of molecular sciences*, 20(7), 1538.
- 502 Kachur, K., & Suntres, Z. (2020). The antibacterial properties of phenolic isomers,
 503 carvacrol and thymol. *Critical Reviews in Food Science and Nutrition*, 60(18), 3042-3053.
- 504 Kokou, F., Sarropoulou, E., Cotou, E., Rigos, G., Henry, M., Alexis, M., & Kentouri,
 505 M. (2015). Effects of fish meal replacement by a soybean protein on growth, histology, selected
 506 immune and oxidative status markers of gilthead sea bream, Sparus aurata. *Journal of the*
 507 *World Aquaculture Society*, 46(2), 115-128.
- 508 Kong, Y. D., Li, M., Xia, C. G., Zhao, J., Niu, X. T., Shan, X. F., & Wang, G. Q. (2021).
 509 The optimum thymol requirement in diets of Channa argus: effects on growth, antioxidant
 510 capability, immune response and disease resistance. *Aquaculture Nutrition*, 27(3), 712-722.
- 511 Kwasek, K., Terova, G., Lee, B. J., Bossi, E., Saroglia, M., & Dabrowski, K. (2014).
 512 Dietary methionine supplementation alters the expression of genes involved in methionine
 513 metabolism in salmonids. *Aquaculture*, 433, 223-228.
- 514 Lau, S. K., Woo, P. C., Tse, H., Leung, K. W., Wong, S. S., & Yuen, K. Y. (2003).
 515 Invasive streptococcus iniae infections outside north america. *Journal of clinical*
 516 *microbiology*, 41(3), 1004-1009.
- 517 Liang, D., Li, F., Fu, Y., Cao, Y., Song, X., Wang, T., ... & Zhang, N. (2014). Thymol
 518 inhibits LPS-stimulated inflammatory response via down-regulation of NF- κ B and MAPK
 519 signaling pathways in mouse mammary epithelial cells. *Inflammation*, 37(1), 214-222.
- 520 Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data
 521 using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *methods*, 25(4), 402-408.
- 522 Manage, P. M. (2018). Heavy use of antibiotics in aquaculture; emerging human and
 523 animal health problems—a review.
- 524 Mashoof, S., & Criscitiello, M. F. (2016). Fish immunoglobulins. *Biology*, 5(4), 45.
- 525 Mastelic, J., Jerkovic, I., Blažević, I., Poljak-Blaži, M., Borović, S., Ivančić-Baće, I., ...
 526 & Müller, N. (2008). Comparative study on the antioxidant and biological activities of
 527 carvacrol, thymol, and eugenol derivatives. *Journal of agricultural and food chemistry*, 56(11),
 528 3989-3996.
- 529 Mirzargar, S. S., Taheri Mirghaed, A., Hoseini, S. M., Ghelichpour, M., Shahnazi, M.,
 530 & Yousefi, M. (2021). Effects of adding Thymol to water on hematological and immunological
 531 parameters of common carp (Cyprinus carpio) during transportation. *Aquaculture*
 532 *Sciences*, 9(1), 27-34.
- 533 Morselli, M. B., Baldissera, M. D., Souza, C. F., Reis, J. H., Baldisserotto, B., Sousa,
 534 A. A., ... & Da Silva, A. S. (2020a). Effects of thymol supplementation on performance,
 535 mortality and branchial energetic metabolism in grass carp experimentally infected by
 536 *Aeromonas hydrophila*. *Microbial Pathogenesis*, 139, 103915.
- 537 Morselli, M. B., Reis, J. H., Baldissera, M. D., Souza, C. F., Baldisserotto, B., Petrolli,
 538 T. G., ... & Da Silva, A. S. (2020b). Benefits of thymol supplementation on performance, the
 539 hepatic antioxidant system, and energetic metabolism in grass carp. *Fish Physiology and*
 540 *Biochemistry*, 46(1), 305-314.

- 541 Nagoor Meeran, M. F., & Stanely Mainzen Prince, P. (2012). Protective effects of
542 thymol on altered plasma lipid peroxidation and nonenzymic antioxidants in isoproterenol-
543 induced myocardial infarcted rats. *Journal of Biochemical and Molecular Toxicology*, 26(9),
544 368-373.
- 545 Ortuno, J., Esteban, M. A., & Meseguer, J. (2001). Effects of short-term crowding stress
546 on the gilthead seabream (*Sparus aurata* L.) innate immune response. *Fish & shellfish*
547 *immunology*, 11(2), 187-197.
- 548 Panchenko, L. F., Brusov, O. S., Gerasimov, A. M., & Loktaeva, T. D. (1975).
549 Intramitochondrial localization and release of rat liver superoxide dismutase. *Febs*
550 *Letters*, 55(1-2), 84-87.
- 551 Rao, Y. V., Das, B. K., Jyotirmayee, P., & Chakrabarti, R. (2006). Effect of
552 *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas*
553 *hydrophila*. *Fish & Shellfish Immunology*, 20(3), 263-273.
- 554 Riella, K. R., Marinho, R. R., Santos, J. S., Pereira-Filho, R. N., Cardoso, J. C.,
555 Albuquerque-Junior, R. L. C., & Thomazzi, S. M. (2012). Anti-inflammatory and cicatrizing
556 activities of thymol, a monoterpene of the essential oil from *Lippia gracilis*, in rodents. *Journal*
557 *of Ethnopharmacology*, 143(2), 656-663.
- 558 Robyt, J. F., Whelan, W. J., in "Starch and Its Derivatives", Radley, J. A., Ed., 4th ed,
559 Chapman & Hall, LTD, London, 1968, pp 430-433
- 560 Romero, J., Feijoó, C. G., & Navarrete, P. (2012). Antibiotics in aquaculture—use, abuse
561 and alternatives. *Health and environment in aquaculture*, 159.
- 562 Ross, N. W., Firth, K. J., Wang, A., Burka, J. F., & Johnson, S. C. (2000). Changes in
563 hydrolytic enzyme activities of naive Atlantic salmon *Salmo salar* skin mucus due to infection
564 with the salmon louse *Lepeophtheirus salmonis* and cortisol implantation. *Diseases of aquatic*
565 *organisms*, 41(1), 43-51.
- 566 Rothmann, C., Levinshal, T., Timan, B., Avtalion, R. R., & Malik, Z. (2000). Spectral
567 imaging of red blood cells in experimental anemia of *Cyprinus carpio*. *Comparative*
568 *Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 125(1), 75-83.
- 569 Saadat Shad, H., Mazhari, M., Esmailipour, O., & Khosravinia, H. (2016). Effects of
570 thymol and carvacrol on productive performance, antioxidant enzyme activity and certain
571 blood metabolites in heat stressed broilers. *Iranian Journal of Applied Animal Science*, 6(1),
572 195-202.
- 573 Sakai, M., Hikima, J. I., & Kono, T. (2021). Fish cytokines: Current research and
574 applications. *Fisheries Science*, 87(1), 1-9.
- 575 Salinas, I., Zhang, Y. A., & Sunyer, J. O. (2011). Mucosal immunoglobulins and B cells
576 of teleost fish. *Developmental & Comparative Immunology*, 35(12), 1346-1365.
- 577 Siwicki, AX.. Anderson, D, p, 1993: Nonspecific defence mechanisms assay in fish. II.
578 Potential killing activity of neutrophils and macrophages, lysozyme activity in serum and
579 organs and total immunoglobulin (TIg) levels in serum. *Fish Diseases Diagnosis and*
580 *Prevention's Methods*, FAO-Project GCPIINT/52611PN, IF! Olsztyn: 105-112
- 581 Soltani, M., Ghodrathnama, M., TAHERI, M. A., Zargar, A., & Rouhollahi, S. H. (2013).
582 The effect of *Zataria multiflora* Boiss. and *Rosmarinus officinalis* essential oil on *Streptococcus*
583 *iniae* isolated from rainbow trout farms.

584 Syahidah, A., Saad, C. R., Daud, H. M., & Abdelhadi, Y. M. (2015). Status and
585 potential of herbal applications in aquaculture: A review.

586 Tan, P., Dong, X., Xu, H., Mai, K., & Ai, Q. (2017). Dietary vegetable oil suppressed
587 non-specific immunity and liver antioxidant capacity but induced inflammatory response in
588 Japanese sea bass (*Lateolabrax japonicus*). *Fish & Shellfish Immunology*, *63*, 139-146.

589 Tocher, D. R., Mourente, G., Van der Eecken, A., Evjemo, J. O., Diaz, E., Bell, J. G.,
590 ... & Olsen, Y. (2002). Effects of dietary vitamin E on antioxidant defence mechanisms of
591 juvenile turbot (*Scophthalmus maximus* L.), halibut (*Hippoglossus hippoglossus* L.) and sea
592 bream (*Sparus aurata* L.). *Aquaculture Nutrition*, *8*(3), 195-207.

593 Toghiani, M., Tohidi, M., Gheisari, A. A., & Tabeidian, S. A. (2010). Performance,
594 immunity, serum biochemical and hematological parameters in broiler chicks fed dietary thyme
595 as alternative for an antibiotic growth promoter. *African journal of biotechnology*, *9*(40), 6819-
596 6825.

597 Tukmechi, A., & Bandboni, M. (2014). Effects of *Saccharomyces cerevisiae*
598 supplementation on immune response, hematological parameters, body composition and
599 disease resistance in rainbow trout, *Oncorhynchus mykiss* (Walbaum, 1792). *Journal of*
600 *Applied Ichthyology*, *30*(1), 55-61.

601 Utley, H. G., F. Bernheim & P. Hochstein: Effect of sulphhydryl reagents on
602 peroxidation in microsomes. 1967. *Arch. Biochem. Biophys.* *118*, 29-32.

603 Wang, T., & Secombes, C. J. (2013). The cytokine networks of adaptive immunity in
604 fish. *Fish & shellfish immunology*, *35*(6), 1703-1718.

605 Vijayaram, S., Sun, Y. Z., Zuurro, A., Ghafarifarsani, H., Van Doan, H., & Hoseinifar,
606 S. H. (2022). Bioactive immunostimulants as health-promoting feed additives in aquaculture:
607 A review. *Fish & Shellfish Immunology*, *130*, 294-308.

608 Yousefi, M., Ghafarifarsani, H., Hoseini, S. M., Hoseinifar, S. H., Abtahi, B., Vatnikov,
609 Y. A., ... & Van Doan, H. (2022). Effects of dietary thyme essential oil and prebiotic
610 administration on rainbow trout (*Oncorhynchus mykiss*) welfare and performance. *Fish &*
611 *Shellfish Immunology*, *120*, 737-744.

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632 **Table 1.** Dietary formulation (g/kg) and proximate composition of the basal diet (Farsani et
 633 al., 2019)
 634

Ingredients	(g/kg)	Ingredients	(g/kg)
Fish meal	350	L-Carnithine	0.3
Corn gluten	175	salt	7.5
Wheat flour	180	Vitamin C	0.2
Soybean meal	130	Proximate composition of ingredients	%
Rice bran	36	Dry matter	91
Fish oil	23	Crude protein	40
Sunflower oil	48	Crude lipid	15
¹ Vitamin	15	Ash	9
² Mineral premix	15	Phosphorous	1.2
Molasses	20	Fiber	3

635
 636 ¹Vitamin contains amounts per kg of feed: retinol acetate (A): 10,000 IU; Cholecalciferol (D3): 6000
 637 IU; DL-a-tocopheryl acetate (E): 600 mg; menadione sodium bisulfite (K3):15 mg; L-ascorbic acid (C):
 638 5400 mg; Dbiotin (H2): 2.4 mg; thiamin mononitrate (B1): 45 mg; riboflavin (B2):75 mg; calcium D-
 639 pantothenate (B3): 7200 mg; niacin amide (B5): 135 mg; pyridoxine hydrochloride (B6): 45 mg; folic
 640 acid (B9): 24 mg; cyanocobalamin (B12): 120 mg;

641 ²Mineral premix (mg/kg): Fe: 60 mg; Cu: 9 mg; Co: 0.7 mg; Se: 0.75 mg; Zn: 90 mg; Mn: 39 mg; I: 3
 642 mg; cholinechloride: 150,000 mg

643

644

645 **Table 2.** The sequences of the primers used for the real-time PCR in this study.

Genes	Primers	Primer sequences	ID number
lysozyme	Forward	TGCCTGTCAAAATGGGAGTC	NM_001124716.1
	Reverse	CAGCGGATACCACAGACGTT	
Complement (C3)	Forward	GAGATGGCCTCCAAGAAGATAGAA	L24433.1
	Reverse	ACCGCATGTACGCATCATCA	
TNF- α	Forward	GGGACAAACTGTGGACTGA	AJ277604
	Reverse	GAAGTTCTTGCCCTGCTCTG	AJ401377
IL-8	Forward	CACAGACAGAGAAGGAAGGAAAG	NM_001124362.1
	Reverse	TGCTCATCTTGGGTTACAGA	
β -actine	Forward	GGACTTTGAGCAGGAGATGG	U04616.1
	Reverse	ATGATGGAGTTGTAGGTGGTCT	
Ig	Forward	AAAGCCTACAAGAGGGAGACCGAT	NM_001124235.1
	Reverse	AGAGTTATGAGGAAGAGTATGATGAAGGTG	

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662 **Table 3.** The growth and survival (Mean \pm SD) of rainbow trout, *Oncorhynchus mykiss* over
 663 60 days feeding with dietary levels of thymol (TYM). For each parameter, the differences
 664 between the means are indicated as different superscripted letters (P<0.05).

Parameters	Thymol concentrations (g/kg diet)					P values
	Non-supplemented	1	1.5	2	2.5	
Initial weight (g)	35.8 \pm 4.3	36.4 \pm 3.2	37.3 \pm 5.2	34.7 \pm 4.6	35.5 \pm 6.5	0.31
Final weight (g)	52.1 \pm 5.3 ^a	55.9 \pm 3.7 ^a	65.2 \pm 4.4 ^b	69.2 \pm 4.3 ^b	73.1 \pm 6.1 ^b	0.01
Weight gain (%)	45.5 \pm 9.1 ^a	53.57 \pm 10.3 ^a	74.7 \pm 9.5 ^b	99.4 \pm 10.8 ^c	105.9 \pm 11.5 ^c	0.006

Specific growth rate (%/d)	0.63±0.11 ^a	0.71±0.08 ^a	0.93±0.11 ^b	1.15±0.13 ^{bc}	1.21±0.11 ^c	0.014
Feed conversion ratio	2.1±0.25 ^a	2.2±0.3 ^a	1.9±0.15 ^a	1.4±0.15 ^c	1.45±0.2 ^c	0.022
Survival (%)	93.4±2.3	94.4±3.2	96.2±1.5	95.2±2.1	95.2±2.2	0.032
Feed intake (kg)	53.2±1.1 ^a	55.1±2.3 ^{ab}	58.1±2.6 ^b	63.5±2.4 ^c	65.1±1.7 ^c	0.03

665
666
667
668
669
670
671
672
673
674

675 **Table 4.** The body composition (%) of rainbow trout, *Oncorhynchus mykiss* (n= 15) over 60
676 days feeding with dietary levels of thymol (TYM). For each parameter, the differences between
677 the means (Mean ± SD) are indicated as different superscripted letters (P<0.05).

Parameters	Non-supplemented	Thymol concentrations (g/kg diet)				P values
		1	1.5	2	2.5	
Protein	12.2±0.6 ^a	13.1±0.5 ^a	14.5±0.3 ^b	14.8±0.4 ^b	15.3±0.3 ^b	0.04
Lipid	10.9±0.7	11.4±0.5	12.1±0.6	12.4±0.3	12.3±0.2	0.25
Moisture	59.2±2.3	60.1±2.4	61.4±2.5	60.3±1.8	60.6±1.3	0.11
Ash	6.2±0.3	6.5±0.5	5.8±0.6	5.7±0.4	5.8±0.5	0.33

678
679
680
681
682
683
684
685
686
687
688

689
690
691
692
693
694

695 **Table 5.** The activity of digestive enzymes (IU/mg protein) in rainbow trout, *Oncorhynchus*
696 *mykiss* (n= 15) over 60 days feeding with dietary levels of thymol (TYM). For each parameter,
697 the differences between the means (Mean \pm SD) are indicated as different superscripted letters
698 (P<0.05).

Parameters	Thymol concentrations (g/kg diet)					P values
	Non-supplemented	1	1.5	2	2.5	
Amylase	0.56 \pm 0.15	0.59 \pm 0.13	0.68 \pm 0.07	0.65 \pm 0.09	0.72 \pm 0.08	0.15
Protease	2.34 \pm 0.05 ^c	2.28 \pm 0.09 ^{bc}	3.72 \pm 0.07 ^b	3.9 \pm 0.11 ^b	4.4 \pm 0.15 ^c	0.02
Lipase	0.77 \pm 0.03 ^b	0.95 \pm 0.03 ^b	1.40 \pm 0.25 ^b	1.63 \pm 0.19 ^{bc}	1.98 \pm 0.18 ^c	0.01

699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715

716 **Table 6.** The immune components of blood and mucus (Mean \pm SD) in rainbow trout,
 717 *Oncorhynchus mykiss* (n= 15) over 60 days feeding with dietary levels of thymol (TYM). For
 718 each parameter, the differences between the means are indicated as different superscripted
 719 letters (P<0.05).

		Thymol concentrations (g/kg diet)				P values
	Non-supplemented	1	1.5	2	2.5	
Plasma immune components						
Lysozyme activity (IU/ml)	66.4 \pm 10.3 ^a	70.6 \pm 11.1 ^{ab}	95.8 \pm 12.5 ^b	121.7 \pm 9.2 ^c	130.3 \pm 10.2 ^c	0.011
C3 activity (IU/ml)	19.11 \pm 6.3 ^a	21.23 \pm 4.5 ^a	28.1 \pm 3.3 ^{ab}	32.6 \pm 5.4 ^b	38.6 \pm 6.3 ^b	0.021
Ig (mg/ml)	3.35 \pm 0.8 ^a	4.71 \pm 0.6 ^a	6.8 \pm 1.1 ^b	8.9 \pm 1.11 ^b	9.6 \pm 2.12 ^b	0.03
Bactericidal activity (NO. of CFUs)	122.5 \pm 10.3 ^a	115.5 \pm 9.1 ^a	135.6 \pm 12.1 ^b	152.6 \pm 10.2 ^b	149.3 \pm 8.3 ^b	0.01
Total protein (g/dl)	3.4 \pm 0.5 ^a	4.1 \pm 0.52 ^b	5.1 \pm 0.21 ^b	7.2 \pm 0.18 ^c	6.5 \pm 0.31 ^c	0.013
		Thymol concentrations (g/kg diet)				P values
	Non-supplemented	1	1.5	2	2.5	
Mucosal immune components						
Lysozyme activity (IU/ml)	25.3 \pm 5.2 ^a	30.5 \pm 4.1 ^a	45.7 \pm 5.2 ^b	50.6 \pm 6.1 ^b	65.4 \pm 4.2 ^c	0.02
Ig (mg/dl)	1.4 \pm 1.1 ^a	1.3 \pm 1.2 ^a	4.5 \pm 1.3 ^b	5.3 \pm 1.2 ^b	8.2 \pm 1.2 ^c	0.014
Protease activity (IU/ml)	6.3 \pm 1.4 ^a	8.1 \pm 2.1 ^{ab}	11.3 \pm 2.2 ^{bc}	14.4 \pm 1.4 ^{bc}	14.6 \pm 1.5 ^c	0.01
Alkaline phosphatase activity (IU/ml)	1.25 \pm 0.4 ^a	1.34 \pm 0.5 ^a	2.55 \pm 0.32 ^b	3.3 \pm 0.17 ^c	4.58 \pm 0.21 ^d	0.022

720 * C3: complement C3, Ig: total immunoglobulin.

721

722

723

724

725 **Table 7.** The hematological alternations (Mean \pm SD) in rainbow trout, *Oncorhynchus mykiss*
 726 (n= 15) over 60 days feeding with dietary levels of thymol (TYM). For each parameter, the
 727 differences between the means are indicated as different superscripted letters (P<0.05).

		Thymol concentrations (g/kg diet)				P values
Parameters	Non-supplemented	1	1.5	2	2.5	
RBC ($\times 10^6/\mu\text{l}$)	1.5 \pm 0.12 ^a	1.7 \pm 0.21 ^a	1.45 \pm 0.12 ^a	2.5 \pm 0.1 ^b	2.8 \pm 0.13 ^c	0.01
WBC	5.5 \pm 1.2 ^a	5.8 \pm 1.4 ^a	7.2 \pm 1.8 ^{ab}	10.6 \pm 2.3 ^b	11.5 \pm 2.5 ^c	0.015

$(\times 10^3/\mu\text{l})$						
Hct (%)	24.5±2.4 ^a	25.4±1.8 ^a	27.9±2.6 ^a	33.2±2.4 ^b	35.1±2.7 ^b	0.008
Hb (g/dl)	1.99±0.15 ^a	2.02±0.51 ^a	2.1±0.13 ^a	3.41±0.11 ^b	3.6±0.21 ^b	0.011
MCV (fl)	163.3±10.3 ^a	149.4±11.3 ^{ac}	192.4±10.1 ^b	132.8±14.1 ^c	125.3±13.4 ^c	0.02
MCH (pg)	13.2±2.2	11.88±1.7	14.4±2.2	13.64±2.1	12.85±2.2	0.035
MCHC (%)	0.81±0.12 ^a	0.79±0.13 ^a	0.75±0.1 ^a	1.02±0.12 ^b	1.03±0.13 ^b	0.002

728 RBC: red blood cell, WBC: white blood cell, Hct: hematocrit, Hb: hemoglobin, MCV: mean corpuscular volume,
729 MCH: mean corpuscular hemoglobin, MCHC: mean corpuscular hemoglobin concentration.

730

731

732

733

734

735

736

737 **Table 8.** The liver antioxidant changes (IU/mg) in rainbow trout, *Oncorhynchus mykiss* (n=
738 15) over 60 days feeding with dietary levels of thymol (TYM). For each parameter, the
739 differences between the means (Mean ± SD) are indicated as different superscripted letters
740 (P<0.05).

Parameters	Non-supplemented	Thymol concentrations (g/kg diet)				P values
		1	1.5	2	2.5	
SOD	4.4±1.1 ^a	5.1±1.2 ^a	8.1±1.7 ^b	9.3±2.13 ^b	11.5±2.14 ^b	0.02
CAT	5.2±1.3 ^a	7.9±1.55 ^b	8.1±1.7 ^b	11.2±1.3 ^c	12.3±1.2 ^c	0.01
GPx	3.1±0.08 ^a	4.11±0.1 ^a	7.8±0.13 ^b	8.5±0.1 ^b	9.7±0.13 ^b	0.04

741 * SOD: superoxide dismutase, CAT: catalase, GPx: glutathione peroxidase.

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756


757

758

759


760

761



762

763




764

765


766

767




768
769
770
771
772
773
774

Fig. 1. Relationships between the dietary TYM levels and growth parameters of the rainbow trout (n = 3).




775
776
777
778
779
780
781
782
783
784
785
786

Fig. 2. The changes of malondialdehyde (MDA) levels in the rainbow trout, *Oncorhynchus mykiss* over 60 days feeding with dietary levels of thymol (TYM). The differences between the means are indicated as different superscripted letters (P<0.05).




787
788
789
790
791
792
793
794

Fig. 3. The expression of immune-related genes, complement (C3), Ig and lysozyme in the rainbow trout, *Oncorhynchus mykiss* over 60 days feeding with dietary levels of thymol (TYM). The differences between the means are indicated as different superscripted letters (P<0.05).



795
796
797
798
799
800

Fig. 4. The expression of inflammatory-related genes, TNF-α and IL-8 in the rainbow trout, *Oncorhynchus mykiss* over 60 days feeding with dietary levels of thymol (TYM). The differences between the means are indicated as different superscripted letters (P<0.05).



801
 802
 803
 804
 805
 806
 807
 808

Fig. 5. Cumulative mortality of rainbow trout, *Oncorhynchus mykiss* (n = 10/tank) fed different concentration of TYM in the diet throughout 10-day challenge with *Streptococcus iniae*. Significant differences are shown with different lower-case letters (P < 0.05). Data: Mean ± SD.

Dietary thymol improved growth, body composition, digestive enzyme activities, 2 hematology, immunity, antioxidant defense, and resistance to Streptococcus iniae in the 3 rainbow trout (Oncorhynchus)

ORIGINALITY REPORT

9%

SIMILARITY INDEX

2%

INTERNET SOURCES

8%

PUBLICATIONS

%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

< 1%

★ Ramasamy Harikrishnan, Chellam Balasundaram.
" In Vitro and In Vivo Studies of the Use of Some Medicinal Herbals against the Pathogen in Goldfish ", Journal of Aquatic Animal Health, 2008

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On