D: 2087684534			Similarity by Source
Vord Count: 6496 Submitted: 1		Similarity Index	Internet Sources: 14
14217-38137-1-PB.pdf By Anonymous		17%	Publications: 11 Student Papers: 19
6% match (Internet from 22-Jul-2021) <u>https://repository.its.ac.id/80915/</u>			
2% match (Internet from 12-Feb-2023) https://www.researchgate.net/publication/341972256 Bivariate zero inf	flated genera	alized Poisson regre	ssion model in the
1% match (Internet from 24-Nov-2020) http://repository.its.ac.id/45708/1/1314201011-Master Thesis.pdf			
1% match (Nendy Septi Arniva, Purhadi, Sutikno. "Parameter Estimation Geographically Weighted Bivariate Poisson Inverse Gaussian Regression", Intelligent Informatics (SAIN), 2018) <u>Nendy Septi Arniva, Purhadi, Sutikno. "Parameter Estimation and Statistic</u> <u>Weighted Bivariate Poisson Inverse Gaussian Regression", 2018 Internation Informatics (SAIN), 2018</u>	2018 Intern	ational Symposium c <u>xed Model of Geogra</u>	n Advanced <u>phically</u>
1% match () <u>Khaeruddin, Khaeruddin, Nur, Mohammad, Wasis, Wasis. "Critical Thinking Learning Science-Physics", 2016</u>	<u>g Skills Profil</u>	<u>e of High School Stu</u>	<u>dents In</u>
1% match (Jaka Nugraha, Welly Nur Armawati. "Modelling on human imm regression", AIP Publishing, 2018) <u>Jaka Nugraha, Welly Nur Armawati. "Modelling on human immunodeficien AIP Publishing, 2018</u>			
1% match (Internet from 03-Sep-2022) https://www.semanticscholar.org/paper/Multivariate-generalized-Poisson- Purhadi/bc735dbd5e02a9ba5eea73f6e5f0cc4fd5028095	-regression-n	nodel-a-Berliana-	
1% match (Sischa Wahyuning Tyas, Gunardi, Lila Aryani Puspitasari. "Geo regression model with the best kernel function in the case of the number MethodsX, 2023) <u>Sischa Wahyuning Tyas, Gunardi, Lila Aryani Puspitasari. "Geographically</u> with the best kernel function in the case of the number of postpartum ma	of postpartu	m maternal mortality	ression model
1% match (I U Utami, I Setiawan, D Daniaty. "Modelling the number of H Conference Series, 2021) <u>I U Utami, I Setiawan, D Daniaty. "Modelling the number of HIV/AIDS in (</u> <u>Series, 2021</u>		,	,
1% match (Internet from 26-Apr-2023) https://en.tatoli.tl/2021/07/05/1-374-people-have-been-infected-with-hi	v-aids/06/		
1% match (Internet from 16-Mar-2016) <u>http://edepot.wur.nl/44380</u>			
1% match (student papers from 26-Jun-2021) Submitted to Bellevue Public School on 2021-06-26			
1% match (student papers from 08-Feb-2023) <u>Submitted to Universitas Indonesia on 2023-02-08</u>			
ol.11 (2021) No. 6 ISSN: 2088-5334 <u>Bivariate Zero-Inflated Poisson Inverse</u> <u>pplication Purhadi</u> a, <u>Ermawati</u> a,b,*, <u>Rossy Noviyana</u> a, <u>Sutikno</u> b a Depar epuluh Nopember, Surabaya, 60111, Indonesia b Department of Mathemati lakassar, 92118, Indonesia Corresponding author: *erma.mat@gmail.com A ero-Inflated Poisson Inverse Gaussian Regression (BZIPIGR) model to prese stimation and modeling of the number of HIV and AIDS cases in each sub-g egencies to determine the factors that have a significant effect. This model verdispersion cases caused by extra zeros in the response variables. The per ses the Maximum Likelihood Estimation (MLE). The first derivative of the B2 nerefor it has continued with the Berndt Hall Hall Hausman (BHHH) iteration stimators, while the hypothesis testing of the BZIPIGR model is derived usi pproach. <u>Based on the AICc</u> value <u>obtained</u> , the BZIPIGR model is a feasible f HIV and AIDS cases in Trenggalek and Ponorogo Districts, East Java Provin n the increase in the number of HIV and AIDS cases was the percentage of the <u>variables that</u> had a significant <u>effect</u> on <u>reducing the number of HIV and</u> population aged 25-29 years, the percentage of reproductive-age couples us	rtment of Sta cs, Universit: Abstract— <u>Th</u> ents the form district in Tre can be used (arameter esti ZIPIGR mode <u>1 to</u> obtain <u>th</u> ng Maximum e model to b nce. The vari <u>the populati</u> d AIDS case	tistics, Institut Tekno as Islam Negeri Alau is study developed a of BZIPIGR parame inggalek and Ponorog on data that have mation of the BZIPIG as obtained not cl maximum likelihoo be i Likelihood Ratio Tes e applied to data on able that had a signi on with low educatio s were the percentage	ologi ddin <u>Bivariate</u> ter <u>30</u> GR <u>model</u> osed form, od t (<u>MLR</u>) the <u>number</u> ficant effect <u>n (SMA</u>). <u>39</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u> <u>30</u>

https://www.turnitin.com/newreport_printview.asp?eq=0&eb=1&esm=-1&oid=2087684534&sid=0&n=0&m=2&svr=29&r=95.10892275514458&lan... 1/5

Ordinary Least Square (OLS) regression because it will violate the assumption that the error value follows a normal

Turnitin - Originality Report - 1._14217-38137-1-PB.pdf

distribution and has heteroscedasticity. One of the models that can apply for modeling count data is Poisson regression, but this model must fulfill the Equi dispersion assumption (mean and variance are equal). This assumption is often violated. That is, the variance value is smaller than the mean (under dispersion) or otherwise (overdispersion)[1]. The number of zeros (excess zeros) can cause a large enough variance that is known as overdispersion[2]. Occasionally, an event rarely occurs. Therefore, many responses are zero (excess zero). If the case is disregarded, it will underestimate and wrong decision when testing the hypothesis[3]. Several models have been developed to solve the overdispersion problem. The modeling can use Generalized Poisson[4] or a mixed Poisson distribution both discrete and continuous. Some of the mixed-Poisson distributions that have been developed are Negative Binomial [5], and Poisson Inverse Gaussian (PIG) [6]. Poisson Inverse Gaussian (PIG), a mixed Poisson distribution with random effects, has an Inverse Gaussian distribution. These distributions are not good enough to handle overdispersion and extra zeros cases. Several other mixed Poisson distributions can manage this problem, they are Zero Inflated Negative Binomial (ZINB)[5], Zero Inflated Generalized Poisson (ZIGP)[7], and Zero Inflated Poisson Inverse Gaussian (ZIPIG)[8]. Based on the results of research by Hilbe that the ZIPIGR model is the best compared to ZIP or ZINB in modeling count data with many significant zero values[9]. Constraints in <u>the parameter estimation</u> process <u>of the ZIPIGR model</u> using <u>Maximum</u> <u>Likelihood Estimation (MLE</u>) is <u>the</u> function of the model is complicated, thereby to overcome this problem in obtaining parameter estimator must be followed by numerical iteration. One of the numerical iterations that can apply is the Berndt Hall Hall Hausman (BHHH) algorithm[10], which in this algorithm only uses the first derivative of its likelihood function. The ZIPIGR model is still limited to one response variable only. Meanwhile, model cases with two response variables will be developed to the Bivariate Zero Inflated Poisson Inverse Gaussian Regression (BZIPIGR) model. This study will be applying it to modeling the number of HIV and AIDS cases. In this case, there are many zeroes in the response variable, so the BZIPIGR model is suitable for the application. HIV (Human Immunodeficiency Virus) is a virus that attacks white blood cells, which causes a decrease in the body's immune system. AIDS (Acquired Immune Deficiency Syndrome) is a group of diseases that arise due to decreased immunity caused by HIV infection. HIV and AIDS have become an epidemic worldwide because no medicine can cure the sufferer. Also, the symptoms of this disease are not visible, and the course of this disease takes a long time. The fifth highest rankings of HIV infections that occurred in Indonesia from 1987 to 2019 were East Java, DKI Jakarta, Papua, Bali, Riau, and West Java [11]. In opposition to HIV, the three highest rankings of AIDS infections were central Java, Papua, and East Java. In 2019 the number of HIV 8,885 cases and 920 cases of AIDS in East Java, this number increased from the previous year[12]. Many HIV and AIDS infections in East Java, which is extremely high, need to be addressed. Trenggalek and Ponorogo regencies are districts in East Java Province. This study will apply the BZIPIGR model to data on the number of HIV and AIDS cases in Trenggalek and Ponorogo Districts. II. MATERIALS AND METHOD A. Poisson Inverse Gaussian Regression (PIGR) The PIG distribution is a mixed Poisson distribution consisting of two parameters, namely µ (mean) as the location parameter and τ (dispersion parameter) as the shape parameter. Let Y is the response variable that PIG distribution and can be denoted 11, y?0 (1) with s ? y ? 1 and z ? 2? . Where according to 1 2 ? 2? ? Wilmot, Ks ?z? ? Ky?12 ????1 ?2?? ?1? ?? is the third kind of the ? modified Bessel function [14]. The expected value and variance of the PIG distribution are E?Y ? ? E? E?Y | ?v?? ? E??v? ? ?. Var?Y? ?Var?E?Y | ?v??? E?Var?Y | ?v??? ? ???2. Y is the number of events in a unit of observation in a certain period, ? is the average of these events, and tau is an overdispersion parameter and is the same as Var (v), which is due to the heterogeneity or diversity associated with the unit of observation with certain characters [12]. Suppose Yi is the response for the ith observation and xiT ? ??1 x1i x2i ... x pi ?? is the explanatory vector for the T ith observation with the dimension (k+1) x1, so the PIG Regression Model is as follows: ?i ?exp?xiT β ? or log??i??xiT β , where β????0 ?1 ?2 ... ?k??T . B. Bivariate Poisson Inverse Gaussian Regression (BPIGR) The Bivariate Poisson Inverse Gaussian (BPIG) distribution has two count variables that are correlated. Suppose that two random variables have a Poisson distribution and are independent of each other, Y1 and Y2, which have the mean vµ1 and vµ2 respectively, and the variance Var(Y1) ? ?1 ?? 2? and Var(Y2) ? ?2 ?? 2?? . Variable v is a random variable with Gaussian inverse distribution. Hence Y1 and Y2 have mixed Poisson distribution, namely Poisson Inverse Gaussian. The BPIG distribution 2 where s ? y1 ? y2 ? 1 2 and z ? ?1 ???1 ? 2? ?h?1 ?h ??2 so ? ? 1 Ks ?z?u,v?? ? K y1?y2?12 ???? ?? ?1? 2 ? 1 ? 2? ?h?1 ?h ??? 2 ???? BPIGR is a regression model with two correlated responses. Suppose yih is the hth response to the ith observation and is give a random sample (Y1i ,Y2i) ~ BPIG ??hi ,? ? where i ? 1, 2,..., n and h ? 1, 2 . Then the BPIGR model can be written as follows. log?E(Yih)?? xTiβh and E(Yih)?exp?xTiβh?, where. xi ? ??1 x1i x2i ... x pi ??T is the explanatory vector for the ith observation, i = 1, 2, ..., n β h ? ???h0 ? ? h2 ... ?hp ?? is the vector of the T h1 regression and Y2 are random variables with a bivariate Poisson distribution ?Y1,Y2 ? ~ BP ??1, ?2 , ?0 ? where the combined probability function is[16] ? p ? ?1? p?exp????1 ? ?2 ? ?0 ?? ? y1 ? y2 ? 0 ?? ? y1 ? j y2 ? j ? j P ?Y1,Y2 ? ? ???1 ? n? min??y1,y2 ? 1 ? 2 0 ?y1? j?!?y2? j?!j! (3) ? ? j?0 exp????1 ? ?2 ? ?0 ?? y1 ? 0 y2 ? 0 The mean, the variance and the covariance values of the BZIP distribution are. E ?YI ? ? ?1 ? p ? ? ?I ? p ? ?! ? ?0 ? Var ?YI ? ? E ?YI ? ??1 ? p ? ?! ? ?0 ??? , I ? 1, 2 Cov?Y1,Y2?? ?1? p????0 ? p??1 ? ?0???2 ? ?0???? The BZIP regression model is as follows. ?hk ??0 ?exp?xTk?h?;h? 1,2 (4) Method of estimation in the BZIPR model is a MLE. The method for calculating the test statistics on the parameter test is the Maximum Likelihood Ratio Test (MLRT)[17]. D. Zero Inflated Poisson Inverse Gaussian Regression (ZIPIGR) If Y=0 with the probability value v and Y=Y1 with the probability value (1-v), then Y has a Zero-Inflated Poisson Inverse Gaussian distribution, which can be written as withthe probability function follows[18]. P.?Y.?.y.l.?., p <u>? ? ????? p</u> ??1??1 ?p?pP??PY?<u>Y? ?y</u> |0?| ,??,?? ? ; ; <u>y</u> ? 0 <u>y</u> ?1,2,3,... (5) The ZIPIGR model can be written as follows. p?1?exepx?px?Tx?T??? and ?? exp?xT??E. Bivariate Zero Inflated Poisson Inverse Gaussian Regression (BZIPIGR) Let Y1 and Y2 are random variable which BZIPIG iid distribution with ?Y1i ,Y2i ? ~ BZIPIG ? ?0 , ?1, ? 2.,?1,? 2., p then the joint probability function of Y1,Y2 as follows [8] a. If y1 ? 0 and y2 ? 0 , then P ?Y1 ? 0,Y2 ? 0? . P ? y1i , y2i ? ? p2 ? xi ? p1 ? xi ? ? ???1 ? p2 ? xi ? ? p1 ? xi ? ??? ??? ?1 2?2i? ?1??2 exp???1? ? ? K?12 ?? ?1 2?2i? ?1??? ? ?? p2 ? xi ? ?1 ? p1 ? xi ? ? ? ? ??? ?2? ??? 12 exp ?? ?1 ? ? 2?1i? ? 1? 4 K 1 ?? ?1 2?1i? ? 1 1 ? 1 ? 2 ???? ? ????1 ? p1 ? xi ? ? ? 1 ? p2 ? xi ? ? ? 2 ?2 ?? ?? ?? exp ?? ?1 ? ?2 ?? ?1 ? ?2 ?? 1? 4 K? 1 ? ?1 2? ??1i ??2i ? ? 1??? 1 ? 2 ?(6) b. If y1 ? 0 and y2 ? 0 , then P ?Y1 ? 0, Y2 ? 0 ? . P ? y1i , y2i ? ? ? 1 ? p1 ? xi ?? ?1 ? p2 ? xi ?? e?1 ??? ?2? 1 2? ??1i ? 2i ? ? 1 ??2 ?1i y1i ?2i y2i ? exp?? ?1? Ky1i?y2i?1 ?? ?1 2? ??1i ? ?2i ? ?1? 2 ? 1? 2? ??1i ? ?2i ??y1i?y2i y1i !y2i ! (7) The BZIPIGR model can be written as follows. p1 ? xi ? ? 1 ?exepx?p????1?x1Tix β Ti1 β ?1 ? 1 ? p1 ? xi ? ? 1 ? exp ? ??1xTi β 1 ? 1 p2 ? xi ? ? 1?exepx?p????2xZTixBTi2B?2 ? 1? p2 ?xi ? ? 1? exp???2xTiB2 ? 1 logit ? p1i ?=log ??? 1 ?p1p?1x?ix?i ? ??? = ? ? 1xTi β1 ?1i ? exp ? xTi β1 ? logit ? p2i ? ? log ???1?p2p?2x?ix?i ? ???= ??2xTiβ2 ?2i ? exp?xTiβ2 ? (8) F. HIV and AIDS and Factors Affecting the Number of HIV and AIDS Cases HIV is a type of virus that attacks/infects white blood cells, which will cause a decrease in human immunity and impair its function. If HIV infection is not handled immediately, it will develop into serious health called AIDS, which is the final stage of infection caused by HIV, which can infect the body's organ systems, including the brain, thus damaging the immune system. Until now, there is no cure for HIV and However, there are drugs to slow the progression of the disease and increase the life expectancy of sufferers. HIV and AIDS cases were determined by three main factors, namely[19]. 1. Predisposing factors. The factors that encourage a person's behavior include knowledge, attitudes, religion, beliefs, values, traditions, etc. The driving factors for HIV exposure are low knowledge of HIV, accepting attitudes about the behaviors that cause HIV, and traditions that can lead to HIV. 2. Enabling factors. Factors that facilitate a person's behavior to become infected with HIV such as facilities and infrastructure for the occurrence of health behaviors, enabling factors that can cause HIV, namely the distance between prostitution and prostitution localization close to home, distance to health services that are far away so that there is less information about HIV. 3. Reinforcing factors are factors that amplifier the behavior of a person infected with HIV. Sometimes even though people know and can behave healthily, they do not. Examples include family, health workers, and community leaders. In cases of HIV exposure, reinforcing factors are family rules that do not prohibit HIV-causing behavior and the enabling environment for HIV-causing behavior. G. Data Source This study used secondary data obtained from the Health Service of Trenggalek and Ponorogo Districts Profile in 2012. This data has been used by Wijaya [20] to know the factors that can affect the number of HIV and AIDS cases. The observation units used in this study were 27 districts in Trenggalek and Ponorogo regencies. Research variables used in this study consisted of two

https://www.turnitin.com/newreport_printview.asp?eq=0&eb=1&esm=-1&oid=2087684534&sid=0&n=2&svr=29&r=95.10892275514458&lan... 2/5

Turnitin - Originality Report - 1. 14217-38137-1-PB.pdf

responses ?Y ? and five predictor variables (X). The variables in this study are given in Table 1: TABLE I RESEARCH VARIABLES Response Variables Y1 The number of HIV Y2 The number of AIDS Predictor Variables X1 The percentage of age group 25-29 years X2 The percentage of population with senior high school level X3 The percentage of couples of reproductive ages using condoms X4 The Percentage of health education activities about HIV and AIDS X5 The percentage of community health insurance (Jamkesmas) H. Steps to determine the factors that influence the number of HIV and AIDS cases using BZIPIGR Model The steps to determine the factors that influence the number of HIV and AIDS cases are as follows. 1. Make a descriptive analysis of variables. 2. Correlation test between responses using the Pearson test. 3. Detect multicollinearity from predictor variables using the VIF test criteria. 4. Testing of overdispersion using the Variance Test (VT) 5. Perform data analysis using the BZIPIGR model. a. Specifies the parameter estimator value using MLE with BHHH algorithm. b. Conducting hypothesis testing simultaneously using MLRT on the regression parameters. c. Perform partial hypothesis testing on regression parameters. d. Interpret the analysis results. e. Conclusion. III. RESULT AND DISCUSSION A. Parameter Estimation of BZIPIGR The parameter estimation of the BZIPIGR model uses MLE method. The probability distribution for BZIPIG is in equation (8). The first stage is to estimate the parameters using MLE by forming the likelihood function, which is defined as follows. n n L ? β1 , β 2 ,? 1 ,? 2 ,? ? ??? Ai ???? Bi ? (9) i? 1 i ?1 where. Ai ?1?exp???2xTiβ2?1?exp???1xTiβ1??01?O2?O3 exp???2xTiβ2 exp???1xTiβ1 ? ? O1 ? 1 ? exp ? ?? 2 xTi β 2 ? 1 ?exepx?p????1?x1TixβTi1β?1 ? ??? ?2? 2 exp ? xTi β 2 ?? ? 1 ?? 2 1 1 ? exp?? ?1 ??2 ?? ?1 2exp?xTiβ2 ?? ?1? K 1 O2 ? 1 ?exepx?p????2?x2TixβTi2β?2 ? 1 ? exp ? ??1xTi β1 ? ??? ?2? ??? 2 ? 2? exp ? xTi β1 ? ? 1?4 1 1 1 exp ?? ?1 ? ?? ?1 2? exp ? xTi β1 ? ? 1 ? K ? 1 2 1 1 03??1?exp????1x?Ti?β?1???1?exp???2xTiβ2???2??exp?xTiβ1??exp?xTiβ2???1?4 ? 2 ?2exp???1? ? ? K?12???1 2??exp?xTiβ1??exp?xTiβ2???1? 1 1 Bi ? e? ????1???exp2?????e1xxpTi?βx1Ti?β??11??? (11) i ?1 i ?1 bi is a dummy variable equal to 1. Furthermore, to obtain parameter estimation, the $l(\beta h,?h,?;h$?1,2) function is derived against the parameter $\beta_{1,\beta_{2,2},1,2}$ and? then equated with zero. These first derivative does not yield closed form, hence parameter estimation was continued to a numerical method, namely the BHHH iteration[21], where the initial value γ ?[β T1 β T2 ?1???]T and m?0 with γ \$?0? ?0 corresponds to the ZIPIGR function for convergence tolerance limits. The initial values for $\beta_{1,\beta_{2,2},1,2}$, and? are obtained from the estimator in the ZIPIGR model. The iteration starting from m?0 in the following equation. γ\$m?1? γ\$m?H?1?γ\$m?g?γ\$m? . Iteration will stop if value $\gamma = 1$, where ϵ is a very small positive number close to 0, where the gradient vector g? γ (m)? is ?L(.)?T?? i?,γ[^]βm1,?β2,?1,?2,? ? ? B. Hypothesis Testing of Parameter BZIPIG Model Testing Hypotheses are examining that simultaneously and partially to determine the significance of each parameter. 1) SimultaneousTestingParameters β 1, β 2 Hypothesis testing of parameters BZIPIGR model, determined by the likelihood function L???? and L???? , where L???? is the Maximum Likelihood value under the population which involved the predictor variables, and otherwise L??? ? is the likelihood function under H0 . Simultaneous testing of the parameters BZIPIGR model is to determine the significance of the parameters β_1 , β_2 together, with the following hypothesis. H0 : ?1k ? ?2k ? 0with k ?1,2,K, p H1 : at least one of ? hk =0 with <u>h ?1,2</u> and <u>k ?1,2,K</u>, p Before managing the test statistic to be used, it will define the parameter set under the population , namely , then from the set of parameters, the likelihood function can be formed under the population like in equation (9). The maximum likelihood value under population L??? is L??^? were obtained from section 3.1. 20 ? (13) the maximum likelihood value under H0 L??? is L???? obtained in the following way. Substitute equation (14) in equation(6)and(7)toobtain P ? y1i , y2i ? ? ???? DCii ,,?? yy11ii ,, yy22ii ?? ?? ??00,,00?? (14) The first step is to estimate the parameters using MLE by forming the likelihood function, which is defined as follows. n n L ? 10 , ?20 ,?10 ,?20 ,? ???? ?Ci ???? Di ? (15) i ?1 i ?1 where. Ci ? 1 ? exp ? ??20 ? 1 ? exp ? ??10 ? ? 01 ? 02 ? 03 exp ? ??20 ? exp ? ??10 ? O1 ?1?exp???20?1?exepx?p????1?01?0?????2? 2exp?ß20?? ?1??2 1 1 ? exp ?? ?1 ? K ?1 ? 1 2 ? ? 2exp?ß20?? ? 1? O2 ? 1 ?exepx?p????2?O2?O ? 1 ? exp ? ??10 ? ??? ?2? ??? 2 ? 2? exp ? β10 ? ? 1? 4 1 1 1 exp ?? ?1 ? K ?12 ?? ?1 2? 12 ?? ?1 2? ? exp ? β10 ? ? exp ? β20 ? ? ? 1 ? e? ?? 2??exp?β10??exp?β20???1??2 P1 1 1 2 Di ? ????1?exp???1???1? exp?????? P1 ? exp ? β20 ? y1 exp ? β20 ? y2 K 12 ?? ?1 2? ? exp ?β10 ? ? exp ? β20 ? ? ? 1? ? ? 1? ?? exp ?β10 ? ? exp ? $\beta 20$? ? ? y1? y2 y1 ! y2 ! exp????? The second stage is to determine the log from the likelihood function. I ? log L(?10 , ?20 ,?10 ,?2, 0?; 0?; ?) ? log ? Ci ?1?bi ? Di ?bi n (16) i ?1 n n I ? ??1? bi ? log ?Ci ? ??bi log ? Di ? (17) i ?1 i ?1 bi is a dummy variable with a value of 1. Furthermore, to obtain parameter estimation, the function I(?10,?20,?10,?20,?) defines a derivative for parameters then equates to zero. This first derivative for each parameter does not yield close form. Then for getting L??^? is using BHHH algorithm. From these two maximum likelihood values, a ratio is formed between the maximum likelihood value under population L??? and the maximum likelihood value under H 0 L ??? , which is called the odds ratio, as follows: ?? L???? L??? with the rejection area of H0? < ?0 where 0 < ?0 < 1. The next is to determine the distribution of test statistics with the n big sample approach as follows. G 2 ? ? log ? ? 2 ? with G 2 ? c and ? ? P ? G 2 ? c | H 0 is true ? c will obtain if G2 is known, so this simultaneous test uses the test statistics. G2 ??log??2? ? ?2log ?? L??^? ?? ?? L??^? ?? ? 2 ?log ? L ??^?? ? log ? L ??^ ???. will compare with ??2,?a?b? , where a is the number of parameters under the population, and b is the number of parameters under H0. The rejection area of H0 if G2 ? ??2, ?a?b? . 2) Partial Parameter Testing If the test results produce a rejecting conclusion, then the test can be continued by partially testing with the Wald test statistics. This test, to find out the effect of the predictor variable 1:? h ? 0 with h ? 1, 2 iii) H 0:? ? 0 H 1:? ? 0 with the test statistics which used are: Z? se??h khk? where se?? hk? va^r??^hk? ^ ? ? ? Z? ? s?e^h???^h1k? where se ??^hk ? ? va^r ??^hk ? Z? ? se??\$? wherese??\$?? va^r??\$? ? \$ Where the value of va^r??[^]hk?, va^r??[^]hk?, and va^r??^{\$}? are obtained from the main diagonal elements of the matrix? H?1??^? . H0 reject if Zscore ? Z? . 2 C. Descriptive statistics of Response and predictor variable In each sub-district in Trenggalek and Ponorogo Districts, the highest number of HIV and AIDS cases was 4 cases and almost 50% of the subdistricts had no cases or more than 50% of the sub-districts had cases. The highest mean and variance from the variable of percentage public health insurance and the lowest in the percentage health extension activities variable. This shows that there is still low awareness of the importance of health education, which is followed by a low percentage of reproductive-age couples (PUS) using condoms. The average percentage for the population of the lowest level of education in a senior high school is 15.43%, with the highest in the Pudak sub-district. The average percentage of couples of reproductive ages using condoms is 3.283%, with the highest in the Tugu sub-district and the least in the Ngebel sub-district. The percentage of health education activities is 0.6407%, with the highest in the Pudak sub-district and the least in the Gandusari sub-district. The percentage of public health insurance is 42,75%, with the highest in the Pudak sub- district and the least in the Trenggalek sub-district. Based on the value of the correlation coefficient that there are positive relationship patterns of the number cases of HIV and AIDS variable with the percentage of the age group 25- 29 years, the percentage of the lowest level of education (SMA), and percentage of couples of reproductive ages using condoms. Meanwhile, the percentage of health education activities and the percentage of public health insurance are negative relationship pattern to the response variable. 1) Pearson Correlation Test The two responses in the bivariate regression analysis must correlate. To determine whether there is a relationship is tested using the Pearson correlation coefficient with the following hypothesis[10]. H0 : ? y1, y2 ? 0 H1 : ?y1, y2 ? 0 Test Statistics. T ? ry1, y2 n ? 2 0.851 27 ? 2 ? ? 8.102 1? r2 1? 0.8512 The T value (8.102) is greater than t0.025,25 ? 2.059 therefore H0 is rejecting or the number cases of HIV and AIDS are correlated, so this research can continue with the bivariate model. 2) Multicollinearity Test There are many ways to detect multicollinearity, one of these is the Variance Inflation Factor (VIF) value or a value that describes the increase in the variance of the estimated parameters between predictor variables. If

https://www.turnitin.com/newreport_printview.asp?eq=0&eb=1&esm=-1&oid=2087684534&sid=0&n=2&svr=29&r=95.10892275514458&lan...____3/5

Turnitin - Originality Report - 1._14217-38137-1-PB.pdf

the VIF ?10 , it can be said that there is multicollinearity, on the other hand, if the VIF ?10 , there is no multicollinearity[22]. The VIF value of predictor variables are less than 10 (X1=1.22, X2=2.32, X3=1.93, X4=1.06, and X5=1.06). Therefore, the predictor variables do not experience multicollinearity, or there is no correlation between the predictor variables. 3) Overdispersion Test This study conducted an overdispersion test to determine whether the data used violates the assumptions or not. One way that can do to detect the presence or absence of overdispersion is using the variance test (VT), where overdispersion occurs if VT>1. The value of VT can be calculated using[10]. ?? yi ? y ?2 n VT1 ? i ?1 y ? 47.4074074 1.148148 ? 41.29 n VT2 ? i?1 ??yi ? y?2 y ? 24.0740741 ? 29.545 0.814815 According to VT value for Y1 and Y2 (HIV and AIDS cases) have VT1 and VT2 are more than 1, therefore the data had overdispersion problems, hence this study continued using the BZIPIGR model. D. Regression Modeling of the Number of HIV and AIDS Cases in Trenggalek and Ponorogo Districts using the BZIPIG method. The number of HIV and AIDS cases in Trenggale and Ponorogo Districts is count data with overdispersion problems. Therefore, this research modeling using BZIPIGR. Simultaneoustestingofparametersiscarried outtodetermine whether the significant $\beta 1, \beta 2$ parameter jointly affects the response. Simultaneous testing of the parameters of the BZIPIGR model with a hypothesis H0 : ?1k ? ?2k ? 0 with k ? 1,2,K,5 H1: at least one of?<u>hk =0</u> with <u>h ?1,2</u> and <u>k ?1,2,K</u>,5 The value of test statistics G (73.18336) is greater than ? (20.05;10) ?18.30704 , hence rejecting H0 . <u>The conclusion is</u> predictor <u>variables</u> together <u>affect the response</u>. After testing simultaneously, the next step is testing the parameters BZIPIGR model partially. Thehypothesis for this testing is: i) For ? H0 : ?<u>hk ? 0 ; h ?1,2; k ?1,2</u>,3,4,5 H1 : ?hk ? 0 ii) For ? H0 :?h ? 0 ; h ?1,2 H1 :?h ? 0 iii) For ? H 0 : ? ? 0 H1 : ? ? 0 The partial hypothesis testing on the parameters of the BZIPIGR model can be seen in Table 2. TABLE II PARAMETER ESTIMATION OF BZIPIGR MODEL Parameter Estimate Se Z ????? -1.117 ????? -0.097 ????? 0.064 ????? -0.042 ????? -0.27 ????? -0.015 ????? -1.963 ????? -0.019 ????? 0.056 ????? -0.014 ????? -0.01 ????? -0.022 ??? 0.05 ??? 0.057 ?? 0.006 0.001417051 0.001422267 0.001409661 0.001405293 0.001415502 0.001417234 0.001515899 0.001539334 0.001515931 0.001507484 0.001469292 0.001510367 1.3956E-07 2.59258E-07 9.81499E-05 -788.257 -68.201 45.401 -29.887 -190.745 -10.584 -1294.941 -12.343 36.941 -9.287 -6.806 -14.566 358267.700 219858.600 61.131 P ? value of the all parameters are less than the ? ? 0.05 (P-Value < 0.05), hence H 0 is rejecting. Therefore, all the predictor variables partially influence the number of HIV and AIDS cases in Trenggalek and Ponorogo Districts. They are the percentage of the age group 25-29 years, the percentage of the lowest level of education is the senior high school variables, the percentage of couples of reproductive ages using condoms, the percentage of health education activities, and the percentage of public health insurance. The equation of the BZIPIGR model according to table 2 for the number of HIV cases is divided into 2 models, such as the model for Poisson state and the model for zero state. The Poisson state regression model for HIV as follows. log??1i???1.1169? 0.0966 X1i? 0.0643 X 2i? 0.0423 X 3i?? 0.2703 X 4i ? 0.0150 X 5i The estimation of parameters in the equation above can be interpreted based on the independent variables' coefficient and coefficient sign. For instance, every 1% increase of the population aged 25-29 years ? X1? will reduce the average number of HIV cases in each sub-district in Trenggalek and Ponorogo districts equal to exp?0.0966? ?1.1011 times, assuming the other variables are constant. The interpretation of the other independent variables can carry out in the same way. The Poisson state regression model for AIDS as follows. log??^2i ? ? 1.9625? 0.0187X1i ? 0.0560X2i ?0.0141X3i ? ?0.0103X4i ? 0.0221X5i Like the first model above, can interpret every 1% increase of population aged 25-29 years ? X1? will reduce the average number of AIDS cases in each sub-district in Trenggalek and Ponorogo districts equal to exp??0.187? 20.9815 times, which assuming the other variables are constant. The interpretation of the other independent variables can carry out in the same way. There are differences influence of variables by Wijaya, where in Wijaya's research there are 3 variables that have a positive effect on the number of HIV and AIDS, namely the percentage of age group 25-29 years, The percentage of couples of reproductive age using condoms, and The percentage of community health insurance (Jamkesmas), while the other 2 variables have a negative effect. In this study only the percentage of population with senior high school level has a positive effect (it can increase the number of HIV and AIDS as the percentage of this variable increases), while other variables have a negative effect. The second model of BZIPIGR is zero state regression model for HIV and AIDS. The zero-state regression model for HIV is as follows: logit ? p^1i ? ? ?0.05? ?1.1169 ? 0.0966X1i ? 0.0643X2i ? 0.0423X3i ? ? 0.2703X4i ? 0.0150X5i ? = 0.0558+0.0048X1i ? 0.0032X2i ? 0.0021X3i ? 0.0135X4i ? 0.0008X5i The logit model equation above can interpret the Opportunities in every district in Trenggalek and Ponorogo did not have HIV cases increased equal to exp (0,0048)=1.0048 times if there is an increase of 1% of the population aged 25-29 years (X1), assuming other variables constant. Similarly, to the variable X3, X4, and X5. However, if there is an increase of 1% of percentage of the population with senior high school level X2 with the assumption that the other variables are constant, then the chance of not having an HIV case will decrease by exp (0.0032)= 1.0032 times. This is related with the Poisson state regression model, which can increase the number of HIV is variable X2. The zero-state regression model for AIDS cases is as follows. logit?p^2i???0.057??1.9625?0.0187X1i ?0.0560X2i ?0.0141X3i ? ?0.0103X4i ?0.0221X5i? =0.1119+0.0011X1i 20.0032X2i 20.0008X3i 20.0006X4i 20.0013X5i The logit model equation above can interpret the chance that each sub-district in Trenggalek and Ponorogo districts does not have AIDS cases increases by exp?0.0011?? 1.0011 times if there is an increase of 1% of the population aged 25- 29 years, assuming the other variables are constant. Similarly, to the variable X3, X4, and X5. However, if there is an increase of 1% of percentage of the population with senior high school level X2 with the assumption that the other variables are constant, then the chance of not having an AIDS case will decrease by exp (0.0032)= 1.0032 times. This is related to the Poisson state regression model, which can increase the number of AIDS variables X2. Based on these two regression models (Poisson state and zero state), an increase in the number of HIV and AIDS can be caused by an increase in the percentage of the population aged 2 years (X2). According to Akaike Information Criterion Corrected (AICc) value. The AICc value uses if the objective of regression modeling is to identify the influencing factors. The AICc value using the BZIPIGR model obtained is 317.96. Previous research with the same data using the BZIPR method on the number of HIV and AIDS cases got an AICc value of 340.6977. Therefore, the data on the number of HIV and AIDS cases in Trenggalek and Ponorogo Districts is better using the BZIPIGR model. IV. CONCLUSION Based on the results and discussion above, it can conclude that: The estimation parameters of the BZIPIGR model using the MLE method has obtained that the first derivative does not yield closed form. Therefore, it is followed by numerical iteration using the Berndt Hall Hall Hausman (BHHH) algorithm. For Simultaneous hypothesis testing of the BZIPIGR model using the MLRT method has obtained test statistics that follow Chi-square distribution. Based on the AICc value were obtained from the BZIPIGR model, it has shown that the model is feasible to apply for data on the number of HIV and AIDS cases. All predictor variables significantly affected the number of HIV and AIDS cases. Based on the Poisson state regression model, the variable whose effect is to increase the number of HIV and AIDS cases is the percentage of the population with low education (SMA). The variables that can reduce the number of HIV and AIDS cases are the percentage of the population aged 25-29 years, the percentage of reproductive-age couples (PUS) using condoms, the percentage of health education activities, and the percentage of community <u>health insurance (Jamkesmas</u>). Based on the Zero state regression model, REFERENCES [1] R. Fitriani, L. N. Chrisdiana, and A. Efendi, "Simulation on the Zero Inflated Negative Binomial (ZINB) to Model Overdispersed, Poisson Distributed Data," IOP Conf. Ser. Mater. Sci. Eng., vol. 546, no. 5, 2019, doi: 10.1088/1757-899X/546/5/052025. [2] R. D. Kusuma and Y. Purwono, "Zero-Inflated Poisson Regression Analysis On Frequency Of Health Insurance Claim PT. XYZ," in Proceedings of the 12th International Conference on Business and Management Research (ICBMR 2018), 2019, vol. 72, no. Icbmr 2018, pp. 321-325, doi: 10.2991/icbmr-18.2019.52. [3] D. Karlis and E. Xekalaki, "Mixed Poisson distributions," Int. Stat. Rev., vol. 73, no. 1, pp. 35–58, 2005, doi: 10.1111/j.1751- 5823.2005.tb00250.x. [4] Purhadi, Sutikno, S. M. Berliana, and D. I. Setiawan, "Geographically weighted bivariate generalized Poisson regression: application to infant and maternal mortality data," Lett. Spat. Resour. Sci., vol. 14, no. 1, pp. 79–99, 2021, doi: 10.1007/s12076-021-00266-5. [5] L. Amaliana, U. Sa'Adah, and N. W. Surya Wardhani, "Modeling Tetanus Neonatorum case using the regression of negative binomial and zero-inflated negative binomial," J. Phys. Conf. Ser., vol. 943, no. 1, 2018, doi: 10.1088/1742-6596/943/1/012051. [6] S. Mardalena, Purhadi, J. T. D. Purnomo, and D. D. Prastyo, "Bivariate Poisson Inverse Gaussian Regression Model with Exposure Variable: Infant and Maternal Death Case Study," J. Phys. Conf. Ser., vol. 1752, no. 1, 2021, doi: 10.1088/1742-6596/1752/1/012016. [7] X. F. Huang, G. L. Tian, C. Zhang, and X. Jiang, "Type I multivariate zero-inflated generalized Poisson distribution with applications," Stat. Interface, vol. 10, no. 2, pp. 291-311, 2017, doi: 10.4310/SII.2017.v10.n2.a12. [8] M. S. Nur, Purhadi, and A. Choiruddin, "Parameter

https://www.turnitin.com/newreport printview.asp?eq=0&eb=1&esm=-1&oid=2087684534&sid=0&n=0&m=2&svr=29&r=95.10892275514458&lan... 4/5

Turnitin - Originality Report - 1._14217-38137-1-PB.pdf

Estimation and Hypothesis Testing of Geographically Weighted Bivariate Zero Inflated Poisson Inverse Gaussian Regression Models," IOP Conf. Ser. Mater. Sci. Eng., vol. 1115, no. 1, p. 012043, 2021, doi: 10.1088/1757 899x/1115/1/012043. [9] J. M. Hilbe, Cambridge Books Online, vol. 48, no. 04. 2010. [10] Q. Aini, Purhadi, and Irhamah, "Bivariate zero inflated generalized Poisson regression model in the number of pregnant maternal mortality and the number of postpartum maternal mortality in the Central Java Province in 2017," J. Phys. Conf. Ser., vol. 1511, no. 1, 2020, doi: 10.1088/1742-6596/1511/1/012055. [11] Dinas kesehatan Provinsi Jawa Timur, Profil Kesehatan Provinsi Jawa Timur Tahun 2018. Jawa Timur, Indonesia: Dinas Kesehatan Provinsi Jawa Timur, 2019. [12] Kementerian Kesehatan Republik Indonesia, "Infodatin (Situasi HIV dan AIDS di Indonesia)," Pusdatin, 2020, [Online]. Available: https://pusdatin.kemkes.go.id/resources/download/pusdatin/infodatin/ infodatin-2020-HIV.pdf. [13] I. Purnamasari and I. N. Latra, "Parameter Estimation and Statistical Test in Modeling Geographically Weighted Poisson Inverse Gaussian Regression," in International Conference Research: Implementation and Education of Mathematics and Science, 2016, no. May, pp. 16-17. [14] S. Mardalena, P. Purhadi, J. D. T. Purnomo, and D. D. Prastyo, "Parameter Estimation and Hypothesis Testing of Multivariate Poisson Inverse Gaussian Regression," Symmetry (Basel)., vol. 12, no. 10, p. 1738, Oct. 2020, doi: 10.3390/sym12101738. [15] J. Amalia, Purhadi, and B. W. Otok, "Application of geographically weighted bivariate poisson inverse Gaussian regression," AIP Conf. Proc., vol. 2268, no. September 2020, doi: 10.1063/5.0017393. [16] J. Pangulimang, "Parameter Estimation and Hypothesis Testing Geographically Weighted Bivariate Zero-Inflated Poisson," in International Conference Research: Implementation and Education of Mathematics and Science, 2016, no. May, pp. 16-17, [Online]. Available: http://seminar.uny.ac.id/icriems/sites/seminar.uny.ac.id.icriems/files/ prosiding/M-12.pdf. [17] S. M. Berliana, Purhadi, Sutikno, and S. P. Rahayu, "Parameter estimation and hypothesis testing of geographically weighted multivariate generalized poisson regression," Mathematics, vol. 8, no. 9, pp. 1-14, 2020, doi: 10.3390/math8091523. [18] J. N. Gonçalves and W. Barreto-Souza, "Flexible regression models for counts with high-inflation of zeros," Metron, vol. 78, no. 1, pp. 71– 95, 2020, doi: 10.1007/s40300-020-00163-9. [19] H. Cho, T. Porras, D. Baik, M. Beauchemin, and R. Schnall, "Understanding the predisposing, enabling, and reinforcing factors influencing the use of a mobile-based HIV management app: A real- world usability evaluation," Int. J. Med. Inform., vol. 117, pp. 88-95, 2018, doi: 10.1016/j.ijmedinf.2018.06.007. [20] S. U. Wijaya, Pendugaan Parameter dan Pengujian Hipotesis pada Bivariate Poisson Inverse Gaussian Regression (Studi Kasus: Jumlah Kasus Baru HIV dan AIDS di Kabupaten Trenggalek dan Ponorogo Tahun 2012). Indonesia: Repository ITS, 2017. [21] S. M. Berliana, Purhadi, Sutikno, and S. P. Rahayu, "Multivariate generalized Poisson regression model with exposure and correlation as a function of covariates: Parameter estimation and hypothesis testing," AIP Conf. Proc., vol. 2192, 2019, doi: 10.1063/1.5139171. [22] N. Shrestha, "Detecting Multicollinearity in Regression Analysis," Am. J. Appl. Math. Stat., vol. 8, no. 2, pp. 39-42, 2020, doi: 10.12691/ajams-8-2-1. 2407 2408 2409 2410 2411 2412 2413 2414 2415