

Turnitin Originality Report

Processed on: 08-May-2023 23:07 WIB
ID: 2087684543
Word Count: 8640
Submitted: 1

2._symmetry-14-00277-v2.pdf By Anonymous

Similarity by Source	
Similarity Index	Internet Sources: 11%
17%	Publications: 14%
	Student Papers: 6%

5% match (Internet from 08-May-2023)

[2% match \(Selvi Mardalena, Purhadi Purhadi, Jerry Dwi Trijoyo Purnomo, Dedy Dwi Prastyo. "Parameter Estimation and Hypothesis Testing of Multivariate Poisson Inverse Gaussian Regression", Symmetry, 2020\)](https://watermark.silverchair.com/030021_1_online.pdf?token=AQECAHi208BE490oan9kkhW_Ercy7Dm3ZL_9CF3qfKAc485ysgAABRQwggUQBgkqhkig9w0BBWagggUBMIIE_QIBADCCBPYGCSqGSIb_416SXvb4Rgk4f0vSNeGgAXMgDNe9wI71TL7ZPmMBwrB-R_bWX7kACa64IF3iaftPjYC-XgxWJNkl_Wjc2WB0-naA37ZG-sn9WKB-NYQi2Yt6dGyJ9G5GQdSAGVrk96Or0c0BYUzApRs-pOn5W9Qfp-e_sZJqon230iUP8zHm4ESkl2S9i7EX3yM9dvMBYdCUMGcXbWA20Q8fMWXBWlqWmcXdoA7nHjTj4lXwH4DaK2rIlWjB6exxdAbw-_VmsjNxfis3GhN7bJ8EIJkvfUsXBac3MUAUV2bc9E8TDviIHhXQcwIf6_rMbMronnyu5PjBY_cmepMWloPg36etPludNKJ-rXQzs-Jme0W_5FG71tKga6W2gJXr9wnoncYAmGnk2FoUlca8lxymwddLI7VLWlqVu4THIDeGtAjKJTffKwPdsB9sQxseJrJPHeZpAfGM_kPCuwZyzL0Uc-wdTA_AYUJ3untWruFa89W4nzSDIERFVfob2X3g7AIIGjOSZhbDmPnGYH3-W6lzkisMJ_m_A3FvKpXGrXhLyz1NuVzUUi766m4f5BBFtgn9-yEoVqxOdWhbouzGmml6AX0CuFoDg7wSYIoBtx_ZKehUJa-aVSgUUNZJjaqht5Wyg4JowGa9GFvt3VTg3L0ecN9EJMSqoKFQyfQpXGPUSro-RcqayGKMYSmBfpoUY6fu2g56jL3EyyyzVduyHeco4pZQAGYj3FVH2t_Itd9HxKk7ySKsCyyUV-XOcEediQy7KcJxF9Dz3ea8J-nUDr9kMHN6wGmmu0oPTKUD45ZCJDG802QXgIBQWYwnyP_rfqNXefaeLZh-66lzRcYFKOcl0rBly0Z6WSajkQJITd6EctUZ2RMHv32jR7WFWVlv18v5OXCod2WXWVVB8g6a4bL1-VEqVwnQtCCTITf4RW3YDUqXTFCp3cid1UkBJIGRbXWVxWlNhQgDg_wyMqSMqsC7qMdT-aAWXrQGc-gCjp4ampeBptU101GL7K7DJNhUisL_yXU-Ol61qgZfMsMW7HM7Wct_4EspVNHcFe01VpoDzJsdnlicUFai3tv7Gx2kLwq3QwFsnHji-9L3x0yzE-9GZupdU4FgLEqHtA3Ec_u-9hDvZwzPwErcWPMoHMJ6ja6A9VmS_YCWCGZUDYANbdZMVXp7w8lGeSa6cPSBFF1a9sbDXvbeE5RwFpFbUTHzW6wHI_qmFarxJBJIgb6SX4T-RWT4dxJZnt9VBIXybEMCSivFXT1sB7ZkwgD_7KNv24Lf99xqFVPhdiSOXpHWR8i2g4_2CYx7PNuJ5KifhRNbnNeu9Tz-TeLeL-fVGvz5-2AXJhUEU6bolVvwWQuK77s55aR1zmdLU0eDzIrDQyUAgZpBr9XR932eTylUzFXwk0SMe43WDXuHleM-16soIAY</p>
</div>
<div data-bbox=)

[Selvi Mardalena, Purhadi Purhadi, Jerry Dwi Trijoyo Purnomo, Dedy Dwi Prastyo. "Parameter Estimation and Hypothesis Testing of Multivariate Poisson Inverse Gaussian Regression", Symmetry, 2020](#)

2% match (Internet from 29-Apr-2023)

<https://www.biorxiv.org/content/10.1101/2021.02.10.430561v2.full.pdf>

1% match (Dewi Novita Sari, Purhadi Purhadi, Santi Puteri Rahayu, Irhamah Irhamah. "Estimation and Hypothesis Testing for the Parameters of Multivariate Zero Inflated Generalized Poisson Regression Model", Symmetry, 2021)

[Dewi Novita Sari, Purhadi Purhadi, Santi Puteri Rahayu, Irhamah Irhamah. "Estimation and Hypothesis Testing for the Parameters of Multivariate Zero Inflated Generalized Poisson Regression Model", Symmetry, 2021](#)

1% match (student papers from 27-Apr-2023)

[Submitted to Universitas Airlangga on 2023-04-27](#)

1% match (Selvi Mardalena, Purhadi Purhadi, Jerry Dwi Trijoyo Purnomo, Dedy Dwi Prastyo. "The Geographically Weighted Multivariate Poisson Inverse Gaussian Regression Model and Its Applications", Applied Sciences, 2022)

[Selvi Mardalena, Purhadi Purhadi, Jerry Dwi Trijoyo Purnomo, Dedy Dwi Prastyo. "The Geographically Weighted Multivariate Poisson Inverse Gaussian Regression Model and Its Applications", Applied Sciences, 2022](#)

1% match (Purhadi, Dewi Novita Sari, Qurotul Aini, Irhamah. "Geographically weighted bivariate zero inflated generalized Poisson regression model and its application", Heliyon, 2021)

[Purhadi, Dewi Novita Sari, Qurotul Aini, Irhamah. "Geographically weighted bivariate zero inflated generalized Poisson regression model and its application", Heliyon, 2021](#)

1% match (Internet from 25-Apr-2023)

<https://www.mdpi.com/2071-1050/14/13/7949>

1% match (Dewi Novita Sari, Purhadi, Santi Puteri Rahayu, Irhamah. "Bivariate zero-inflated generalized Poisson regression on modelling stillbirth and maternal death", AIP Publishing, 2022)

[Dewi Novita Sari, Purhadi, Santi Puteri Rahayu, Irhamah. "Bivariate zero-inflated generalized Poisson regression on modelling stillbirth and maternal death", AIP Publishing, 2022](#)

1% match (Internet from 17-Feb-2023)

<https://journal.unhas.ac.id/index.php/ESTIMASI/article/view/24113>

1% match (M S Nur, Purhadi, A Choiruddin. "Parameter Estimation and Hypothesis Testing of Geographically Weighted Bivariate Zero Inflated Poisson Inverse Gaussian Regression Models", IOP Conference Series: Materials Science and Engineering, 2021)

[M S Nur, Purhadi, A Choiruddin. "Parameter Estimation and Hypothesis Testing of Geographically Weighted Bivariate Zero Inflated Poisson Inverse Gaussian Regression Models", IOP Conference Series: Materials Science and Engineering, 2021](#)

1% match (student papers from 26-Jun-2020)

[Submitted to State Islamic University of Alauddin Makassar on 2020-06-26](#)

1% match (Internet from 18-Mar-2023)

<https://resmilitaris.net/menu-script/index.php/resmilitaris/article/download/3136/2504/4358>

S Symmetry Article A Comparison of Bivariate Zero-Inflated Poisson Inverse Gaussian Regression Models with and without Exposure Variables Ermawati [Ermawati 1,2](#), [Purhadi Purhadi 1,*](#) and [Santi Puteri Rahayu 1](#) 1 Faculty of Science and Data Analytics, Department of Statistics, Institut Teknologi Sepuluh Nopember, Surabaya 60111, Indonesia; ermawati@uin-alauddin.ac.id (E.E.); santi_pr@statistika.its.ac.id (S.P.R.) 2 Faculty of Science and Technology,

Department of Mathematics, Universitas Islam Negeri Alauddin Makassar, Makassar 92118, Indonesia * Correspondence: purhadi@statistika.its.ac.id Abstract: In this paper, we focus on the comparison of the [bivariate zero-inflated Poisson inverse Gaussian regression \(BZIPIGR\) type II model](#) in two cases: with and without exposure variables. The BZIPIGR type II model is applied to analyze the occurrence of [maternal and early neonatal mortality in South Sulawesi](#) Province, Indonesia using [2019](#) data, which contain many zero values and have the issue of overdispersion in the response variable. Furthermore, to analyze the number of deaths in various areas, the exposure variable is considered. The maximum likelihood estimation (MLE) is used in parameter estimation, which involves numerical iteration and application of the Berndt-Hall-Hausman (BHHH) algorithm. Sum square error (SSE) serves as the criterion of model selection when exposure variables are included. The existence of exposure variables strongly affects the model's accuracy, especially using the BZIPIGR type II model. According to the SSE and RMSE values, the BZIPIGR type II [model with exposure](#) variables performs [better than the model without exposure](#) variables in estimating parameter values. All predictors with exposure variables in [????????? ??????](#) this study had a significant influence on the number of maternal and early neonatal mortalities. Citation: Ermawati, E.; Purhadi, P.; Keywords: BZIPIGR type II; overdispersion; excess zero; exposure variable; MLE; BHHH; number Rahayu, S.P. A Comparison of mortalities Bivariate Zero-Inflated Poisson Inverse Gaussian Regression Models with and without Exposure Variables. *Symmetry* 2022, 14, 277. <https://doi.org/10.3390/sym14020277> Count data that follow a Poisson distribution can be modeled by Poisson regression. Academic Editor: Dumitru Baleanu [Poisson regression requires the assumption that the mean and variance of the response](#) Received: 12 December 2021 variables must be equal (equidispersion), i.e., one parameter gives both the mean and Accepted: 20 January 2022 variance of the distribution. [In reality, this assumption may not hold, either with a variance](#) Published: 29 January 2022 [larger than the mean \(overdispersion\) or otherwise \(underdispersion\)](#) [1–5]. [Such a violation can result in errors in decision making in hypothesis testing due to the occurrence of](#) Publisher's Note: MDPI stays neutral underestimates [6–8]. A model was developed based on mixed Poisson distribution to [with regard to jurisdictional claims in published maps and institutional](#) affil- overcome [the](#) overdispersion problem, constituting a combination of Poisson distribution iations. and other distributions, both discrete and continuous, including Poisson Inverse Gaussian (PIG) distribution [9]. Poisson regression can be used to model count data, but between events within a certain period, the data must be independent. For example, maternal mortality at a Copyright: © 2022 by the authors. Switzerland. number of mothers who die within a certain period in a given area may be influenced This article is an open access article by other factors; for example, more births results in a higher occurrence of maternal distributed under the terms and mortality in an area. Thus, when comparing the number of mortalities between one conditions of the Creative Commons area and another, it is necessary to consider the existence of other variables that can Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>). affect the number of occurrences, referred to as the exposure variable [10]. Frome and creativecommons.org/licenses/by/ Morris (1989) [11] conducted research involving exposure variables with Poisson regression 4.0/). modeling. Violation of the equidispersion assumption, especially overdispersion within *Symmetry* 2022, 14, 277. <https://doi.org/10.3390/sym14020277> <https://www.mdpi.com/journal/symmetry> the Poisson distribution, can be caused by the number of [zero values \(excess zero\) in the response variable](#) [11–13]. Phang and Loh recommend a [zero-inflated Poisson inverse Gaussian regression \(ZIP- IGR\) model for data with excess zeros and overdispersion](#) [14]. [Zero-inflated](#) count data contains many zeros in the response variable. Hilbe conducted simulations with data containing excess zeros and overdispersion. Several models were included in this simu- lation, such as zero-inflated negative binomial regression (ZINBR), zero-inflated hurdle negative binomial regression (ZIHNB) and zero-inflated generalized regression (ZIGPR). The simulation obtained the lowest [AIC and BIC values with the ZIPIGR model, so](#) the conclusion is that the ZIPIGR model is preferable to the other models [5]. Several other researchers have also used ZIPIGR to model with count data containing excess zeros and overdispersion [15,16]. The ZIPIGR model is limited to univariate cases; therefore, if the relevant application has more than one response variable, excess zeros or overdispersion, this model cannot be used. In this study, we used two correlated response variables: the number of maternal and early neonatal mortalities. Maternal mortality markedly affects the survival of the child, especially in cases of babies younger than one month old (early neonatal), where the basic needs of the baby are dependent on the mother, such as breast feeding. More than 50% of response variables in this study contained zero values and exhibited overdispersion. Therefore, a bivariate zero-inflated Poisson inverse Gaussian regression (BZIPIGR) model was developed. The BZIPIGR type I model consists of two probability functions that accommodate the state of the response-variable pairs: all response variables are zero, and all response variables are not zero. However, not all cases consist of two response conditions. In the case of [the number of maternal and early neonatal mortalities in South Sulawesi](#) Province [in](#) 2018, there were four combinations of zero values in the response variable. Therefore, to identify [the factors that significantly](#) affected [the number of maternal and early neonatal mortalities](#), a BZIPIGR type II model was developed to accommodate all possible combinations of zero values in the response variable. This is similar to the formation of the probability function in the BZIGPR model [17]. Several researchers have [used Newton-Raphson iterations to maximize the likelihood function using the MLE method when the derivative of the likelihood function for the model parameters was in non-closed form. This algorithm requires a second derivative to form a Hessian matrix. Therefore, for a more complex probability distribution function, it is difficult to obtain a second derivative.](#) According to Nugraha, [the first-derivative approach can be used to inform the Hessian matrix using a Berndt-Hall-Hausman \(BHHH\) iteration](#) [18]. [The complexity of the probability function in the BZIPIGR type II model makes it difficult to obtain the second derivative.](#) Therefore, we used the BHHH iteration method as an alternative to overcome this problem. 2. [Bivariate Zero-Inflated Poisson Inverse Gaussian Regression Type II \(BZIPIGR\)](#) The BZIPIGR type II model [is](#) derived from the ZIPIGR model. The ZIPIGR model is a mixed of [zero-inflated \(ZI\) and Poisson inverse Gaussian \(PIG\) distributions](#) [16]. The zero value in the ZIPIG distribution can be divided into two sources, namely structural zero (zero state) with probability p_0 and sampling zero (mixed Poisson distribution state or PIG state) with probability $(1 - p_0)$. If there are two random variables, Y_1 and Y_2 , with BZIPIG distri- bution with parameters (η_h, γ_h, τ) where $h = 1, 2$ or $(Y_1, Y_2) \sim BZIPIG(\eta_1, \eta_2, \tau, \gamma_1, \gamma_2)$, and where η_1 and η_2 are the parameters of the Poisson distribution; τ [is the dispersion parameter, and](#) γ_1, γ_2 [is the](#) parameter of the [zero-inflated](#) distribution, [the probability function of](#) Y_1 and Y_2 can be obtained with: $(Y_1, Y_2) \sim (0, 0)$ with p_0 probability $\sim (PIG(\eta_1, \tau), 0)$ with p_1 probability $\sim (0, PIG(\eta_2, \tau))$ with p_2 probability $\sim (BPiG(\eta_1, \eta_2, \tau))$ with p_3 probability where $p_3 = 1 - p_2 - p_1 - p_0$. Hence, [the joint probability function of](#) Y_1 and Y_2 [can be written as](#) F_1 ; if $y_1 = 0, y_2 = 0$ $P(Y_1 = y_1, Y_2 = y_2) = \begin{cases} F_2; & \text{if } y_1 = 0, y_2 > 0 \\ F_3; & \text{if } y_1 > 0, y_2 = 0 \\ F_4; & \text{if } y_1 > 0, y_2 > 0 \end{cases}$ where $F_1 = \gamma_1 \gamma_2 + \gamma_1(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 - 2} K_{-12}(\sqrt{2\tau(\eta_1 + \eta_2)}) \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (2) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (3) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (4) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (5) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (6) \gamma_1 \gamma_2 + \gamma_1(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 - 2} K_{-12}(\sqrt{2\tau(\eta_1 + \eta_2)}) \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (7) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (8) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (9) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (10) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (11) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (12) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (13) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (14) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (15) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (16) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (17) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (18) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (19) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (20) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (21) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (22) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (23) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (24) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (25) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (26) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (27) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (28) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (29) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (30) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (31) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (32) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (33) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (34) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (35) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (36) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (37) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (38) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (39) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (40) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (41) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (42) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (43) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (44) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (45) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (46) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (47) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (48) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (49) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (50) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (51) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (52) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (53) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (54) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (55) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (56) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (57) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (58) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (59) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (60) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (61) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (62) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (63) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (64) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (65) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (66) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (67) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (68) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (69) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (70) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (71) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (72) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (73) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (74) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (75) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (76) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (77) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (78) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (79) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (80) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (81) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (82) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (83) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (84) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (85) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau) \tau^{\eta_2} (1 + 2\sqrt{2\tau\eta_1 - 12}) (1 + 2\sqrt{2\tau\eta_1 - 12})^{-1} \tau + (86) (1 - \gamma_1)(1 - \gamma_2) \exp(-\tau) \tau^{\eta_1 + 2\tau\eta_2 + \gamma_2(1 - \gamma_1)} \exp(\tau$

$$\partial \log L_4(\boldsymbol{\tau}) / \partial \boldsymbol{\tau} = \sum_{n=1}^4 \left(\left(-1 + \frac{1}{1 + \gamma_1 + \gamma_2} \left(2(q_1 \exp(x_i \tau_1) + q_2 \exp(x_i \tau_2)) - 1 \right) - 2 + i = 1 \right) \tau_2 \right) \left(\frac{1}{1 + 2\tau(q_1 \exp(x_i \tau_1) + q_2 \exp(x_i \tau_2))} \right) \tau_2 \gamma_1 > 0, \gamma_2 > 0 \left(-4K\gamma_1 + \gamma_2 + 21(z_4) + (\gamma_1 + \gamma_2 - 12)Ks_4(z_4) \right) \left(-1 + \tau(q_1 \exp(x_i \tau_1) + q_2 \exp(x_i \tau_2)) \right) K \gamma_1 + \gamma_2 - 12(z_4) \frac{1}{1 + 2\tau(q_1 \exp(x_i \tau_1) + q_2 \exp(x_i \tau_2))} \right) \left| \right. \text{The first derivative of the likelihood for the model without/exposure variables is like the BZIPIGR type II model with exposure variables, but } q_i \text{ is not involved in the equation above.}$$

References 1. Sari, D.N.; Purhadi, P.; Rahayu, S.P.; Irhamah, I. Estimation and Hypothesis Testing for the Parameters of Multivariate Zero Inflated Generalized Poisson Regression Model. *Symmetry* 2021, 13, 1876. [CrossRef] 2. Zha, L.; Lord, D.; Zou, Y. The Poisson Inverse Gaussian (PIG) Generalized Linear Regression Model for Analyzing Motor Vehicle Crash Data. *J. Transp. Saf. Secur.* 2014, 8, 18–35. [CrossRef] 3. Famoye, F.; Sighn, K.P. Zero-Inflated Generalized Poisson Regression Model with an Application to Domestic Violence Data. *J. Data Sci.* 2006, 4, 117–130. [CrossRef] 4. Purhadi; Sari, D.N.; Aini, Q.; Irhamah. Geographically weighted Bivariate Zero Inflated Generalized Poisson Regression. *Heliyon* 2021, 7, e07491. [CrossRef] [PubMed] 5. Cameron, A.C.; Trivedi, P.K. *Regression Analysis of Count Data*; Cambridge University Press: New York, NY, USA, 1998. 6. Hilbe, J.M. *Poisson Inverse Gaussian Regression. In Modeling Count Data*; Cambridge University Press: New York, NY, USA, 2014. [CrossRef] 7. Ozmen, I.; Famoye, F. Count Regression Models with an Application to Zoological Data Containing Structural Zeros. *Data Sci. J.* 2007, 5, 491–502. [CrossRef] 8. McCullagh, P.; Nelder, J.A. *Generalized Linear Models*, 2nd ed.; Chapman and Hall: London, UK, 1989. 9. Karlis, D.; Xekalaki, E. A Simulation Comparison of Several Procedures for Testing the Poisson Assumption. *Statistician* 2000, 49, 355–382. [CrossRef] 10. Mardalena, S.; Purhadi; Purnomo, D.T.; Prastyo, D.D. Parameter Estimation and Hypothesis Testing of Multivariate Poisson Inverse Gaussian Regression. *Symmetry* 2020, 12, 1738. [CrossRef] 11. Frome, E.L.; Morris, M.D. Evaluating Goodness of Fit of Poisson Regression Model in Cohort Studies. *Am. Stat.* 1989, 43, 144–147. 12. Leffondré, K.; Jager, K.J.J.; Boucquemont, J.; Stel, V.S.; Heinze, G. Representation of exposures in regression analysis and interpretation of regression coefficients: Basic concepts and pitfalls. *Nephrol. Dial. Transplant.* 2014, 29, 1806–1814. [CrossRef] [PubMed] 13. Long, D.L.; Preisser, J.S.; Herring, A.H.; Golin, C.E. A Marginalized Zero-Inflated Poisson Regression Model with Overall Exposure Effects. *NIH Public Access* 2014, 33, 5151–5165. [CrossRef] [PubMed] 14. Phang, Y.N.; Loh, E.F. Statistical Analysis for Overdispersed Medical Count Data. *Int. J. Math. Comput. Sci.* 2014, 8, 292–294. 15. Silva, J.G. Zero-Inflated Mixed Poisson Regression Models. Master's Thesis, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil, 17 February 2017. 16. Gonzalves, J.N.; Souza, W.B. Flexible regression models for counts with high-inflation of zeros. *Metron* 2020, 78, 71–95. [CrossRef] 17. Zhang, C.; Tian, G.; Huang, X. Two New Bivariate Zero-Inflated Generalized Poisson Distribution with a Flexible Correlation Structure. *Stat. Optim. Inf. Comput.* 2015, 3, 105–137. [CrossRef] 18. Nugraha, J.; Guritno, S.; Haryatmi, S. Probit Model on Multivariate Binary Response Using Simulated Maximum Likelihood Estimator. *J. Ilmu Dasar* 2010, 11, 70–75. (In Indonesia) 19. Shoukri, M.M.; Asyali, M.H.; Kelton, D. The Poisson Inverse Gaussian Regression Model of Clustered count Data. *J. Data Sci.* 2014, 2, 17–32. [CrossRef] 20. Willmot, G.E. The Poisson-Inverse Gaussian distribution as an alternative to the negative binomial. *Scand. Actuar. J.* 1987, 1987, 113–127. [CrossRef] 21. Beren, C.; Thamrin, S.A.; Jaya, A.K. Estimasi Parameter Distribusi Exponentiated Modified Weibull Extension. In Proceedings of the National Conference Variansi, Makassar, Indonesia, 10 October 2016; ISBN 978-602-99837-2-2. 22. Mardalena, S.; Purhadi; Purnomo, D.T.; Prastyo, D.D. Bivariate Poisson Inverse Gaussian Regression Model with Exposure Variable: Infant and Maternal Death Case Study. *J. Phys. Conf. Ser.* 2021, 1752, 012016. [CrossRef] [Symmetry 2022, 14, 277 2 of 18](#) [Symmetry 2022, 14, 277 3 of 18](#) [Symmetry 2022, 14, 277 4 of 18](#) [Symmetry 2022, 14, 277 5 of 18](#) [Symmetry 2022, 14, 277 6 of 18](#) [Symmetry 2022, 14, 277 7 of 18](#) [Symmetry 2022, 14, 277 8 of 18](#) [Symmetry 2022, 14, 277 9 of 18](#) [Symmetry 2022, 14, 277 10 of 18](#) [Symmetry 2022, 14, 277 11 of 18](#) [Symmetry 2022, 14, 277 12 of 18](#) [Symmetry 2022, 14, 277 13 of 18](#) [Symmetry 2022, 14, 277 14 of 18](#) [Symmetry 2022, 14, 277 15 of 18](#) [Symmetry 2022, 14, 277 16 of 18](#) [Symmetry 2022, 14, 277 17 of 18](#) [Symmetry 2022, 14, 277 18 of 18](#)