

Turnitin Originality Report

Processed on: 08-May-2023 23:07 WIB

ID: 2087684559

Word Count: 4317

Submitted: 1

4_030021_1_online.pdf By Anonymous

Similarity Index	Similarity by Source
13%	Internet Sources: 10% Publications: 10% Student Papers: 5%

5% match (student papers from 29-Apr-2023)

[Submitted to Universitas Indonesia on 2023-04-29](#)

2% match (Internet from 12-Jan-2023)

<https://pdfcoffee.com/the-implementation-of-matematika-gasing-on-multiplication-concept-toward-interest-study-motivation-and-student-learning-outcome-pdf-free.html>

1% match (Selvi Mardalena, Purhadi, Jerry Dwi Trijoyo Purnomo, Dedy Dwi Prastyo. "Chapter 21 A Modified Inverse Gaussian Poisson Regression with an Exposure Variable to Model Infant Mortality", Springer Science and Business Media LLC, 2021)

[Selvi Mardalena, Purhadi, Jerry Dwi Trijoyo Purnomo, Dedy Dwi Prastyo. "Chapter 21 A Modified Inverse Gaussian Poisson Regression with an Exposure Variable to Model Infant Mortality", Springer Science and Business Media LLC, 2021](#)

1% match (Internet from 03-Feb-2023)

https://mdpi-res.com/d_attachment/symmetry/symmetry-13-00326/article_deploy/symmetry-13-00326-v3.pdf?version=1614329008

1% match (Ivan K. Kostov, Bénédicte Ponsot, Didina Serban. "Boundary Liouville theory and 2D quantum gravity", Nuclear Physics B, 2004)

[Ivan K. Kostov, Bénédicte Ponsot, Didina Serban. "Boundary Liouville theory and 2D quantum gravity", Nuclear Physics B, 2004](#)

1% match (Pratnya Paramitha Oktaviana, Imam Safawi Ahmad, Nuri Wahyuningsih, Yeni April Lina, Annisaa Rahmaah Nurul Syawal. "The modeling of frequency-magnitude of earthquakes in Indonesia using Poisson regression", AIP Publishing, 2022)

[Pratnya Paramitha Oktaviana, Imam Safawi Ahmad, Nuri Wahyuningsih, Yeni April Lina, Annisaa Rahmaah Nurul Syawal. "The modeling of frequency-magnitude of earthquakes in Indonesia using Poisson regression", AIP Publishing, 2022](#)

1% match (Sarni Maniar Berliana, Purhadi, Sutikno, Santi Puteri Rahayu. "Parameter Estimation and Hypothesis Testing of Geographically Weighted Multivariate Generalized Poisson Regression", Mathematics, 2020)

[Sarni Maniar Berliana, Purhadi, Sutikno, Santi Puteri Rahayu. "Parameter Estimation and Hypothesis Testing of Geographically Weighted Multivariate Generalized Poisson Regression", Mathematics, 2020](#)

1% match ()

[LIU, YAN. "Market Penetration of New Vehicle Technology: A Generalized Dynamic Approach for Modeling Discrete-Continuous Decisions", 'Wiley', 2017](#)

1% match (Internet from 12-Apr-2023)

[https://repository.iain-ternate.ac.id/id/eprint/138/1/32.%20ICosMEE%202021%20ASTUTI%20MUH.AMIN%20\(Conference%20International\).r](https://repository.iain-ternate.ac.id/id/eprint/138/1/32.%20ICosMEE%202021%20ASTUTI%20MUH.AMIN%20(Conference%20International).r)

1% match (Internet from 08-May-2023)

<https://elar.usfeu.ru/bitstream/123456789/12374/1/2-s2.0-85151386169.pdf>

RESEARCH ARTICLE | DECEMBER 19 2022 Parameter estimation and statistical test on bivariate zero inflated Poisson Inverse Gaussian with exposure variable Ermawati ?; Purhadi; Santi Puteri Rahayu AIP Conference Proceedings 2641, 030021 (2022) <https://doi.org/10.1063/5.0115074> ?? [CrossMark View](#) [Export Online Citation Articles You May Be Interested In](#) Parameter estimation and statistical test on zero inflated Poisson inverse Gaussian regression model AIP Conference Proceedings (January 2023) Bivariate binary logistic regression with fisher scoring and BHHH iteration AIP Conference Proceedings (January 2023) [Parameter estimation and hypothesis testing of multivariate adaptive bivariate Generalized Poisson Regression spline AIP Conference Proceedings](#) (January 2023) [Downloaded from http://pubs.aip.org/aip/acp/article-pdf/doi/10.1063/5.0115074/16227267/030021_1_online.pdf](#) Parameter Estimation and Statistical Test on Bivariate Zero Inflated Poisson Inverse Gaussian with Exposure Variable Ermawati_{1,2}, a), Purhadi₁, b) and Santi Puteri Rahayu₁, c) 1Department of [Statistics, Institut Teknologi Sepuluh Nopember \(ITS\), Surabaya](#), 60111, [Indonesia](#) 2Department of mathematics, Universitas Islam Negeri Alauddin Makassar, 92118, Indonesia a) Corresponding author: erma.mat@gmail.com b) purhadi@statistika.its.ac.id c) santi_pr@statistika.its.ac.id [Abstract](#). This research focused on [parameter estimation and hypothesis testing of Bivariate Zero-Inflated Poisson Inverse Gaussian Regression](#) (BZIPIGR) type II. BZIPIGR model is a mixed Poisson distribution used to

model data with excess-zero and overdispersion. The parameter estimation in this study used the [Maximum Likelihood Estimation \(MLE\) method](#). The first derivative of the log-likelihood function from the BZIPIGR model had obtained not closed-form. Therefore, to maximize the likelihood function, it had used numerical iteration, namely the [Berndt-Hall-Hall-Hausman \(BHHH\)](#). The determination of the critical area and the significance of the model parameters had used the [Maximum Likelihood Ratio Test \(MLRT\)](#). INTRODUCTION Violation of the equidispersion assumption in Poisson regression consists of underdispersion and overdispersion. Overdispersion cases are more encountered than underdispersion cases in count data. Violation of this assumption can result in errors in decision-making due to occurrences underestimates in the [standard error](#) [1]. [Several models have been developed to overcome the overdispersion](#) problem using a mixed Poisson distribution, including the Poisson Inverse Gaussian (PIG) and Negative Binomial (NB) [2]. Both of these models have been investigated and resulted that the PIGR model is better than the [Negative Binomial regression model](#) [3]. Furthermore, the Multivariate Poisson Inverse Gaussian Regression (MPIGR) model was developed which was estimated using MLE with Newton Raphson (NR) iteration [4]. Many zero values (excess zero) in the response variable can cause overdispersion in the Poisson distribution. The models that can use for data with excess zeros and overdispersion are Zero-Inflated Poisson Inverse Gaussian Regression (ZIPIGR) [5], Zero-inflated Generalized Poisson (ZIGPR) [6],[7]. Based on the simulation results conducted by Hilbe for data that has excess zero and overdispersion by comparing ZIGPR (Zero-Inflated Generalized Poisson Regression), and ZIPIGR, it is found that [the smallest AIC and BIC values in the ZIPIGR model](#) [8], so in this study will be developed ZIPIGR model. The model for two response variables can use Bivariate Zero-Inflated Poisson Inverse Gaussian Regression (BZIPIGR). The BZIPIGR is divided into two models based on the zero condition in the response variable, they are type I and type II BZIPIGR. Type I BZIPIGR only consists of two distribution functions where accommodates two conditions from each pair of response variables, where all response variables are zero or positive integers. The type II BZIPIGR consists of four distribution functions according to the possible value conditions in the response variable. [7th International Conference on Mathematics: Pure, Applied and Computation](#) AIP Conf. Proc. 2641, 030021-1-030021-11; <https://doi.org/10.1063/5.0115074> [Published by AIP Publishing. 978-0-7354-4291-7/\\$30.00](#) Several previous researchers used Newton Raphson iteration to maximize the likelihood function in the MLE method when the first derivative of the likelihood function for the model parameters was not in closed-form [9]. This Newton Raphson iteration requires the 2nd derivative to form a Hessian matrix. Therefore for more complicated distribution functions, it will be heavy to get the 2nd derivative. So to overcome the difficulty in obtaining the second derivative, the first derivative approach can use in forming the Hessian matrix by using an NR iteration development called the Berndt Hall Haussman (BHHH) iteration [10].

DESCRIPTION OF METHOD The type II BZIPIGR model is a developed ZIPIGR model, where the ZIPIGR model comes from a mixed distribution between the Zero-Inflated (ZI) and Poisson Inverse Gaussian (PIG) distributions. Suppose there are two response variables Y_1 and Y_2 which are random variables with BZIPIG distribution with parameters $(\alpha_1, \alpha_2, \tau)$ where $h=1, 2$ or $(Y_1, Y_2) \sim BZIPIG(\alpha_1, \alpha_2, \tau, p_1, p_2)$, α_1 and α_2 are parameters of Poisson model, τ is dispersion parameter, and p_1, p_2 are parameters of zero inflated model. [The joint probability function of \$Y_1\$ and \$Y_2\$ can be written as follows](#)

: 1) For $Y_1 = 0$, and $Y_2 = 0$ $P(Y_1 = 0, Y_2 = 0) = p_1 p_2 + p_1 (1 - p_2) \exp(-\tau) + p_2 (1 - p_1) \exp(-\tau) + (1 - p_1)(1 - p_2) \exp(-\tau)$ $2) \text{ For } Y_1 = 0, \text{ and } Y_2 = 1, 2, \dots P(Y_1 = 0, Y_2 = y_2) = p_1 (1 - p_2) \exp(-\tau) + p_2 (1 - p_1) \exp(-\tau) + (1 - p_1)(1 - p_2) \exp(-\tau)$ $3) \text{ For } Y_1 = 1, 2, \dots, \text{ and } Y_2 = 0 P(Y_1 = y_1, Y_2 = 0) = p_2 (1 - p_1) \exp(-\tau) + p_1 (1 - p_2) \exp(-\tau) + (1 - p_1)(1 - p_2) \exp(-\tau)$ $4) \text{ For } Y_1 = 1, 2, \dots, \text{ and } Y_2 = 1, 2, \dots P(Y_1 = y_1, Y_2 = y_2) = (1 - p_1)(1 - p_2) \exp(-\tau) + p_1 (1 - p_2) \exp(-\tau) + p_2 (1 - p_1) \exp(-\tau) + (1 - p_1)(1 - p_2) \exp(-\tau)$

With model (4) (5) $\alpha_h = q_h \exp(xT \beta_h)$; $h = 1, 2$ (6) where q_h is the exposure variable of the h -th response variable, x is the matrix of the predictor variable and the parameter β_h is $\log(\beta_h) - \text{ph}$ $\text{ph} = \exp(xT \delta_h) + \exp(xT \delta_h) + 1$ and $(1 - \text{ph}) = 1 + \exp(xT \delta_h)$ (7) RESULT AND DISCUSSION Parameter Estimation. The parameter estimation of the Type II BZIPIGR model uses the Maximum Likelihood Estimation (MLE) method. This method has been done taking n random samples, namely $(Y_{1i}, Y_{2i}, X_{1i}, X_{2i}, \dots, X_{ki})$. The likelihood function of BZIPIGR is: $n L(\beta_h, \delta_h, \tau; h = 1, 2) = \prod_{i=1}^n P(y_i | \beta_h, \delta_h, \tau; h = 1, 2)$ The Type II BZIPIGR has four independent function, so the likelihood function can be written as follows: $L(\beta_h; \tau; \delta_h; h=1,2) = (L_1(\beta_h; \tau; \delta_h; h=1,2))(L_2(\beta_h; \tau; \delta_h; h=1,2))(L_3(\beta_h; \tau; \delta_h; h=1,2))(L_4(\beta_h; \tau; \delta_h; h=1,2))$ (8) where $n_1 L_1(\beta_h; \tau; \delta_h; h=1,2) = \prod P(y_i | \beta_h, \delta_h, \tau, h=1,2) = \prod_{i=1}^{n_1} P(y_i = 0, y_{2i} = 0) = \prod_{i=1}^{n_1} P(y_i | \beta_h, \delta_h, \tau, h=1,2) = \prod_{i=1}^{n_1} P(y_i = 0, y_{2i} > 0) = \prod_{i=1}^{n_1} L_3(\beta_h; \tau; \delta_h; h=1,2) = \prod_{i=1}^{n_1} P(y_i | \beta_h, \delta_h, \tau, h=1,2) = \prod_{i=1}^{n_1} P(y_i > 0, y_{2i} = 0) = \prod_{i=1}^{n_1} L_4(\beta_h; \tau; \delta_h; h=1,2) = \prod_{i=1}^{n_1} P(y_i | \beta_h, \delta_h, \tau, h=1,2) = \prod_{i=1}^{n_1} P(y_i > 0, y_{2i} > 0)$ After getting the likelihood function as in equation (8), then the log likelihood function is determined from equation (8) so that the following equation has obtained: $\log L(\beta_h; \tau; \delta_h; h = 1, 2) = \log(L_1(\beta_h; \tau; \delta_h; h = 1, 2)) + \log(L_2(\beta_h; \tau; \delta_h; h = 1, 2)) + \log(L_3(\beta_h; \tau; \delta_h; h = 1, 2)) + \log(L_4(\beta_h; \tau; \delta_h; h = 1, 2))$ (9) If $L(\beta_h; \tau; \delta_h; h = 1, 2) = L(\bullet)$, the equation (9) becomes: $\log L(\bullet) = \log(L_1(\bullet)) + \log(L_2(\bullet)) + \log(L_3(\bullet)) + \log(L_4(\bullet))$ (10) if $\phi = (\beta_1, \beta_2, \delta_1, \delta_2, \tau)$ is the vector parameters of the BZIPIGR model, therefore to maximize the log likelihood function in equation (10), then derived from each parameter and then equated with zero, as follows: $\partial \log L(\bullet) / \partial \log L_1(\bullet) = \partial \log L_2(\bullet) / \partial \log L_3(\bullet) = \partial \log L_4(\bullet) / \partial \phi = \partial \phi + \partial \phi + \partial \phi + \partial \phi$ (11) The first derivative of the first log likelihood $(\log L_1(\bullet))$ function to parameter β_1 is: $\partial \log L_1(\bullet) / \partial \beta_1 = \sum_{i=1}^{n_1} \exp(-x_i T \beta_1) x_i - \sum_{i=1}^{n_1} \exp(-x_i T \beta_1) x_i^2 = \sum_{i=1}^{n_1} x_i \exp(-x_i T \beta_1) - \sum_{i=1}^{n_1} x_i^2 \exp(-x_i T \beta_1)$ (12) $\partial \log L_2(\bullet) / \partial \beta_1 = \sum_{i=1}^{n_2} \exp(-x_i T \beta_1) x_i - \sum_{i=1}^{n_2} \exp(-x_i T \beta_1) x_i^2 = \sum_{i=1}^{n_2} x_i \exp(-x_i T \beta_1) - \sum_{i=1}^{n_2} x_i^2 \exp(-x_i T \beta_1)$ (13) $\partial \log L_3(\bullet) / \partial \beta_1 = \sum_{i=1}^{n_3} \exp(-x_i T \beta_1) x_i - \sum_{i=1}^{n_3} \exp(-x_i T \beta_1) x_i^2 = \sum_{i=1}^{n_3} x_i \exp(-x_i T \beta_1) - \sum_{i=1}^{n_3} x_i^2 \exp(-x_i T \beta_1)$ (14) $\partial \log L_4(\bullet) / \partial \beta_1 = \sum_{i=1}^{n_4} \exp(-x_i T \beta_1) x_i - \sum_{i=1}^{n_4} \exp(-x_i T \beta_1) x_i^2 = \sum_{i=1}^{n_4} x_i \exp(-x_i T \beta_1) - \sum_{i=1}^{n_4} x_i^2 \exp(-x_i T \beta_1)$ (15)

)))- - τ q1i exp (xiT β1) + ? 4 2 δβ1 ? i =1 ? y1i =0, y2i >0 ? ? (-z22) K 1 (z22) + ?? y22 - 12
 ??? K y ? ? ? (13) B24 ?? y2i + 2 ? 2i - 12 (z22) ??τ q1i exp (xTi β1) ???? 1 + 2τ (q1i exp (xiT β1
) + q2i exp (xTi β 2)) ? ? ? ? ? ? ? ? Where A2 = A21A22A23A24A25 1 1 1 + 2τ q2i exp (xiT β 2))
 4 - y2i A21 = (1+exp(xiTδx1p))(x(1Tiδ+1e)xp(xiTδ2)); A22 =???n2τ ???2 y2-i!1; A23 =(2 A24 =
 q2yi2i exp (xiT β 2 y2i + τ -1) ; A25 = K s2 (z21) ; z21 = 1 + 2τ q2i exp (xiT β 2) 1 τ B2 =
 B21B22B23B24B25 1 1 y1i 1 B21 = (1+exp(xTiδ1))(1+exp(xiTδ2)); B23 =
 (1+2τ(q1iexp(xiTβ1)+q2iexp(xiTβ2)))4- ; B22 =???n2τ ???2 y2-i!1; 2 B24 = q2yi2i exp (xiT β 2 y2i
 + τ -1) ; B25 = K s2 (z22) ; z22 = 1 + 2τ (q1i exp (xiT β1) + q2i exp (xiT β 2)) 1 τ The first
 derivative of log L3 (•) function to parameter β1 is: ? 1 ∂ log L3 (•) n3 1 ? ? ∂β1T = ? ? A3 + B3 ???
 A31A32 ? ? ? A34 A35 ??? 2 ? - y1i ???τ q1i exp (xiT β1) xi ? 3 y1i + A33 A34 xi y1i A35 + i=1 y1i
 >0,y2i =0 ? + 2 ? ? ? (1 + 2τ q1i exp (xTi β1)) 4 A313A+342τq11ieexpp((xxTiTββ11))xi ????
 -z31Ky1i+1 (z31)+???y1i - 12???K y1i - 1 (z31)?????? + 2 2 ? ? B31B32 ? ? B34 B35 ??? 12 - y1i
 ??τ q1i exp (xiT β1) xi ? ? ? ? (1 + 2τ q1i exp(xiTβ1)+ q2i exp(xiTβ2)))4 3 (+ y1i +B33B34xiy1iB35
 + 2 1 + 2τ (qB1i33eBx3p4τ(qx1Tii βe1x)p+(xqiT2iβe1x)px(i xTi β 2)) ??? - z32 ? K y2i + 2 1 (z32
) + ?? y2i - 12 ??? K y2i - 12 (z32) ????????????? (14) ? where A3 = A31A32A33A34Ks3 (z31) (1
 + exp (xTiexδp2 ())x(iT1δ+2e)xp (xiTδ1)) ; A32 = ??? n2τ ??? 2 y1-i!1 ; A33 = (1 + 2τ q1i exp (xiT
 β1)) 4 1 1 y1i A31 = - 2 A34 = q1yi1i exp (xTi β1 y1i + τ -1) ; A35 = K s3 z31) (B3 =
 B31B32B33B34B35 1 B31 = (1+exp(xiTδ2))(1+exp(xiTδ1)); B32 =???n2τ ???2 1 y1-i!1; B33 =
 (1+2τ(q1iexp(xiTβ1)+q2iexp(xiTβ2)))4 1 - y1i 2 ; B34 = q1yi1i exp (xiT β1 y1i + τ -1) ; B35 = s3
 32) ; K y1i - 1 ??? τ1 K (z 2 1 + 2τ q1i exp (xTi β1) + q2i exp (xiT β 2)) ? = s3 z32) (? ? K (The
 first derivative of log L4 (•) function to parameter β1 is: ∂ log L4 (•) y1i >?0i=,1y2i >0 ?????? 14 n 4
 ? = - y1i + y2i ? 2τ q1i exp (xTi β1) xi ∂β1T 2 ? ? 1 + 2τ (q1i exp (xTi β1) + q2i exp (xiT β2))
 +y1ixi + (15) (-z4Ks4+1(z4)+ s4Ks4 (z4))τq1i exp(xiTβ1) ? x i ? K y1i+y2i- 1 2 (z4)(1+2τ (q1i
 exp(xiTβ1)+q2i exp(xiTβ2))) ? ? ? ? The first derivative of the logL1(•) function to parameter β2
 obtained as: = ?n1 ??- ? q2i exp (xTi β 2) xi ? ∂ log L1(•) ? ? ? A11 + A12 + A13 + A14 ? 1 + 2τ q2i
 exp (xiT β 2) ? A12 + A13 ∂β2T ? ? ? ? ? i=1 y1i =0, y2i =0 ? ? 1 + 2τ (q1i exp (xiT β1) + q2i exp (xiT
 β 2)) ???? ? The first derivative of log L2 (•) function to parameter β2 is: ∂ log L2 (•) n2 ? ? xi ?
 A21 A22 ?? A24 A25 ??? 14 ? 2 δβ 2T = ?? A2 + B2 ? ? - + i =1 ? ? y22i ???(1+ 2τqi exp(xiTβ2))
 -34-y2i 2τqi exp(xiTβ2) y1i =0, y2i >0 A23 A25 y2i (qi exp (xiT β 2 y2i)) y2i exp ??? τ1 ???
 A213A+224τ qqii eexpp((xxiTiT ββ22)) xi ??? - z21 y2i + z21) + ?? y2i - 12 ??? K + K 1 (? y2i -
 1 (z21 + 2 2)) ??? ??? ? ? ? ? (1 + 2τ qi (exp (xiT β1) + exp (xiT β2))) - τ qi exp (xiT β 2) + 3
 y1i B21B22 ? ?B24B25 ??? 2 - y2i 1 ? ? 4 2 B23B25y2i(qiexp(xiTβ2y2i))y2i exp???τ1???+B23B24 ?
 -z22Ky2i+1 (1z+222)τ+q??i(ye2xip-(2xiT??β1y)2i+-1ex(pz2(2x)iT??βτ2q)i)exp(xiTβ2) ? 1 ? ? ? ? ?
 K ? ? ? ? xi ??
 The first derivative of log L3 (•) function to parameter β2 is: ? ? ? ? ∂ log L3 (•) n 3 ? = ? ?? A3 + B3 ? 1 ?? B31B32 B34 ? ? ??? 12 - y1i ???τ q2i exp (xiT β
 2) xi ∂β 2T ? 3 i =1 ? (q1i exp(xTiβ q2i exp(xTiβ2)))4 + y1i B35 + y1i >0, y2i =0 ? ? ? ? ? ? (1 + 2τ
 1) + 2 ? ? - z32 K y13 + 1 (z32) + ?? y13 - 2 ?? y13 - 12 (z32) ?τ q2i exp (xiT β 2 1 ? ? ? ? B33 ?
 2 ? ?) xi ?????? ? ? ? 1 + 2τ (q1i exp (xiT β1) + q2i exp (xiT β 2)) ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? The
 first derivative of log L4 (•) function to parameter β2 is: ∂ log L4 (•) n 4 ? = ? ?? 1 ∂β 2T i=1 y1i
>0,y2i =0 ????? 2 -(y1i + y2i)????1+2τ(q1iexpxp2i(exxiTpβ(1x)iT+βq22)iexexp(xiTβ2)) + y2i xi + ?-
 z4 K ? ? ? ? y1i + y2i + 1 (z4) + ?? y1i + y2i - 1 ? y1i + y2i - 2 ? ? ? ? K 1 (z4)τq2i exp(xiTβ2 ? ?
 ? ?) xi ?? K y1i + y2i - 1 2 (z4) ???1 + 2τ (q1i exp (xiT β1) + q2i exp (xTi β2))???? ? ? ? ? where
 z 4 = τ -1 1 + 2τ qi (exp (xiT β1) + exp (xiT β 2)) The first derivative of the log likelihood
 function to parameter δ1 obtained as: ? ∂ log L(•) n1 δδ1T = xi ? ? ? ? ? A11 + A12 exp (xiT δ1) ? n2
 i=1 ? A11 + A12 + A13 + A14 - 1 + exp (xiT δ1) ? ? ? + ? (A2 - B2 exp (xiT δ1)) + ? ? ? y1i =0,
 y2i =0 ? i=1 y1i =0, y2i >0 (A2 + B2) (1 + exp (xiT γ 1)) n 3 exp (xiTδ1) (A3 + B3) n 4 - ? -
 exp (xiT δ1) ? ? i=1 y1i >0, y2i =0 (A3 + B3) (1 + exp (xTiδ1)) ? i=1 y1i >0, y2i =0 1 + exp (xiT
 δ1))???? The first derivative of the log likelihood function to parameter δ 2 is: ? ∂logδL2T1(•) =
 xi ? n1 ? ? ? ? A11 + A14 exp (xiT δ 2) ? ? n2 ?? y1i = i0=,1y2i =0 ?? A11 + A12 + A13 + A14 - 1 +
 exp (xiT δ 2) ?? - ?? exp (xiT δ 2) (A2 + B2) xi ?? + ? ? i =1 y1i =0, y2i >0 ??? (A2 + B2) (1 +
 exp (xiT δ 2)) ??? n 3 xi B3) xi ?? n 4 - ? ?? (A3 + exp (xiT δ1) 0i=,1y2i = 0 ??? (A3 + B3) (1 +
 exp (xiT δ 1))) ? - ? y1i > ? ? y1i >0i=,y12i =0 ????? 1ex+pe(xxpIT(δx2iT)δx2i) ?????????? The first
derivative of the first log likelihood function (log L1(•)) to parameter τ obtained as: ∂ log L(•) n 1 ? =
 ? ? 1 ? ? ? ? A12 ? - 2 + ? 1 1 δτ ? i =1 ?? A11 + A12 + A13 + A14 ?? ?? τ 2 y1i =0, y2i =0 τ 1 +τ
 q2rieqixep(pxIT(xβiTβ 2)) ????? + A13 ?? - 12 + ? 1 + τ qi (exp (xiT β1) + exp (xiT β 2)) ?? ?
 1 + 2τ qi (exp (xiT β1) + exp (xTi β 2)) ??? ? + A14 ??? - τ12 + 1 + τ qi exp (xiT β1) ?? ? ? ? ? τ
 τ 2 ? τ 2 1 + 2τ qi exp (xiT β1) ??? ? The first derivative of log L2 (•) function to parameter τ is: ∂
 logδτL2 (•) = ?in=21 ??? A2 1+ B2 ?? ? A21 A22 ?????? ∂∂Aτ23 ??? A24 A25 + A23 ??? ∂∂Aτ24 ???
 A25 + A23 A24 ??? ∂∂Aτ25 ?????? + y1i =0, y2i >0 B21B22 ?? ∂B23 ? ??? ∂τ ?? B24B25 + B23 ???
 ∂∂Bτ24 ??? B25 + B23B24 ??? ∂∂Aτ25 ?????????? where ∂∂Aτ23 = ??? 12 - y2i ??? (1 + 2τ q2i exp
 (xTi β 2)) - y22i q2i exp (xiT β 2) ; ∂∂Aτ24 = - (q2i exp (xTi β 2)) y2i exp ??? τ1 ???τ -2 3 4
 ∂∂Aτ25 = -τ (11++τ2τq2qi2eixexp(px(iTxiβTβ2)2)) -z21 Ky2i+1 (z21)+ ??y2i - 12???Ky2i-12
 (z21) ? ? ? ? ? ? ? ∂∂Bτ23 ??? 12 - y2i ? 1 + 2τ ((q11ii eexpp((xxiTiT ββ11)) ++ qq22ii eexpp((
 xxiTiT ββ22)))) 4 = ? ∂B24 ? 3 y2i δτ ; (+ 2) = - (q2i exp (xTi β 2)) y2i exp ??? τ1 ???τ -2 ∂B25
 = τ-(z12+2K2xy2i(+q112i(ezx2p2(x+Ti)?βy12)i+-q122i)?xpy2(i-x12Ti(βz22)2)))
 (-1+τq1iexp(xTiβ1)+q2iexp(xiTβ2))) ? K δτ The first derivative of log L3 (•) function to parameter τ
 is: ∂ log L3 (•) = n3 δτ ? ??? A3 + B3 ? ? 1 ? A31 i =1 ?????? ∂∂Aτ32 ??? A33 A34 A35 + A32 ???
 ∂∂Aτ33 ??? A34 A35 + A32 A33 ??? ∂∂Aτ34 ??? A35 + y1i >0, y2i =0 A32 A33 A34 ??? ∂∂Aτ35 ??????
 + B31B32B34 ?? ∂B32 ? ??? ∂τ ?? B33 B34 B35 + B32 ??? ∂∂Bτ33 ??? B34 B35 + B32 B33 ???
 ∂∂Bτ34 ??? B35 + B32 B33 B34 ??? ∂∂Bτ35 ?????? ?????????? where ∂∂Aτ32 = ??? n2 ??? 2 y1-i!1 ???
 - 12 ??? (τ)- 32 ; 1 (1??+22-τqy11iie??xqp1i(exxTipβ(1x)iT)34β+1)y1i ; ∂∂Aτ34 = -q1yi1i
 (exp(xTiβ1))y1i ? 1 ∂A33 δτ = 2 exp ??? τ1 ???τ -2 ∂∂Aτ35 = ? - z31K y1i + (z31) + ? y1i - 12 ???
 ? ? ? ? 1 ? ? K y1i - 1 2 2 (z31)??????-τ (11++τ2τq1qi1eixexp(px(iTxiβTi1β)1))???? ∂∂Bτ32 = ??? n2
 ???2 y1-i!1????- 12 ???(τ)-2 ; ∂∂Bτ33 = (1????+122-τ(q11ii)??
 e(xqp1i(exxiTpβ(1x)iT+βq1)2i+eqxp2i(exxiTpβ(2x)Ti)β)342+)y1i 1 3 ; 2 ∂∂Bτ34 = -q1yi1i
 (exp(xiTβ1))y1i exp????τ1???τ-2; ∂B35 ? δτ T = -τz3(21K+y21iτ+q12i((ze3x2p)+(x??Tiyβ11i)-
 +12ex???pK(y1xiIT-12β2(z)3)2))??-(1+τ (q1i exp(xiTβ1)+q2i exp(xiTβ2)))??? The first derivative of
 log L4 (•) function to parameter τ is: ∂ log L4 (•) n 4 ? δτ = ? - 1 2 + i =1 2 τ y1i >0, y2i >0 ??????
 - 2τ12 ???+???14 - y1i + y2i ???1+22(τq1(iqe1ixpex(px(TixβiT1β)1)+q+2iqe2ixepx(px(TixβiT2β)2))
 ? ? ? ?-z4K y1i + y2i + 12 (z4)+??y1i + y2i - 12???Ks4 (z4)????-(1+τ (q1i exp(xiTβ1)+q2i

Downloaded from http://pubs.aip.org/aip/acp/article-pdf/doi/10.1063/5.0115074/16227267/030021_1_online.pdf