

Rhizosphere bacteria community shift from vegetative to reproductive stage of Indonesia aromatic rice var. Pulut mandoti EMAS

Eka Sukmawaty^{a,*}, Hafsan Hafsan^a, Masriany Masriany^a, Selis Meriem^a, Kurnia Makmur^a, Widya Hastuti^{a,b}

^a Department of Biology, Faculty of Science and Technology, UIN Alauddin Makassar, 92118, Gowa, Indonesia

^b Undergraduate Student in Department of Biology, Faculty of Science and Technology, UIN Alauddin Makassar, 92118, Gowa, Indonesia

ARTICLE INFO

Keywords:

2-acetyl pyrroline (2AP)
Bacteria community
Rhizobacteria
Rice nutrition

ABSTRACT

The popular Indonesian aromatic rice, var. Pulut Mandoti Emas (PME) is produced with high economic value in Salukanan Village, Enrekang Regency, however, the best aroma quality of mandoti rice is restricted to the village. Therefore, this study aims to analyze the rhizosphere microbial group and the nutrients of the Mandoti rice paddy field in Salukanan Village at the vegetative and reproductive phases. Metagenome analysis was carried out with 16S rRNA amplification and Library preparation to examine the macro and micronutrients. The results showed that the most dominant taxa in the rhizosphere soil of vegetative mandoti rice was phylum Acidobacteria (35%), Proteobacteria (30%), and Firmicutes (12%), while in the reproductive phase, it was Phylum Firmicutes (42%), Proteobacteria (29%), and Planctomycetes (14%), respectively. The most dominant genus in vegetative phase was *Candidatus koribacter*, while at the reproductive phase was *Aquisphaera*. The analysis of macronutrients P, K, Mg, Ca, and micronutrients Fe, Mn, Cu, Na, Zn showed a reduction in the nutrient concentration at the reproductive phase compared to the vegetative phase excluding S element. Furthermore, the concentration of S element affects the abundance of bacteria in the reproductive phase, while the other elements of nutrients affect the vegetative phase. Based on these results, further strategies can be formulated to increase yield and guide breeding of mandoti aromatic rice.

1. Introduction

Aromatic rice has a fairly high economic value, which is in demand by consumers due to its delicious and fluffier taste (Syamsia, 2016). This includes Pulut Mandoti Emas (PME) that had a strong pandan fragrance, where 1 L can contribute aroma to 40 L of ordinary rice (Karim et al., 2020). PME rice is endemic to Enrekang Regency, Salukanan Village, Baraka District, therefore, it will not produce similar quality when cultivated outside the village. Although it has a red color when produced, it does not produce a strong distinctive aroma (Latif et al., 2020). The selling price from farmers is Rp. 55,000.00 per liter in 2019 and increased to Rp. 70,000.00 in 2020. This high selling price is commensurate with the long harvest period, which is six months after planting, with harvest time only once a year. Furthermore, not all rice fields in Salukanan Village are planted with PME rice and the planted area is only about 250 ha of the existing 312 ha of rice fields (Karim et al., 2020). This limited the production and was insufficient to meet the increasing

market demand within Enrekang Regency and others.

The discovery and application of microbial inoculants to enhance plant growth and health is a promising alternative for addressing society's aspirations for more food and a more nutritious and sustainable food supply (Doni et al., 2022). The dynamics of soil microbes significantly determine the potential for plant productivity and stimulate growth regulation and nitrogen fixation (Ahmad et al., 2008; Kumar and Verma, 2019). *Pseudomonas otitidis*, *Enterobacter* sp., *Chryseobacterium* sp. have reported promote Basmati rice growth (Jasrotia et al., 2021).

Several studies have been conducted on the importance of rhizosphere symbiotic bacteria in the production of secondary metabolites by plants. The production of 2-acetyl pyrroline (2AP) in basmati aromatic rice, the compound responsible for the aroma enhanced by rhizobacteria (Deshmukh et al., 2016). It was also stated that a consortium of rhizosphere bacteria mediates the increase in the production of 2AP compounds in Basmati aromatic rice (Dhondge et al., 2021).

It is necessary to explore bacterial species associated with PME rice

* Corresponding author. Microbiology Laboratory, Department of Biology, Faculty of Science and Technology, UIN Alauddin Makassar, 92118, Gowa, Indonesia.
E-mail address: Ekasukmawaty@uin-alauddin.ac.id (Eka Sukmawaty).

as the initial stage to increase its products nationally and internationally. Therefore, this study was conducted to investigate the effects of soil properties, the soil macro and micronutrient contents, and the relative abundance of bacterial community in the rhizosphere between vegetative and reproductive phase.

2. Materials and methods

2.1. Sampling and measurement of environmental parameters

Rhizosphere soil samples were collected from the rice fields of PME in Salukanan Village, Baraka District, Enrekang Regency, with coordinates 3°22'59.1"S, 119°53'20.0"E. The soil was removed carefully from the rice plant and the soil attached to the roots was taken as a sample. Meanwhile, the excavations were carried out around the roots at a depth of approximately 20 cm, where 50 g of soil samples were taken, put in sterile plastic bags, and labeled according to the time and sampling location. The sampling time was carried out in two phases of rice growth, namely the vegetative and the reproductive phases. The environmental parameters measured were pH, humidity, and temperature of the rhizosphere soil.

2.2. Analysis of macro and micro nutrients

This stage aims to determine the content of macro and micro-nutrients in the rhizosphere of PME rice. It was started by precisely weighing 0.5 g of fine soil into a digest tube. Subsequently, approximately 5 ml of nitric acid pa and 0.5 ml of perchloric acid pa were added to the tube and allowed to stand overnight, while the tube was heated at 100 °C for 1.5 h the day after. The temperature was increased to 130 °C for 1 h and later to 150 °C for 2 h 30 min until the yellow steam fades out, and in case there is still yellow steam remaining, the heating time is further increased. When the yellow vapor have disappeared, the temperature was raised to 170 °C for 1 h and increased to 200 °C for another 1 h until white steam is formed. The destruction is completed with the formation of a white precipitate or commonly referred to as a clear solution of approximately 0.5 ml.

The clear extract was used for the measurement of macro elements, namely P, K, Ca, Mg, Na, S, and the microelements Fe, Al, Mn, Cu, Zn, and B.

2.2.1. Measurement of P

A total of 1 ml of soil extract and each standard series P were pipetted into a chemical tube, 9 ml of deionized water was added and shaken. Subsequently, 1 ml of aqueous extract and standard series of P was pipetted into a test tube, and 10 ml of dye reagent P was also added. The solution was shaken using a shaker tube until homogeneous and left for 30 min. The Phosphorus (P) in the solution was measured with a spectrophotometer at a wavelength of 889 nm.

2.2.2. Measurement of K, Ca, Mg, and Na

A total of 1 ml of extract and standard series were pipetted into a chemical tube and 9 ml of 0.25% solution, where Mg, the extract was first diluted with zero standard 5 times before pipetting. The solution was shaken using a tube shaker until homogeneous, while the K, Ca, Mg, and Na in the extract were measured by AAS using a standard series as a comparison.

2.2.3. Measurement of S

In this study, 1 ml of extract and standard series 5 each into a chemical tube. Subsequently, 7 ml of mixed acid and 2.5 ml of BaCl₂-tween solution were added to each tube and were shaken with a tube shaker until homogeneous. The solution was allowed to stand for 30 min and measured with a spectrophotometer at a wavelength of 494 nm.

2.2.4. Measurement of Fe, Al, Mn, Cu, and Zn

Fe, Al, Mn, Cu, and Zn were measured directly from the extract using AAS, with their respective standard series as a comparison. Al was used a gaseous mixture of NO₂-acetylene flame, while others were used an air-acetylene mixture flame.

2.3. DNA extraction and metagenome sequencing

Genomic DNA (GDNA) from each sample was isolated using Quick-DNA Fungal/Bacterial Kits from Zymo Research. The concentration of DNA was determined using NanoDrop spectrophotometers and Qubit fluorometer, while GDNA was amplified using Primer 16S 27F-1429R. The PCR products were observed by 1% agarose gel electrophoresis. Library preparations were carried out using Kits from Oxford Nanopore Technology (Oxford Science Park, UK) and the GridION sequencing was operated by MinKNOW version 20.06.9. Furthermore, the base-calling was using Guppy version 4.0.11 with high accurate mode. The quality of FASTQ files was visualized using NanoPlot and the filtered reads were classified using Centrifuge classifier. The bacteria and Archaea index was downloaded from the centrifuge website (<https://ccb.jhu.edu/software/centrifuge>). Downstream analysis and visualizations were carried out using Pavian (<https://github.com/fbreitwieser/pavian>) and Krona Tools (<https://github.com/marbl/Krona>).

2.4. Statistical analysis

The data of environmental soil factors including temperature, pH, and moisture between two growth phases of vegetative and reproductive were analyzed using Independent Sample *t*-test to establish the significance. The data were conducted using IBM version 25.0 at $p < 0.05$ and presented as Mean \pm SD ($n = 3$).

3. Results

3.1. Environmental parameters

The microbial community in the rhizosphere is influenced by several biotic and abiotic factors such as temperature, pH, salinity, as well as, soil moisture, organic compounds, type, and structure, respectively (Fierer, 2017). Based on the results of environmental observations in the two growth phases of PME rice, it was discovered that the soil pH changed from acid (pH 4.6) in the vegetative phase to normal (pH 7) in the reproductive phase (Table 1).

3.2. Macro and micro-nutrients

Another factor that affects soil pH is chemical cations in the soil. This is because cation exchange capacity plays a significant role as a buffer for changes in soil pH by controlling the supply of exchangeable cations such as Ca²⁺, Mg²⁺, K⁺, N⁺, Al⁺, and Fe³⁺ (Fang et al., 2017). The supply of these cations can be obtained from the nutrients in the soil. The measurement results of the macro and micro soil nutrients (Table 2) in the PME rice fields showed that the nutrients decreased from the vegetative to the reproductive phase.

Table 1

Changes in soil temperature, pH, and moisture between vegetative and reproductive phases in PME rice fields.

Phase growth	Temperature (°C)	pH	Moisture (%)
Vegetative	25.33 \pm 0.58**	4.6 \pm 0.28**	67.77 \pm 1.66**
Reproductive	35.57 \pm 0.15**	7.0 \pm 0.00**	57.27 \pm 1.50**

Note. Abbreviation (**) $P < 0.01$, Independent Sample *t*-test. Data show Mean \pm SD.

Table 2
Analysis of macro and micro nutrient levels in PME rhizosphere.

Growth Phase	Nutrients									
	P2O5, %	K2O, %	Fe, ppm	Mn, ppm	Mn, ppm	Cu, ppm	Ca, ppm	Mg, ppm	S, %	
Vegetative	0.06	0.04	27622	192	34	30.4	1368	1789	0.04	
Reproductive	0.02	0.03	22189	185	30.18	28.64	1261	1508	0.14	

3.3. Metagenome analysis

The results of metagenome readings showed that there are 40 bacterial phyla, which consisted of 73 classes and 338 families in the vegetative phase of PME rice. Meanwhile, in the reproductive phase, it was shown that there are 42 bacterial phyla, consisting of 77 classes and 357 families. The phyla with the highest relative abundance in the vegetative phase were *Acidobacteria* (35%), *Proteobacteria* (30%), and *Firmicutes* (12%), while the highest in the reproductive phase was shown by the phylum *Firmicutes* (42%), *Proteobacteria* (29%), and *Planctomycetes* (14%) of all bacteria (Fig. 1).

At the genus level, the dominant bacteria in both growth phases are *Acidisarcinia*, *Candidatus koribacter*, *Terriglobus*, *Candidatus solibacter*, *Bacillus*, *Clostridium*, *Gemmata*, *Aquisphaera*, *Rhodoplanes*, *Penibacillus*, *Pseudobacillus*, *Pseudoclostridium*, *Desulfosporosinus* and *Methylocystis* but with different relative abundances. At level class, *Acidobacteria* was the abundant class in the vegetative phase, while *Clostridia* was the highest in the reproductive phase. However, the *Nitrosopira* class was only discovered in the vegetative phase, while the *Negativicutes* class was observed only in the reproductive phase. (Fig. 1A). At the family level, the most abundant bacteria in the vegetative and reproductive phases were *Acidobacteriaceae* and *Colistrideaceae*, respectively (Fig. 1B). Furthermore, *Candidatus koribacter* was the highest abundance genus in the vegetative phase, while at the reproductive phase *Aquisphaera* was the most abundant genus (Fig. 1C).

3.4. Soil physicochemical properties

The CCA analysis was carried out to determine the effect of soil physicochemical on the bacterial community in the rhizosphere of PME rice. The results showed that temperature, pH, and S are factors that affect the relative abundance of *Firmicutes*, *Actinobacteria*, *Chloroflexi*, and *Planctomycetes* in the vegetative phase. Meanwhile, humidity and other nutrients affect the relative abundance of *Cyanobacteria*, *Proteobacteria*, *Verruimicrobia*, *Bacteriodes*, *Nitrospirae*, and *Acidobacteria* in the reproductive phase (Fig. 2).

4. Discussion

In the present study showed that the abundance of rhizosphere bacteria were affected by environmental factors and soil physicochemical properties. The environmental parameters were observed are pH, temperature and humidity. pH of soil increased in reproductive phase. The increased pH was likely influenced by the condition of rice field irrigation. During the vegetative phase, the rice fields are flooded with water, hence, the organic compounds are fermented in the fields, which makes the soil pH acidic. Furthermore, in flooded land conditions, organic material is decomposed under anaerobic conditions by soil microbial activity, supplying electrons and protons, leading to an increase in soil pH in the reproductive phase through decarboxylation of anions of organic compounds (Ding et al., 2019). The increase in soil pH also occurs in line with the increase in temperature and decrease in humidity from the vegetative to the reproductive phase. This was in accordance with Ghazali et al. (2020) who stated that the pH is negatively correlated with moisture and positively correlated with soil temperature.

The concentration of macro and micro-nutrients was observed in this study for evaluating their effect to PME rice. Nutrients affect rice growth

and physiology directly and indirectly (Mengistu et al., 2018; Kumar et al., 2021). The analysis of nutrients from the vegetative and reproductive phases of PME rice showed that all observed nutrients decreased in concentration due to their use by rice plants for growth. However, sulfur increased in concentration from 0.04% in the vegetative phase to 0.14% in the reproductive phase due to the nutrient antagonistic interactions. Macro and micronutrients can influence each other, where some micronutrients affect the absorption of macronutrients and vice versa. The increase in S nutrients affects the nutrients of Mn, Mo, Zn, and Fe (Shrestha et al., 2020).

Rhizosphere bacteria affected the increase in plant growth, however, the mechanism has not been fully established. These NGS approaches such as metagenomics are necessary to assist in their exploration (Babalola et al., 2020). Metagenomics targets the sequence sequenced and identified based on similarities in the gene sequences available in the database (Ivanova et al., 2016). These results are in line with the previous studies, which stated that the Basmati aromatic rice rhizosphere bacterial community from the phylum *Proteobacteria*, *Acidobacteria*, *Actinobacteria*, *Chloroflexi*, and *Firmicutes* were the most abundant bacteria (Ivanova et al., 2016). According to Pittol et al. (2018), the dominant phyla in the flooded rice fields in the vegetative phase were the *Proteobacteria* phylum from the *Alphaproteobacteria*, and *Betaproteobacteria* classes. Ding et al. (2019) also showed that the bacterial community in the rice rhizosphere was mostly occupied by *Proteobacteria*, mainly Alpha-, Beta- and Deltaproteobacteria classes, such as *Acidobacteria*, *Actinobacteria*, and *Chloroflexi*. Meanwhile, Lopes et al. (2014) discovered that the dominant phyla in alpha rice farming were *Acidobacteria*, *Proteobacteria*, *Chloroflexi*, *Actinobacteria*, and *Bacteroidetes*. *Proteobacteria* and *Acidobacteria* play an important role in the C cycle in soil, due to their ability to degrade cellulose and lignin. *Acidobacteria* provides macro and micronutrients, produce exopolysaccharides, express a variety of active transporters, and function in soil acidity (Kielak et al., 2016; Anandan et al., 2016). *Firmicutes*, including the genus *Bacillus*, are the most studied group of bacteria, which are known to promote plant growth. According to Kim et al. (2021), *Firmicutes* function as decomposers that converse organic compounds. Meanwhile, *Actinobacteria* are associated with disease suppression by producing antibiotics. They also possess the ability to increase root nodulation and promote plant growth by providing nutrients and forming humus (Lagos et al., 2015).

At the genus level, *Candidatus koribacter* was the highest abundance genus in the vegetative phase, while at the reproductive phase *Aquisphaera* was the most abundant genus. *Candidatus koribacter* have found as PGPR and known as suppressor to *Fusarium oxysporum* pathogen (Nisrina et al., 2021). *Candidatus koribacter* play role in resistance to fluctuating temperature and nutrient deficient conditions due to carbohydrate enzyme family production (Liu et al., 2021). The presence of these bacteria may play a role in the adaptation of PME in early growth phase.

The structure of the microbial community, particularly the bacterial in the PME rice rhizosphere is diverse and dynamic. This is influenced by soil and plant-related conditions such as geographic location, soil type, and rice genotype. Meanwhile, changes in these conditions can alter the soil environment for microbial growth and proliferation, namely soil pH and root exudation patterns. This will determine the various structure of the rhizosphere microbial community. Agricultural management, which involves cultivation, drainage practices, and growth also influences rice




Fig. 1. Relative abundance of (A) class, (B) family, (C) genus bacteria in PME rice.




Fig. 2. Effect of Soil Physicochemical properties on The Rhizosphere Bacterial Community of PME Rice.

rhizosphere microbiome (Ding et al., 2019). Plant type and soil characteristics can be impactful to the composition of endophytic bacteria in the roots. This is because soil properties affect the structure of microbial communities directly by providing specific habitats while specifying different groups of microorganisms, and indirectly by influencing plant root physiology. However, many reports indicated that plant species and cultivars are also important factors for establishing the root microbiome (Cordero et al., 2020).

5. Conclusion

The results showed that the rhizosphere bacterial community of PME rice in the vegetative and reproductive phases tend to be similar, only differs from the relative abundance in each growth phase. In the vegetative phase, the phyla with the highest relative abundance were *Acidobacteria* (35%), *Proteobacteria* (30%), and *Firmicutes* (12%), while in the reproductive phase the phylum *Firmicutes* (42%), *Proteobacteria* (29%) and *Planctomycetes* (14%) were the most abundant. Furthermore, the bacterial phylum that occupies the rhizosphere habitat of PME rice plays a role in plant growth as PGPR. The analysis of macro and micro-nutrients showed a general decrease from the vegetative to the reproductive phase, except for S that increased in concentration in the reproductive phase. Based on the CCA results, it was discovered that S, temperature, and pH affect the bacterial community in the vegetative phase, while humidity and other nutrients affect the reproductive phase of PME rice. These results provide initial information for further strategies to increase yield and guide breeding of PME rice.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgment

The authors are grateful to the Salukanan village government for the permission and Mr. Middin who allowed his rice fields to be used as sampling sites during this study.

References

- Ahmad, F., Ahmad, I., Khan, M.S., 2008. Screening of free-living rhizospheric bacteria for their multiple plant growth promoting activities. *Microbiol. Res.* 163 (2), 173–181. <https://doi.org/10.1016/j.micres.2006.04.001>.
- Anandan, R., Dharumadurai, D., Manogaran, G.P., 2016. An introduction to actinobacteria. In: *Actinobacteria-Basics and Biotechnological Applications*. Intechopen.
- Babalola, O.O., Fadiji, A.E., Enagbonma, B.J., Alori, E.T., Ayilara, M.S., Ayangbenro, A. S., 2020. The nexus between plant and plant microbiome: revelation of the networking strategies. *Front. Microbiol.* 11, 2128. <https://doi.org/10.3389/fmicb.2020.548037>.
- Cordero, J., de Freitas, J.R., Germida, J.J., 2020. Bacterial microbiome associated with the rhizosphere and root interior of crops in Saskatchewan, Canada. *Can. J. Microbiol.* 66 (1), 71–85. <https://doi.org/10.1139/cjm-2019-0330>.
- Deshmukh, Y., Khare, P., Patra, D., 2016. Rhizobacteria elevate principal basmati aroma compound accumulation in rice variety. *Rhizosphere* 1, 53–57. <https://doi.org/10.1016/j.rhisph.2016.07.001>. July.
- Dhondge, H.V., Pable, A.A., Barvkar, V.T., Dastager, S.G., Nadaf, A.B., 2021. Rhizobacterial consortium mediated aroma and yield enhancement in basmati and non-basmati rice (*Oryza sativa* L.). *J. Biotechnol.* 328, 47–58. <https://doi.org/10.1016/j.jbiotec.2021.01.012>.
- Ding, C., Du, S., Ma, Y., Li, X., Zhang, T., Wang, X., 2019. Changes in the pH of paddy soils after flooding and drainage: modeling and validation. *Geoderma* 337, 511–513. <https://doi.org/10.1016/j.geoderma.2018.10.012>. July 2018.
- Doni, F., Suhaimi, N.S., Mispan, M.S., Fathurrahman, F., Marzuki, B.M., Kusmoro, J., Uphoff, N., 2022. Microbial contributions for rice production: from conventional crop management to the use of 'omics' technologies. In: *International Journal of Molecular Sciences*. <https://doi.org/10.3390/ijms23020737> (Vol. 23, Issue 2), 3088–3097. <https://doi.org/10.1002/2017JG003968>.
- Fang, K., Kou, D., Wang, G., Chen, L., Ding, J., Li, F., Yang, G., Qin, S., Liu, L., Zhang, Q., Yang, Y., 2017. Decreased soil cation exchange capacity across northern China's grasslands over the last three decades. *J. Geophys. Res. Biogeosci.* 122 (11), 3088–3097. <https://doi.org/10.1002/2017JG003968>.
- Fierer, N., 2017. Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat. Rev. Microbiol.* 15 (10), 579–590. <https://doi.org/10.1038/nrmicro.2017.87>.
- Ghazali, M.F., Wikantika, K., Harto, A.B., Kondoh, A., 2020. Generating soil salinity, soil moisture, soil pH from satellite imagery and its analysis. *Inf. Process. Agric.* 7 (2), 294–306. <https://doi.org/10.1016/j.inpa.2019.08.003>.
- Ivanova, N., Gugleva, V., Dobрева, M., Pehlivanov, I., Stefanov, S., Andonova, V., 2016. We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists TOP 1%. Intech 13 i(tourism).
- Jasrotia, S., Salgotra, R.K., Samnotra, R.K., 2021. Identification of basmati rice (*Oryza sativa* L.) rhizobacteria and their effect on plant growth traits for sustainable development in agriculture. *Proc. Indian Natl. Sci. Acad.* 87 (3), 469–486. <https://doi.org/10.1007/s43538-021-00033-6>.
- Karim, A., Amrullah, Junaidin, Syukur, A., 2020. Beras Lokal Pulu 'Mandoti Penopang Ekonomi Dan Ketahanan Pangan Masyarakat Desa Salukanan di Tengah Crisis Pandemi Global 'Covid - 19.' May.
- Kielak, A.M., Barreto, C.C., Kowalchuk, G.A., van Veen, J.A., Kuramae, E.E., 2016. The ecology of *Acidobacteria*: moving beyond genes and genomes. *Front. Microbiol.* 7, 744.

- Kim, S., Samaddar, S., Chatterjee, P., Choudhury, A.R., Choi, J., Choi, J., Sa, T., 2021. Structural and functional shift in soil bacterial community in response to long-term compost amendment in paddy field. *Appl. Sci.* 11 (5), 1–10. <https://doi.org/10.3390/app11052183>.
- Kumar, A., Verma, J.P., 2019. *The Role of Microbes to Improve Crop Productivity and Soil Health* (Issue January 2019). Springer, Singapore. https://doi.org/10.1007/978-981-13-0149-0_14.
- Kumar, S., Kumar, S., Mohapatra, T., 2021. Interaction between macro- and micro-nutrients in plants. In: *Frontiers in Plant Science*, vol. 12. <https://www.frontiersin.org/article/10.3389/fpls.2021.665583>.
- Lagos, M.L., Maruyama, F., Nannipieri, P., Mora, M.L., Ogram, A., Jorquera, M.A., 2015. Current overview on the study of bacteria in the rhizosphere by modern molecular techniques: a mini-review. *J. Soil Sci. Plant Nutr.* 15 (2), 504–523. <https://doi.org/10.4067/s0718-95162015005000042>.
- Latif, J., Busthanul, N., Amrullah, A., 2020. Motivasi Petani dalam Budidaya Padi Pulu mandoti. *Jurnal Sosial Ekonomi Pertanian* 16 (1), 49. <https://doi.org/10.20956/jsep.v16i1.8353>.
- Liu, T.C., Peng, H.M., Wollney, S., Shen, C.H., 2021. Rhizosphere microbiome regulates the growth of mustard under organic greenhouse cultivation. *Agriculture (Switzerland)* 11 (10), 1–10. <https://doi.org/10.3390/agriculture11100987>.
- Lopes, A.R., Manaia, C.M., Nunes, O.C., 2014. Bacterial community variations in an alfalfa-rice rotation system revealed by 16S rRNA gene 454-pyrosequencing. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Ecol.* 87 (3), 650–663. <https://doi.org/10.1111/1574-6941.12253>.
- Mengistu, T., 2018. Role of macro nutrient for rice production: review articles. *Int. J. Res. Agron.* 1 (2), 60–66.
- Nisrina, L., Effendi, Y., Pancoro, A., 2021. Revealing the role of Plant Growth Promoting Rhizobacteria in suppressive soils against *Fusarium oxysporum* f.sp. *cubense* based on metagenomic analysis. *Heliyon* 7 (8), e07636. <https://doi.org/10.1016/j.heliyon.2021.e07636>.
- Pittol, M., Scully, E., Miller, D., Durso, L., Mariana Fiuza, L., Valiati, V.H., 2018. Bacterial community of the rice floodwater using cultivation-independent approaches. *Int. J. Microbiol.* 2018 <https://doi.org/10.1155/2018/6280484>.
- Shrestha, J., Kandel, M., Subedi, S., Shah, K.K., 2020. Role of nutrients in rice (*Oryza sativa* L.): a review. *Agrica* 9 (1), 53. <https://doi.org/10.5958/2394-448x.2020.00008.5>.
- Syamsia, 2016. Peningkatan Pertumbuhan vegetatif Padi Aromatik lokal Enrekang menggunakan cendawan Endofit increased of vegetative growth of aromatic rice local Enrekang using endophyte fungi. *Jurnal Galung Tropika* 5 (2), 80–85.