

Description of screen time and dry eye syndrome in medical students of UIN Alauddin Makassar

by Ulfah Rimayanti

Submission date: 24-May-2023 11:22AM (UTC+0700)

Submission ID: 2100582658

File name: 3710-BMJ-Final_Edit_FIX.docx (114.2K)

Word count: 3351

Character count: 18243

Description of screen time and dry eye syndrome in medical students of UIN Alauddin Makassar

Ulfah Rimayanti^{1,2*}, Gita Yustika¹, Sri Rezki Wahdania Jamaluddin¹

³
¹School of Medicine, Universitas Islam Negeri Alauddin Makassar, Indonesia
²Haji Government Hospital, Makassar, Indonesia

³
*Correspondence: Ulfah Rimayanti. School of Medicine, Universitas Islam Negeri Alauddin Makassar, Indonesia. Email: rimayantiu@gmail.com

Abstract:

Background: During pandemic, the amount of screen time in individuals is tremendously expanding. Excessive screen time may result in eye blinks reduction, tears evaporation, and eventually dry eye syndrome (DES). This study aims to show the descriptions of the duration of screen time and dry eye symptoms in medical students during the pandemic.

Methods: This study uses a descriptive and cross-sectional methods. The subjects were students of The School of Medicine at UIN Alauddin Makassar, who were undergoing online learning. The data were collected through a questionnaire containing subjects' activities using gadgets per day and the Ocular Surface Disease Index (OSDI) to assess dry eye symptoms. Ophthalmological examinations for DES (blink rate, tear meniscus height, tear break-up time, Schirmer test) were also performed on the study subjects.

Results: Most of the subjects are female and 22 years old. The average screen time of the subjects was 17.7 hours per day. Two subjects (4.55%) had mild dry eye symptoms, 14 subjects (31.82%) had moderate dry eye symptoms, and 28 subjects (63.64%), had severe dry eye symptoms. This study also found that the average blink rate of the subjects was 11.36 ± 7.45 times per minute, and the average TBUT value was 10.58 ± 2.69 seconds. Schirmer 1 examination showed a value with a mean of 23.67 ± 12.76 mm, and Schirmer 2 examination showed a value of 20.21 ± 11.64 mm.

Conclusion: The average screen time of the subjects was 17.7 hours. According to the OSDI index, most subjects (63.64%) had severe dry eye symptoms and low blink rate (82%).

Keywords: dry eye, pandemic, screen time

Introduction

Recently, the number of computer users and display devices with a monitor (screen) is increasing. In Indonesia there are 143 million internet users, making Indonesia one of the top five internet users in the world. Most of the internet users are young people and teenagers, even though adults user has also increased.¹ Excessive screen time may result in imperfect blinking, evaporation of tears, and ultimately leading to Dry Eye Syndrome (DES).² Substantially, DES could affect visual acuity and quality of life, as the symptoms often disturb daily activities, such as reading, writing, or

working in front of the screen.³ Undiagnosed dry eye may cause unstable conditions on the ocular surface, leading to corneal damage, ulcer, and cicatrix due to corneal irregularity.⁴

Dry eye is more often found in the elderly, especially in the 4th decade.³⁻⁶ This fact influences the limited number of dry eye research in teenagers and young adults. Moreover, the COVID-19 pandemic has increased screen time among students as learning is generally shifted to online mode and the use of gadgets as learning tools is unavoidable.^{7,8} This study is to show the description of screen time and DES events in students, especially medical students of UIN Alauddin, Makassar.

Material and methods

This study uses descriptive and cross-sectional methods. The study was conducted in the School of Medicine at UIN Alauddin Makassar and RSUD Haji Makassar from December 2020 to April 2021. The subjects were students of The School of Medicine at UIN Alauddin Makassar, aged 17-24 years old, who were undergoing online learning. The inclusion criteria included those who were willing to fill the questionnaires and take the examinations. Exclusion criteria were incomplete questionnaire filling, history of eye surgery, the use of soft contact lenses, abnormal palpebral anatomy, the use of antihistamine eye drops medication, and history of systemic disease.

The data collection was undertaken using online questionnaires and ophthalmological examinations. Written and signed informed consent from the subject was obtained before the data collection was performed. The questionnaires consisted of the a daily log on the research and Ocular Surface Disease Index (OSDI) questionnaire filled in and submitted by the medical students via a Google form. The daily log included the personal identity, such as age and gender, as well as screen time, use of contact lenses, use of fans or air conditioners (AC), non-screen duration per day. The OSDI questionnaire was scored on a range of 1-100 points. The OSDI score was classified as normal (0-12), mild (13-22), moderate (23-32), severe (33-100) dry eye symptoms.

Ophthalmological examinations for DES (blink rate, tear break-up time, Schirmer tests 1 and 2) were performed on the study subjects at RSUD Haji Makassar. The blink rate measurement was done observationally during 1 minute usage of the smartphone. The normal blink rate was 12-18 times/minute. The examination of the anterior segment of the eyeball was performed using a slit lamp biomicroscope. The subjects had Tear Break-up Time (TBUT) examination by applying a fluorescein strip combined with 1 drop of balanced salt solution on the inferior part of conjunctival fornix of the eye. The subjects were asked to blink 3 times then look straight ahead without blinking. The time between the last blink and the onset of dry spot on the ocular surface was identified using a cobalt blue filter and measured using a stopwatch. Normal TBUT was >10 sec, 6-10 sec was classified as having mild dry eye, and ≤5 sec was identified as having severe dry eye.

The Schirmer test was performed by placing the Schirmer strip paper on the inferior conjunctival sac in the temporal third of the inferior palpebral of both eyes. The subjects were asked to keep their eyes to remain open then asked to blink. After 5 min, the strip was taken off, and the length of the wet strip paper was measured. The examination result was considered normal when the wet strip was ≥15 mm, mild dry eye when it was 9-14 mm, medium dry eye if it was 4-8 mm, and heavy dry eye when it was <4 mm long. Each examination was repeated 3 times and the average

value was calculated. All the data were analyzed using SPSS version 25 descriptively and presented in the form of a frequency distribution table.

Results

A total of 44 students became the subjects of this study. The questionnaires contain the subject characteristics, screen time, and the OSDI index. The collected data were presented in tables. Table 1 shows the characteristics of the study subjects, namely age, gender, screen times, use of contact lenses, use of fans or air conditioners (AC), and non-screen duration. Most of the study subjects were 22 years old (27.27%) and female (65.91%). The average screen time for the online class was about 8.14 hours per day and the average screen time for other online activities was 9.57 hours per day. There were nine subjects (9.09%) who wore contact lenses. The average use of the fan/AC was 12.55 hours per day. The average non-screen duration was 6.30 hours per day.

Table 1 Characteristic Distributions of Research Subjects

Subjects Characteristics	Results
Age	
17 years	1 (2.27%)
18 years	5 (11.36%)
19 years	7 (15.91%)
20 years	2 (4.55%)
21 years	9 (20.45%)
22 years	12 (27.27%)
23 years	4 (9.09%)
24 years	4 (9.09%)
Sex	
Male	15 (34.09%)
Female	29 (65.91%)
Screen Time	
Online class	8.1 hours
Other than online classes	9.6 hours
Total screen time per day	17.7 hours
Use of Contact Lenses	4 (9.09%)
Use of Fan/AC	12.55 hours
Non-Screen Time	6.30 hours

Table 2 shows the distribution of dry eye symptoms using the OSDI index. All study subjects had dry eye symptoms. It was found that most study subjects (n=28 (63.64%)) suffered from severe dry eye symptoms. Study subjects suffering from moderate dry eye symptoms were 14 subjects (31.82%). Study subjects who suffered from mild dry eye symptoms were 2 persons (4.55%).

5 **Table 2.** The distribution of Dry Eye Symptoms Using the OSDI index

Dry Eye Symptoms	Results
Normal	0 (0.00%)
Mild	2 (4.55%)
Moderate	14 (31.82%)
Severe	28 (63.64%)
Total	44 (100.00%)

Table 3 shows the distribution of dry eye symptoms based on gender. In the category of mild dry eye, all two subjects (100%) were females. In the moderate dry eye category, there were 8 females (57.14%) and 6 males (42.86%). In the category of severe dry eye, there were 19 females (67.86%) and 9 males (32.14%).

Table 3. Distribution of Dry Eye Symptoms by Sex

Dry Eye Symptoms	Male	Female	Total
Normal	0 (0.00%)	0 (0.00%)	0 (0.00%)
Mild Dry Eye	0 (0.00%)	2 (100%)	2 (100%)
Moderate Dry Eye	6 (42.86%)	8 (57.14%)	14 (100%)
Severe Dry Eye	9 (32.14%)	19 (67.86%)	28 (100%)

Table 4 shows the data of dry eye signs examinations. The average blink rate was 11.36 ± 7.45 times per minute. From the TBUT examination, the average value was 10.58 ± 2.69 seconds. The results of the Schirmer 1 examination showed a mean value of 23.67 ± 12.76 mm. The mean of the results of the Schirmer 2 examination was 20.21 ± 11.64 mm.

Table 4. Distribution of DES Examination

Parameters	Mean \pm Standard Deviation
Blink Rate	11.36 ± 7.45
TBUT	10.58 ± 2.69
Schirmer 1	23.67 ± 12.76
Schirmer 2	20.21 ± 11.64

TBUT: Tear Break-Up Time

The data are expressed as the mean \pm standard deviation.

Discussion

Based on age, this study showed that most subjects were 22 years old. The incidence of dry eye increases with age due to decreased function of the lacrimal gland,⁹ but in this study, we cannot conclude that there is a relationship between age and the incidence of dry eye due to the adjacent population. In the younger age group, it was reported that the incidence of dry eye increased due to

the duration of the screen time which decreased the blink rate and increased the evaporation of the tear film.¹⁰ The results showed that the average duration of screen time was high, which was about 17.7 hours per day. A survey by the Indonesian Internet Service Providers Association (APJII) found that on average internet users in Indonesia (35.3%) access the internet for 1 hour per day, around 9% or 20 million subjects use the internet excessively (3% use 7-9 hours, and 6% use >9 hours).¹¹ Based on the age of internet users, most users in Indonesia are 18-25 years old, which is equivalent to almost half of the total internet users in Indonesia. In a study involving 259 smartphone users aged 19-25 years in Jakarta, 28% used the internet <6 hours, while 72% used the internet >6 hours per day.¹ Research on medical students in India shows that the majority of students (67%) spend 45 hours/week using the internet, the largest percentage is to send messages via email (36.6%), followed by academics (35.2%), pleasure (19.2%), and social networks (10%).¹² The higher screen time in this study was likely due to the online learning system during the pandemic, which encouraged students to spend much more time using computers or other gadgets.

This study found all stages of DES symptoms were more frequent in women than men. Several studies have reported the same results, where the presentation of women is greater than that of men. The prevalence of DES was greater in women (16.7%) than men (11.4%),¹³ other studies by Syawal et al. in Makassar also reported that dry eye was more common in women than men with a ratio of 2:1.¹⁴ A study in Sumatra reported a prevalence of 32.7% of DES in men and 22.8% in women.¹⁵ Genders can cause observable differences at both the molecular and physiological levels, in terms of the production of water layers in the eye, tear stability, the number of blinks, and even ocular immune function that may trigger DES.¹⁶ Hormonal factors, including levels of sex hormones as well as hormonal cycles in women, affect ocular structure and function. Lower levels of androgens in women have also been associated with DES.^{17,18} One of the risk factors of dry eye syndrome is female, it is thought to be due to the role of hormones in the stability of the tear film.¹⁹

Based on the OSDI score, most of our subjects experienced moderate and severe dry eye symptoms. Similar research involving 350 students reported that 27% had normal eye dryness, 16% had mild eye dryness, 28.5% had moderate eye dryness, and 28.5% had severe eye dryness.¹⁵ We found that the average values of TBUT, Schirmer 1, and Schirmer 2 test results were normal, while the average blink rate was lower than normal. Blink intervals vary from person to person, but healthy subjects blink roughly 20 times per minute on average.^{20,21} Phadatare and Messmer in 2015 put forward the theory that the occurrence of dry eye on smartphone use could be triggered by the activity of the eye staying long at the screen, thereby reducing the blink rate and increasing the length of exposure to the ocular surface. This may lead to instability of the tear film that causes dry eye complaints.^{22,23} Moon et al. made a similar statement that the reduced frequency of blinking rates during prolonged smartphone use could lead to faster evaporation of the tear film, which could result in dry eye.²⁴

The result showed that most of the subjects had a low blink rate. This will reduce the amount of moisture that the eyes can absorb from tears. Blinking helps to remove any debris from the eyeball and flushes out the fresh tears that have been produced. The retina will receive a clearer and

brighter image because of the coating of tears on the surface of the eyeballs. Blinking also provides oxygen and nutrients to the eye, ensuring that it remains healthy and comfortable.^{20,21}

Several studies indicated that the appearance of symptoms is adequate to diagnose dry eye as there is no definite single test to diagnose dry eye disease. However, symptoms alone are not sufficient for the diagnosis of dry eye because many ocular surface abnormalities and tear disorders show the same symptoms.²⁵ We performed both subjective and objective examinations for dry eye. We also obtained the average screen time of subjects in this study. However, there were several limitations, namely limited information about other factors that were likely to affect the results of the study, such as smoking habits and disease history. Another limitation is the absence of a control group, therefore the relationship between screen time duration and DES symptoms and signs cannot be evaluated.⁴

From the results of the study, it can be concluded that the average duration of screen time in medical students of UIN Alauddin Makassar is 17.7 hours per day and the subjective examination of DES with the OSDI questionnaire shows that most medical students of UIN Alauddin Makassar experience severe dry eye symptoms and a low blink rate.

Conclusion

This study showed that the average screen time of the subjects was 17.7 hours. Most subjects (63.64%) had severe dry eye symptoms and low blink rate (82%). The ophthalmological examinations (TBUT, Schirmer 1 and 2 tests) revealed normal results and a low blink rate. It is recommended that future studies can include a control group to be able to show the link between screen time duration and DES symptoms and signs.

Acknowledgments

This study was supported by the fund of The School of Medicine, Universitas Islam Negeri Alauddin Makassar. English Language editing was done by Nurhira Abdul Kadir, MD from the University of Wollongong, New South Wales, Australia.

Conflict of Interest

All authors declared that there is no conflict of interest regarding this article.

Ethical Consideration

The procedure follows the principles of the Declaration of Helsinki, and its use has been approved by the Ethics Committee of UIN Alauddin University Makassar, Indonesia (No.B.064a/KEPK/FKIK/II/2021).

Fundings

This study was supported by the fund of the School of Medicine, Universitas Islam Negeri Alauddin Makassar

Author Contribution

All authors contributed equally to this study

References

1. Prabowo H, Purwani Dewi M. Intercorrelation of insomnia, sleep duration, internet use duration and internet addiction on millennial in Jakarta. *Dinasti International Journal of Education Management and Social Science*. 2020;1(5). Available from: <https://dinastipub.org/DIJEMSS>
2. Akkaya S, Atakan T, Acikalin B, Aksoy S, Ozkurt Y. Effects of long-term computer use on eye dryness. *North Clin Istanbul*. 2018;5(4):319–22. Available from: <https://pubmed.ncbi.nlm.nih.gov/30859162>
3. Stapleton F, Alves M, Bunya VY, Jalbert I, Lekhanont K, Malet F, et al. TFOS DEWS II Epidemiology Report. *Ocul Surf*. 2017;15(3):334–65. Available from: <http://dx.doi.org/10.1016/j.jtos.2017.05.003>
4. Alves M, Dias AC, Rocha EM. Dry Eye in Childhood: Epidemiological and Clinical Aspects. *Ocul Surf*. 2008;6(1):44–51. Available from: [http://dx.doi.org/10.1016/s1542-0124\(12\)70104-0](http://dx.doi.org/10.1016/s1542-0124(12)70104-0)
5. Uchino M, Schaumberg DA, Dogru M, Uchino Y, Fukagawa K, Shimmura S, et al. Prevalence of Dry Eye Disease among Japanese Visual Display Terminal Users. *Ophthalmology*. 2008;115(11):1982–8. Available from: <http://dx.doi.org/10.1016/j.ophtha.2008.06.022>
6. Chidi-Egboka NC, Briggs NE, Jalbert I, Golebiowski B. The ocular surface in children: A review of current knowledge and meta-analysis of tear film stability and tear secretion in children. *Ocul Surf*. 2019;17(1):28–39. Available from: <http://dx.doi.org/10.1016/j.jtos.2018.09.006>
7. Manuaba IBAP, Darmayani IGAS, Sindhughosa DA, Yani MVW, Remitha NPSI, Sasmana IGAP, et al. Levels of anxiety and depression of students and responses to online learning methods during the COVID-19 pandemic. *Bali Medical Journal*. 2023;12(1):550–5.
8. Juliari IGAM, Sari Dewi R, Ratnasari NLMN, Tri Handayani A. The mucin 5AC level in medical faculty students with Computer Vision Syndrome (CVS). *Bali Medical Journal*. 2019;8(2):551–4.
9. Uchino M, Nishiwaki Y, Michikawa T, Shirakawa K, Kuwahara E, Yamada M, et al. Prevalence and Risk Factors of Dry Eye Disease in Japan: Koumi Study. *Ophthalmology*. 2011;118(12):2361–7. Available from: <http://dx.doi.org/10.1016/j.ophtha.2011.05.029>
10. Moon JH, Kim KW, Moon NJ. Smartphone use is a risk factor for pediatric dry eye disease according to region and age: a case control study. *BMC Ophthalmol*. 2016;16(1):188. Available from: <https://pubmed.ncbi.nlm.nih.gov/27788672>
11. Asosiasi Penyedia Jasa Internet Indonesia (APJII). Infografis penetrasi & perilaku pengguna internet Indonesia: Survey 2017 [Internet]. 2017. Available from:

- <https://www.kominfo.go.id/content/detail/12638/hasil-survei-penetrasi-dan-perilaku-pengguna-internet-2017-apjii/0/download2/5>
12. Singh B, Gupta R, Garg R, Resident S. Study of Medical Students and Internet Usage. *International Journal of Scientific and Research Publications*. 2013;3(5). Available from: www.ijsrp.org
 13. Moss SE. Prevalence of and Risk Factors for Dry Eye Syndrome. *Archives of Ophthalmology*. 2000;118(9):1264. Available from: <http://dx.doi.org/10.1001/archoph.118.9.1264>
 14. Pietersz EL, Sumual V, Rares L, Manado SR, Kesehatan BI, Fakultas M, et al. Penggunaan lensa kontak dan pengaruhnya terhadap dry eyes pada mahasiswa Fakultas Ekonomi Universitas Sam Ratulangi. Vol. 4, *Jurnal e-Clinic (eCl)*. 2016.
 15. Lee AJ, Lee J, Saw S-M, Gazzard G, Koh D, Widjaja D, et al. Prevalence and risk factors associated with dry eye symptoms: a population based study in Indonesia. *Br J Ophthalmol*. 2002;86(12):1347–51. Available from: <https://pubmed.ncbi.nlm.nih.gov/12446361>
 16. Matossian C, McDonald M, Donaldson KE, Nichols KK, Maciver S, Gupta PK. Dry eye disease: Consideration for women's health. Vol. 28, *Journal of Women's Health*. Mary Ann Liebert Inc.; 2019. p. 502–14.
 17. Ismael DS. Smartphone overuse and vision problems among university students. *ZANCO Journal of Pure and Applied Sciences (ZJPAS)*. 2022;34(3). Available from: <http://dx.doi.org/10.21271/zjpas.34.3.5>
 18. Susila NK, Mahayani NM, Triningrat AA, Widiana IG, Djelantik AA, Jayanegara WG. Blood superoxide dismutase (SOD) level has a negative correlation with dry eye (DE) degree. *Bali Medical Journal*. 2017;6(2):390.
 19. Truong S, Cole N, Stapleton F, Golebiowski B. Sex hormones and the dry eye. *Clin Exp Optom*. 2014;97(4):324–36. Available from: <http://dx.doi.org/10.1111/cxo.12147>
 20. Hirota M, Uozato H, Kawamorita T, Shibata Y, Yamamoto S. Effect of Incomplete Blinking on Tear Film Stability. *Optometry and Vision Science*. 2013;90(7):650–7. Available from: <http://dx.doi.org/10.1097/oxp.0b013e31829962ec>
 21. Argilés M, Cardona G, Pérez-Cabré E, Rodríguez M. Blink Rate and Incomplete Blinks in Six Different Controlled Hard-Copy and Electronic Reading Conditions. *Investigative Ophthalmology & Visual Science*. 2015;56(11):6679. Available from: <http://dx.doi.org/10.1167/iovs.15-16967>
 22. Phadatare SP, Momin M, Nighojkar P, Askarkar S, Singh KK. A Comprehensive Review on Dry Eye Disease: Diagnosis, Medical Management, Recent Developments, and Future Challenges. *Advances in Pharmaceutics*. 2015;2015:1–12. Available from: <http://dx.doi.org/10.1155/2015/704946>
 23. Messmer EM. The pathophysiology, diagnosis, and treatment of dry eye disease. *Dtsch Arztebl Int*. 2015;112(5):71–82. Available from: <https://pubmed.ncbi.nlm.nih.gov/25686388>
 24. Pazare P, Mujawar N, Singh P. Tears In Newborn Babies. *Int J Sci Res*. 2017;6(8).

25. Khanal S, Tomlinson A, Esakowitz L, Bhatt P, Jones D, Nabili S, et al. Changes in corneal sensitivity and tear physiology after phacoemulsification. *Ophthalmic and Physiological Optics*. 2008;28(2):127–34. Available from: <http://dx.doi.org/10.1111/j.1475-1313.2008.00539.x>

Description of screen time and dry eye syndrome in medical students of UIN Alauddin Makassar

ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

11%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Marliyanti Nurrahmah Akib, Suryana Rannu Pirade, Siti Rukiah Syawal, Muhammad Miftah Fauzan, Hasnah Eka, Arifin Seweng. "Association between prolonged use of smartphone and the incidence of dry eye among junior high school students", *Clinical Epidemiology and Global Health*, 2021
Publication 4%
- 2 belitungraya.org
Internet Source 1%
- 3 Batari Todja Umar, Ulfah Rimayanti, Halimah Pagarra, Budu, Nasrum Massi, Habibah Setyawati Muhiddin. "Novel point mutation and intronic mutations of *RB1* gene in retinoblastoma patients in Indonesia", *Medical Journal of Indonesia*, 2023
Publication 1%
- 4 balimedicaljournal.org
Internet Source 1%

5

Danai Tonkerdmongkol, Teera Poyomtip, Chotika Poolsanam, Akarapon Watcharapalakorn, Patarakorn Tawonkasiwattanakun. "Prevalence and associated factors for self-reported symptoms of dry eye among Thai school children during the COVID-19 outbreak", PLOS ONE, 2023

Publication

1 %

6

Alime Gunes, Seden Demirci, Hasan Rifat Koyuncuoglu, Levent Tok, Ozlem Tok. "Corneal and Tear Film Changes After Botulinum Toxin-A in Blepharospasm or Hemifacial Spasm", Cornea, 2015

Publication

1 %

7

Ning Wang, XiaoYun Zhuang, XiaoWei Zhong, Ju Zhang, GuangWei Li, Suxia Li. "Questionnaire Analysis on Incidence and Risk Factors of Dry Eye in Children From a Myopia Outpatient Clinic", Frontiers in Medicine, 2022

Publication

1 %

8

ro.uow.edu.au

Internet Source

1 %

9

www.visasolutions4u.com

Internet Source

1 %

10

oamjms.eu

Internet Source

1 %

11

worldnutrijournal.org

Internet Source

1 %

Exclude quotes *On*

Exclude matches *< 1%*

Exclude bibliography *On*