

Novel point mutation and intronic mutations of RB1 gene in retinoblastoma patients in Indonesia

by Batari Todja Umar Ulfah Rimayanti

Submission date: 05-Jun-2023 08:27AM (UTC+0700)

Submission ID: 2108960160

File name: 6544-Article_Text-50258-4-10-20230418.pdf (546.81K)

Word count: 4752

Character count: 24227

Novel point mutation and intronic mutations of *RB1* gene in retinoblastoma patients in Indonesia

Batari Todja Umar,¹ Ulfah Rimayanti,^{1,2} Halimah Pagarra,¹ Budu,^{1,3} Nasrum Massi,⁴ Habibah Setyawati Muhiddin¹



9

pISSN: 0853-1773 • eISSN: 2252-8083
<https://doi.org/10.13181/mji.oa.236544>
 Med J Indones. 2022;31:218–24

Received: October 22, 2022

Accepted: March 10, 2023

Published online: March 29, 2023

Authors' affiliations:

¹Department of Ophthalmology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia, ²School of Medicine, Universitas Islam Negeri

Alauddin Makassar, Makassar, Indonesia, ³Graduate School of Universitas Hasanuddin, Universitas Hasanuddin, Makassar, Indonesia, ⁴Department of Microbiology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia

¹³Department of Microbiology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia

Corresponding author:

Ulfah Rimayanti

Department of Ophthalmology, Faculty of Medicine, Universitas Hasanuddin, Jalan Perintis Kemerdekaan KM. 11, Tamalanrea, Makassar 90245, South Sulawesi, Indonesia

Tel/Fax: +62-411-580678

E-mail: rimayantiu@gmail.com

ABSTRACT

15

BACKGROUND Retinoblastoma (RB) is an inherited disorder caused by the *RB1* gene mutation in retinal cells or germline mutation. Identifying the specific mutation is crucial for prognosis, inheritance risk assessment, and treatment planning. This study aimed to identify the germline mutation in the *RB1* gene in patients with RB and their parents from the eastern part of Indonesia.

METHODS This observational analytic study recruited patients with RB and their parents between 2016 and 2018 at Dr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia. The normal control subjects were children from the outpatient clinic at the Department of Ophthalmology, Universitas Hasanuddin Hospital. Ophthalmic examinations and peripheral blood tests were performed in RB patients, their parents, and control subjects. Genomic DNA was isolated from blood leukocytes and amplified using conventional PCR. Hotspot exons 8, 10, 14, 17, and 22 were screened for mutations using the Sanger method.

RESULTS There were 21 patients with RB (16 unilateral and 5 bilateral) and 14 normal subjects. Of the 184 variations detected in RB patients, 164 were also found in normal subjects. 19 intronic mutations in introns 10, 16, 17, and 21, and 1 novel missense mutation in exon 17 were identified. Parental testing revealed 8 substitutions in exon 17 and 5 intronic mutations in introns 16 and 17 of the parents. None of the variations in exons were passed to their children.

CONCLUSIONS This study found a novel missense mutation in exon 17 of the *RB1* gene.

20

KEYWORDS germline mutation, *RB1* gene, retinoblastoma

Retinoblastoma (RB) is the most common eye neoplasm in childhood,¹ with approximately 50% of cases being heritable due to a mutation of the *RB1* gene that predisposes the development of retinal tumors. The *RB1* gene mutation occurs in retinal cells (somatic mutation) or gametes (germline mutation) in 98% of RB cases.² This gene comprises 27 exons spanning 178,143 kb genomic DNA at chromosome 13q14^{3,4} and encodes RB protein (pRB), which is a nuclear phosphoprotein and crucial for regulating cell

cycle progression.^{4,5} More than 1,900 mutations in the *RB1* gene have been reported globally,⁶ and identifying these mutations in RB patients is crucial for developing mutational analysis procedures and understanding the molecular mechanism underlying disease penetrance and expressivity.

Familial RB mutations exhibit variable expressivity and phenotypic variability.^{6–10} The *RB1* mutations associated with low penetrance include promoter mutations, missense mutations, and in-frame

7

Copyright © 2022 Authors. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are properly cited. For commercial use of this work, please see our terms at <https://mji.ui.ac.id/journal/index.php/mji/copyright>.

deletions/insertions.^{11,12} These mutations are autosomal dominant; thus, children whose parents have the RB1 gene mutation are at a higher risk of developing RB and other cancers such as osteosarcoma, small cell lung carcinoma, bladder tumor, and breast carcinoma.⁶ Identifying mutation type in patients with RB and their parents is essential for assessing short-term prognosis (intraocular tumor in the same eye or fellow eye), long-term prognosis (the risk of non-ocular malignancies), inheritance risk, and treatment strategy.^{13,14} Mutational screening of the RB1 gene can contribute to clinical management and provide accurate genetic counseling to patients and their families. Despite the importance of RB1 gene mutational analysis, there is a lack of data about germline RB1 mutations in RB patients in Indonesia. This study aimed to identify the RB1 gene mutations in patients with RB in the eastern part of Indonesia.

METHODS

Subjects

All subjects were recruited between 2016 and 2018 at Dr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia, which included patients diagnosed with RB and their parents. The diagnosis of RB was established through standard ophthalmological and histological examinations. Patients with a history of radiotherapy or other malignancies were excluded from the study. The control group was composed of children at the outpatient clinic at the Department of Ophthalmology, Universitas Hasanuddin Hospital, who had no abnormal ophthalmological findings except for mild refractive error and had neither history nor family history of neoplasm.

Sample collection and DNA extraction

A total of 5 ml peripheral blood samples were collected from patients and their parents into standard ethylenediaminetetraacetic acid (EDTA) tubes, and the DNA sample was stored at -20°C. The available parental samples were examined for the RB1 gene mutation at the same nucleotide position as their child who had a germline mutation. The blood samples were analyzed at Hasanuddin University Medical Research Center laboratory. DNA was isolated from peripheral blood leukocytes using gSYNC DNA Extraction Kit (Geneaid Biotech Ltd., Taiwan) according to the manufacturer's protocol.¹⁵

Sample preparation

Transfer up to 200 µl of whole blood to a 1.5 ml microcentrifuge tube. Adjust the volume to 200 µl with phosphate buffered saline. Add 20 µl of proteinase K then mix by pipetting. Incubate at 60°C for 5 min.

Cell lysis

Add 200 µl of gel sample buffer, then mix by shaking vigorously. Incubate at 60°C for 5 min, add proteinase, then incubate again for 5 min.

DNA binding

Add 200 µl of absolute ethanol to the sample lysate and mix immediately by shaking vigorously for 10 sec. Transfer all of the mixture to the GS Column. Centrifuge at 14–16,000 × g for 1 min. Transfer the GS Column to a new 2 ml collection tube.

Wash

Add 400 µl of W1 buffer to the GS Column. Centrifuge at 14–16,000 × g for 30 sec. Add 600 µl of wash buffer (make sure absolute ethanol was added) to the GS Column. Centrifuge at 14–16,000 × g for 30 sec. Centrifuge for 3 min at 14–16,000 × g.

Elution

Transfer the dried GS Column to a 1.5 ml microcentrifuge tube. Add 100 µl of pre-heated elution buffer, Tris-EDTA buffer. Let stand for at least 3 min. Centrifuge at 14–16,000 × g for 30 sec. DNA quality was evaluated with 2% agarose gel electrophoresis at 100 V (Bio-Rad, USA); the DNA quantity was not calculated.

Polymerase chain reaction (PCR) and sequencing

Screening of germline mutations in the RB1 gene was performed by direct Sanger sequencing for exons 8, 10, 14, 17, and 22. The primer sequences followed the previously published primer sequences for the RB1 gene, as presented in Table 1.³ The DNA was amplified in a PCR DNA thermal cycler (Bio-Rad, USA) in a total volume of 50 µl. The PCR component was KAPPA2G fast enzyme, forward and reverse RB1 gene primer (Table 1), nuclease-free water, and DNA template. PCR cycling was initiated with the denaturation step at 95°C for 15 min, then 94°C for 1 min, annealing at 55°C for 30 sec, extension at 72°C for 1.5 min as much as 21 cycles, continued with the last extension at 72°C for 10 min and 12°C for 30 min for storage. The PCR product was sequenced to detect mutation in

Table 1. Sequence of primers used to amplify exons of the *RB1* gene in Indonesian patients with RB

Primer name	Forward primer	Reverse primer	Length (bp)
RB1 exon 8	5'-GCAGAGTAGAAGAGGGATGGC-3'	5'-TGATTCCAGAGTGAGGGAGC-3'	558
RB1 exon 10	5'-TTTATATTGCATGCGAACTCAG-3'	5'-GGTAACTGTTATAGGACACACAATTC-3'	469
RB1 exon 14	5'-GTGATTTCTAAAATAGCAGG-3'	5'-TGCCTTGACCTCTGATCTG-3'	191
RB1 exon 17	5'-TCAAAATTGGAAGGCTATTCC-3'	5'-TTAGATGGTTAGGGTGCTCG-3'	567
RB1 exon 22 + 23	5'-TCCTTTATAATATGTGCTTCTTACCAG-3'	5'-TTCTTGGATCAAATAATCCCC-3'	577

bp=base pair; RB=retinoblastoma

the *RB1* gene, then analyzed using BioEdit Sequence Alignment Editor version 7.0.5.1. software (Tom Hall; North Carolina State University, USA). The results were then compared with the data in Gene Bank of National Center for Biotechnology Information GRCh37 database using basic local alignment search tool. Additional information about mutations in the *RB1* gene was confirmed from the *RB1* variation database *RB1-Isdb* and the *Human Gene Mutation Database*.

Statistical analysis

The demographic data between normal subjects and patients with RB were compared using the Mann-Whitney *U* test. A *p*-value of <0.05 was considered statistically significant. The statistical analyses were performed using the *JMP* software version 13.0 (SAS Institute Inc., USA).

Ethical considerations

The procedures adhered to the tenets of the Declaration of Helsinki and were approved by the Ethics Committee of the Faculty of Medicine, Universitas Hasanuddin (No. 555/UN4.6.4.5.31/PP36/2019). Written and signed informed consent from the parents was obtained before the examinations.

RESULTS

A total of 21 patients with RB and 14 normal subjects were included in this study. RB patients were mostly diagnosed at a later stage, with 81% of patients exhibiting tumor extension. Additionally, 14% of patients were classified as Group V according to the Reese-Ellsworth classification, and 15% had a regressed type of RB. Of the patients, 76% had unilateral RB, and 24% had bilateral RB. Table 2 presents the demographic data for these groups.

Ten patients with unilateral RB and one with bilateral RB had the *RB1* gene variations. The mean paternal age was 36 years old, and the mean maternal age was 32 years old. Ten out of 11 fathers had a smoking habit, whereas none of the mothers smoked. Furthermore, the parents had no history of radiation exposure or a family history of RB.

In this study, 184 variations were discovered in RB patients, with 164 of which were also found in normal subjects. Nineteen intronic mutations in introns 10, 16, 17, and 21, and one missense mutation in exon 17 were identified; these mutations were not found in normal subjects. One patient (5%) (R33) had a germline mutation among the 21 patients. The heterozygous mutation identified in exon 17 of R33 is presented in Figure 1, which caused an amino acid change from tryptophan to glycine. No family members of R33 had a history of RB. The mutation list of the *RB1* gene in RB patients is shown in Table 3.

The analysis of the parental origin of the detected mutations revealed the *RB1* gene variations in seven out of 21 parents whose children tested positive for the *RB1* gene mutations. The variations included eight substitutions in exon 17 and five intronic mutations in introns 16 and 17, as shown in Table 3. However, none of the variations in exons were passed to their children who had RB.

DISCUSSION

This study conducted a comprehensive screening of germline mutations of the *RB1* gene in Indonesian patients with RB and found a 5% germline RB mutation rate. The prevalence of the *RB1* gene germline mutation varies worldwide, ranging from 37% in Italy to 51% in India among patients with RB. Germline mutation occurs in 51.06, 37.14, and 42.4% of patients with RB in India, Italy, and Singapore, respectively.^{8,9,16}

Table 2. Demographic data of patients with RB and normal subjects

Parameter	RB, n (N = 21)	Normal, n (N = 14)	p*
Patient's characteristics			
Age (years), mean (SD)	3.90 (4.07)	4.07 (4.09)	0.057
Male sex	10	5	0.484
Laterality			
Unilateral	10	-	-
Bilateral	1	-	-
Parent's characteristics			
Paternal age (years), mean (SD)	33.76 (9.34)	31.78 (5.67)	0.829
Maternal age (years), mean (SD)	32.33 (7.36)	32.71 (4.78)	0.855
Paternal smoking habit			
Yes	17	9	0.414
No	4	5	
Maternal smoking habit			
Yes	0	1	0.727
No	21	13	
Paternal history of radiation exposure			
Yes	0	3	0.293
No	21	11	
Maternal history of radiation exposure			
Yes	0	1	0.727
No	21	13	
Paternal family history of RB			
Yes	0	0	-
No	21	14	
Maternal family history of RB			
Yes	0	0	-
No	21	14	

RB=retinoblastoma; SD=standard deviation. *Mann-Whitney U test




Figure 1. Sequencing and pedigree of patient with novel exon mutation. (a) Sequencing result of a patient (R33) with heterozygous mutation of exon 17, c.1687T>G (arrow); (b) sequencing result of normal subjects

Moreover, the rate is higher in Vietnam (73.53%).⁶ The prevalence of germline mutation of the RB1 gene in various countries are presented in Table 4.^{8-10,16-22}

Notably, this study found a novel missense mutation in exon 17 in patients with a unilateral tumor, but no germline mutation was detected in

patients with bilateral RB. A study in Singapore found that 71% of patients with bilateral RB and 29% with unilateral RB had a RB1 gene germline mutation.¹⁶ Meanwhile, these mutations were detected in 44% of Vietnamese patients with a unilateral tumor and 84% with a bilateral tumor.⁶

Exon 17 is crucial as it encodes the domains A and B of the pRB pocket, necessary for repressor activity that blocks cell growth by binding to transcriptional factors E2F/DP.^{4,23} Mutations in this repressor domain may disturb growth suppression and cause loss of cell growth control. The identified *de novo* missense mutation in exon 17 induced amino acid changes from tryptophan to glycine. Tryptophan is an essential amino acid with amphipathic properties, exhibiting weak hydrophilic and hydrophobic characteristics.²⁴⁻²⁶ Consequently, it tends to compose the inner core of soluble proteins.²⁷ Meanwhile, glycine is a non-

Table 3. Data of the *RB1* gene mutations in patients with RB

Patient ID	Exon/intron No	g. position	c. position	Type of mutation	Present in mother/father
R6	Intron 17	g.78390A>T	c.1695+110C>T	Intronic	No/No
R6	Intron 17	g.78425T>A	c.1695+145T>A	Intronic	No/No
R6	Intron 17	g.78426C>A	c.1695+146C>A	Intronic	No/No
R6	Intron 17	g.78430T>G	c.1695+150T>G	Intronic	No/No
M6	Intron 17	g.78301G>A	c.1695+21G>A	Intronic	-
M6	Exon 17	g.78176T>G	c.1592T>G	Missense	-
R8	Intron 17	g.78393G>A	c.1695+113G>A	Intronic	No/No
M8	Intron 16	g.77969DelA	c.1499-113DelA	Intronic	-
M8	Exon 17	g.78108T>C	c.1524T>G	Silent mutation	-
M8	Exon 17	g.78143T>G	c.1559T>G	Missense	-
R10	Intron 17	g.78367C>T	c.1695+87C>T	Intronic	No/NA
R13	Intron 17	g.78301G>A	c.1695+21G>A	Intronic	No/No
R15	Intron 10	g.64575InsC	c.1049+135InsC	Intronic	No/No
M15	Exon 17	g.78130T>G	c.1546T>G	Missense	-
R16	Intron 10	g.64570T>C	c.1049+130T>C	Intronic	No/No
R16	Intron 10	g.64576InsC	c.1049+136InsC	Intronic	No/No
R16	Intron 10	g.64449T>C	c.1049+9T>C	Intronic	No/No
R17	Intron 10	g.64599InsA	c.1049+159InsA	Intronic	No/No
R32	Intron 21	g.161917A>T	c.2212-79A>T	Intronic	No/No
R32	Intron 21	g.161917InsC	c.2212-79InsC	Intronic	No/No
R32	Intron 21	g.161937InsT	c.2212-59InsT	Intronic	No/No
M32	Exon 17	g.78271T>G	c.1687T>G	Missense	-
M32	Intron 17	g.78301G>A	c.1695+21G>A	Intronic	-
M32	Exon 17	g.78108T>G	c.1524T>G	Silent mutation	-
M32	Exon 17	g.78130T>G	c.1546T>G	Missense	-
R33	Intron 16	g.77969DelA	c.1499-113DelA	Intronic	NA/No
R33	Exon 17	g.78271T>G	c.1687T>G	Missense*	NA/No
R33	Intron 17	g.78301G>A	c.1695+21G>A	Intronic	NA/Yes
F33	Intron 17	g.78301G>A	c.1695+21G>A	Intronic	-
R35	Intron 16	g.77969DelA	c.1499-113DelA	Intronic	Yes/No
M35	Intron 16	g.77969DelA	c.1499-113DelA	Intronic	-
R36	Intron 16	g.77969DelA	c.1499-113DelA	Intronic	No/No
M36	Exon 17	g.78130T>G	c.1546T>G	Missense	-

Del=deletion; ID=identity; Ins=insertion; NA=not available; RB=retinoblastoma. All patients had unilateral RB except R8 and had intronic mutation except R33. *Novel mutation

essential hydrophobic amino acid with unique features, including its backbone consisting of only one hydrogen atom, which makes it more flexible than other amino acids.²⁵⁻²⁸ Due to this characteristic, glycine is often located at the interface where two polypeptides come into contact. In the *RB1* gene, even a single amino acid substitution may result in a dysfunctional pRB, compromising its role as a tumor

suppressor gene.²⁹ However, additional analysis such as minigene assay or *in silico* analysis may be necessary to confirm the pathogenicity of a mutation.^{30,31} This was another limitation of this study; we did not perform additional assay and pathogenicity assessment. This low prevalence of germline mutation might be due to the inability to identify mutational change in all *RB1* gene exons, as some “hot-spot” exons (8, 10, 14,

Table 4. Prevalence of the RB1 gene germline mutation in various countries

First author, year	Germline mutation prevalence (%)	Country	Year
Alonso, ¹⁷ 2001	69	Spain	2001
Kiran, ⁸ 2003	14.8 (tumor) and 65 (peripheral blood alone)	India	2003
Sampieri, ⁹ 2006	37	Italy	2006
Kontic, ¹⁸ 2006	43.8	Serbia	2006
Macias, ¹⁹ 2008	64.3	Mexico	2008
Ali, ²⁰ 2010	66.2	Saudi	2010
Barbosa, ²¹ 2013	86.2	Brazil	2013
Ayari-Jeridi, ¹⁰ 2015	18.9	Tunisia	2015
Tomar, ¹⁶ 2017	19.5	Singapore	2017
Parma, ²² 2017	20	Argentina	2017

17, and 22) are more commonly affected.¹⁶ Another possibility was due to the small sample size.

This study compared sequencing results between the RB patients and normal subjects (control) to reduce the possibility of normal gene variants or polymorphism. DNA exon primers were used,³ as well as DNA sequence to read the adjacent intronic parts. Of the 20 mutations discovered, 90% were intronic mutations, with deep intronic mutations accounting for 66.67%. Although only a few bases deep in introns are known to be effective targets for oncogenic mutations,³² point mutations deep within introns (>100 bp from a splice junction) may alter normal splicing, transcription regulatory motifs, and non-coding RNA genes.³³⁻³⁵ Some promoters also collaborate with regulatory sequences within the intron.³⁶

In most cases, RB was unilateral and sporadic, and the parents appeared genetically normal. In some families, carriers had no tumor (reduced penetrance) or only unilateral RB or benign retinocytoma (reduced expressivity).¹¹ These variations in the phenotype may be caused by various factors such as immunologic factors, epigenetic mechanisms, delayed mutation, host resistance, and modulatory genes.³⁷⁻⁴¹

Our findings have implications for prognosis and genetic counseling for the patients and their families. We provided genetic counseling to one pregnant patient with unilateral regressed RB (R3), and no germline mutation was detected. However, we advised her about the possibly inheriting the condition on to her offspring, even though the risk was smaller than for patients with a germline mutation. Since most parents were of their active reproductive age, comprehensive genetic counseling about RB is crucial.

In conclusion, one novel missense mutation was identified in exon 17 of the RB1 gene, which was not detected in normal subjects. This finding has implications for prognosis, genetic counseling, and risk prediction for affected families. Additional studies are needed to confirm the pathogenicity of the mutation and explore other mutational changes in the entire RB1 gene sequencing.

6
Conflict of Interest

The authors affirm no conflict of interest in this study.

Acknowledgment

This study was supported by the Universitas Hasanuddin Research Grant (Grant No.2006/UN4.1/KEP/2019). We thank Handayani Halik and Asep M. Ridwanulloh who provided insight and expertise that greatly assisted the research. English language editing and review were done by John Ellis, MD, Ph.D from University of Dundee, United Kingdom.

Funding Sources

This study was funded by Universitas Hasanuddin Research Grant (Grant No.2006/UN4.1/KEP/2019).

REFERENCES

1. Dimaras H, Corson TW. Retinoblastoma, the visible CNS tumor: a review. *J Neurosci Res.* 2019;97(1):29-44.
2. Ramasubramanian A, Shields CL. Staging and treatment strategies. In: Ramasubramanian A, Shields CL, editors. *Retinoblastoma.* New Delhi: Jaypee Brothers Medical Publishers Ltd; 2012. p. 70-8.
3. Kalsoom S, Wasim M, Afzal S, Shahzad MS, Ramzan S, Awan AR, et al. Alterations in the RB1 gene in Pakistani patients with retinoblastoma using direct sequencing analysis. *Mol Vis.* 2015;21:1085-92.
4. Ganguly A, Nichols K. Genetics of retinoblastoma: molecular and clinical aspects. In: Ramasubramanian A, Shields CL, editors. *Retinoblastoma.* New Delhi: Jaypee Brothers Medical Publishers Ltd; 2012. p. 24-33.
5. Rogers KT, Higgins PD, Milla MM, Phillips RS, Horowitz JM. DP-2, a heterodimeric partner of E2F: identification and characterization of DP-2 proteins expressed in vivo. *Proc Natl*

- Acad Sci U S A. 1996;93(15):7594–9.
6. Nguyen HH, Nguyen HTT, Vu NP, Le QT, Pham CM, Huyen TT, et al. Mutational screening of germline RB1 gene in Vietnamese patients with retinoblastoma reveals three novel mutations. *Mol Vis*. 2018;24:231–8.
 7. Lohmann DR. RB1 gene mutations in retinoblastoma. *Hum Mutat*. 1999;14(4):283–8.
 8. Kiran VS, Kannabiran C, Chakravarthi K, Vemuganti GK, Honavar SG. Mutational screening of the RB1 gene in Indian patients with retinoblastoma reveals eight novel and several recurrent mutations. *Hum Mutat*. 2003;22(4):339.
 9. Sampieri K, Hadjistilianou T, Mari F, Speciale C, Mencarelli MA, Cetta F, et al. Mutational screening of the RB1 gene in Italian patients with retinoblastoma reveals 11 novel mutations. *J Hum Genet*. 2006;51(3):209–16.
 10. Ayari-Jeridi H, Moran K, Chebbi A, Bouguila H, Abbes I, Charradi K, et al. Mutation spectrum of RB1 gene in unilateral retinoblastoma cases from Tunisia and correlations with clinical features. *PLoS One*. 2015;10(1):e0116615.
 11. Harbour JW. Molecular basis of low-penetrance retinoblastoma. *Arch Ophthalmol*. 2001;119(11):1699–704.
 12. Hung CC, Lin SY, Lee CN, Chen CP, Lin SP, Chao MC, et al. Low penetrance of retinoblastoma for p.V654L mutation of the RB1 gene. *BMC Med Genet*. 2011;12:76.
 13. Mallipatna A, Marino M, Singh AD. Genetics of retinoblastoma. *Asia Pac J Ophthalmol (Phila)*. 2016;5(4):260–4.
 14. Temming P, Viehmann A, Biewald E, Lohmann DR. Sporadic unilateral retinoblastoma or first sign of bilateral disease? *Br J Ophthalmol*. 2013;97(4):475–80.
 15. Tanti A. Estimate of mismatch repair deficient pancreatic cancer in patients enrolled in the Quebec pancreas cancer study [master's thesis]. [Montreal]: McGill University; 2019.
 16. Tomar S, Sethi R, Sundar G, Quah TC, Quah BL, Lai PS. Mutation spectrum of RB1 mutations in retinoblastoma cases from Singapore with implications for genetic management and counselling. *PLoS One*. 2017;12(6):e0178776.
 17. Alonso J, García-Miguel P, Abelairas J, Mendiola M, Sarret E, Vendrell MT, et al. Spectrum of germline RB1 gene mutations in Spanish retinoblastoma patients: phenotypic and molecular epidemiological implications. *Hum Mutat*. 2001;17(5):412–22.
 18. Kontic M, Palacios I, Gámez Á, Camino I, Latkovic Z, Rasic D, et al. New RB1 oncogenic mutations and intronic polymorphisms in Serbian retinoblastoma patients: genetic counseling implications. *J Hum Genet*. 2006;51(10):90–13.
 19. Macías M, Dean M, Atkinson A, Jiménez-Morales S, García-Vázquez FJ, Saldaña-Alvarez Y, et al. Spectrum of RB1 gene mutations and loss of heterozygosity in Mexican patients with retinoblastoma: identification of six novel mutations. *Cancer Biomark*. 2008;4(2):93–9.
 20. Ali MJ, Parsam VL, Honavar SG, Kannabiran C, Vemuganti GK, Reddy VA. RB1 gene mutations in retinoblastoma and its clinical correlation. *Saudi J Ophthalmol*. 2010;24(4):119–23.
 21. Barbosa RH, Aguiar FC, Silva MF, Costa RA, Vargas FR, Lucena E, et al. Screening of RB1 alterations in Brazilian patients with retinoblastoma and relatives with retinoma: phenotypic and genotypic associations. *Invest Ophthalmol Vis Sci*. 2013;54(5):3184–94.
 22. Parma D, Ferrer M, Luce L, Giliberto F, Szijan I. RB1 gene mutations in Argentine retinoblastoma patients. Implications for genetic counseling. *PLoS One*. 2017;12(12):e0189736.
 23. Chow KN, Dean DC. Domains A and B in the Rb pocket interact to form a transcriptional repressor motif. *Mol Cell Biol*. 1996;16(9):4862–8.
 24. Hung A, Mager M, Hembury M, Stellacci F, Stevens MM, Yarovsky I. Amphiphilic amino acids: a key to adsorbing proteins to nanopatterned surfaces? *Chem Sci*. 2013;4(3):928–37.
 25. Pánek J, Eidhammer I, Aasland R. A new method for identification of protein (sub)families in a set of proteins based on hydrophathy distribution in proteins. *Proteins*. 2005;58(4):923–34.
 26. Chen YL, Li QZ. Prediction of the subcellular location of apoptosis proteins. *J Theor Biol*. 2007;245(4):775–83.
 27. Karp G. Cell and molecular biology: concepts and experiments. 6th ed. New York: John Wiley & Sons; 2010. Chapter 2, The chemical basis of life. p. 31–84.
 28. Razak MA, Begum PS, Viswanath B, Rajagopal S. Multifarious beneficial effect of nonessential amino acid, glycine: a review. *Oxid Med Cell Longev*. 2017;2017:1716701.
 29. Kaye FJ, Kratzke RA, Gerster JL, Horowitz JM. A single amino acid substitution results in a retinoblastoma protein defective in phosphorylation and oncoprotein binding. *Proc Natl Acad Sci U S A*. 1990;87(17):6922–6.
 30. Imperatore V, Pinto AM, Gelli E, Trevisson E, Morbidoni V, Frullanti E, et al. Parent-of-origin effect of hypomorphic pathogenic variants and somatic mosaicism impact on phenotypic expression of retinoblastoma. *Eur J Hum Genet*. 2018;26(7):1026–37.
 31. Gámez-Pozo A, Palacios I, Kontic M, Menéndez I, Camino I, García-Miguel P, et al. Pathogenic validation of unique germline intronic variants of RB1 in retinoblastoma patients using minigenes. *Hum Mutat*. 2007;28(12):1245.
 32. Zhang K, Nowak I, Rushlow D, Gallie BL, Lohmann DR. Patterns of missplicing caused by RB1 gene mutations in patients with retinoblastoma and association with phenotypic expression. *Hum Mutat*. 2008;29(4):475–84.
 33. Vaz-Drágo R, Custódio N, Carmo-Fonseca M. Deep intronic mutations and human disease. *Hum Genet*. 2017;136(9):1093–111.
 34. Highsmith WE, Burch LH, Zhou Z, Olsen JC, Boat TE, Spock A, et al. A novel mutation in the cystic fibrosis gene in patients with pulmonary disease but normal sweat chloride concentrations. *N Engl J Med*. 1994;331(15):974–80.
 35. McConville CM, Stankovic T, Byrd PJ, McGuire GM, Yao QY, Lennox GG, et al. Mutations associated with variant phenotypes in ataxia-telangiectasia. *Am J Hum Genet*. 1996;59(2):320–30.
 36. Stadhouders R, van den Heuvel A, Kolovos P, Jorna R, Leslie K, Grosveld F, et al. Transcription regulation by distal enhancers: who's in the loop? *Transcription*. 2012;3(4):181–6.
 37. Char DH, Ellsworth R, Rabson AS, Albert DM, Herberman RB. Cell-mediated immunity to a retinoblastoma tissue culture line in patients with retinoblastoma. *Am J Ophthalmol*. 1974;78(1):5–11.
 38. Greger V, Passarge E, Hopping W, Messmer E, Horsthemke B. Epigenetic changes may contribute to the formation and spontaneous regression of retinoblastoma. *Hum Genet*. 1989;83(2):155–8.
 39. Matsunaga E. Hereditary retinoblastoma: delayed mutation or host resistance? *Am J Hum Genet*. 1978;30(4):406–24.
 40. Bremner R, Du DC, Connolly-Wilson MJ, Bridge P, Ahmad KF, Mostachfi H, et al. Deletion of RB exons 24 and 25 causes low-penetrance retinoblastoma. *Am J Hum Genet*. 1997;61(3):556–70.
 41. Valverde JR, Alonso J, Palacios I, Pestaña A. RB1 gene mutation up-date, a meta-analysis based on 932 reported mutations available in a searchable database. *BMC Genet*. 2005;6:53.

Novel point mutation and intronic mutations of RB1 gene in retinoblastoma patients in Indonesia

ORIGINALITY REPORT

20%

SIMILARITY INDEX

17%

INTERNET SOURCES

18%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Negeri Semarang Student Paper	2%
2	journals.plos.org Internet Source	2%
3	www.protocols.io Internet Source	1%
4	Submitted to Kirkuk University Student Paper	1%
5	mail.scialert.net Internet Source	1%
6	Mei Riasanti, Ikhwanuliman Putera, Priscilla Jessica, Muhammad Zakiy Waliyuddin et al. "Interferon-gamma release assay and chest X-ray to classify intraocular tuberculosis among clinically undifferentiated uveitis", Medical Journal of Indonesia, 2023 Publication	1%
7	prp.hec.gov.pk Internet Source	1%

8

Sri Anggarini Rasyid, Sanatang Sanatang, Satriani Syarif, Sugireng Sugireng et al. "Effect of *Batissa violacea celebensis* Martens, 1897, Extract on β -Catenin Gene in Mice (*Mus musculus*) Induced by Azoxymethane and Dextran Sulfate Sodium using Polymerase Chain Reaction-restriction Fragment Length Polymorphism Method", Open Access Macedonian Journal of Medical Sciences, 2021

Publication

1 %

9

Muhammad Adi Satrio Lazuardi, Andrie Rhomdhon Kurniawan, Pradana Nurhadi, Besut Daryanto. "Multiple vaginal stones secondary to an ectopic ureter in an adult female patient: a case report", Medical Journal of Indonesia, 2023

Publication

1 %

10

revistabiomedica.org

Internet Source

1 %

11

www.frontiersin.org

Internet Source

1 %

12

Michal Sagi, Avishag Frenkel, Avital Eilat, Naomi Weinberg, Shahar Frenkel, Jacob Pe'er, Dvorah Abeliovich, Israela Lerer. "Genetic screening in patients with Retinoblastoma in Israel", Familial Cancer, 2015

Publication

1 %

13	www.currentpediatrics.com Internet Source	1 %
14	labs.feinberg.northwestern.edu Internet Source	1 %
15	mafiadoc.com Internet Source	1 %
16	Robert E. Ferrell. "Candidate Gene Analysis in Primary Lymphedema", Lymphatic Research and Biology, 06/2008 Publication	1 %
17	academic.oup.com Internet Source	1 %
18	www.hindawi.com Internet Source	1 %
19	repositori.uin-alauddin.ac.id Internet Source	1 %
20	"Advances in Vision Research, Volume I", Springer Nature, 2017 Publication	1 %
21	www3.unisi.it Internet Source	1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

