



- [SEARCH](#)
- [DASHBOARD](#)
- [PAPER SUBMISSION](#)
- [SUBSCRIPTION](#)
- [BOOKS](#)
- [BOOK ADVERTISEMENT](#)
- [MY CART \(0\)](#)
- [MY ORDERS](#)

Paper ID: 2356034457024539
 Title: Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (Chromolaena odorata L.)
 Author(s): Karlina Amir Tahir;Erwin Hafid;Muh. Fitrah;Ahmad Lalo;Nurul Fadilah;Syamsuri Syakri;Syatirah Jalaluddin;Katsuyoshi Matsunami
 (karlina.amir@uin-alauddin.ac.id;erwin.hafid@uin-alauddin.ac.id;muh.fitrah@uin-alauddin.ac.id;ahmadlalo_pom@yahoo.co.id;syamsurisyakri@gmail.com;syatirahjalaluddin@gmail.com;)
 Submitted To: Research Journal of Pharmacy and Technology
 Payment Transaction No.: pay_NVWeQiODHeeiGO
 Paid On: 2/1/2024 12:00:00 AM

INR: 3540.00
 Or USD: 210.00

PAYMENT VERIFIED (03-Feb-2024)

- [Editorial Comments](#)
- [Reviewer Comments](#)
- [Download Acceptance Letter](#)
- [Print Report](#)

6/25/24, 6:32 PM

Article Status



A and V Publications

Report Generated on: 6/25/2024 4:56:55 PM

BASIC INFORMATION

Paper ID: 2356034457024539

Submission Date: May 06, 2023

Paper Title: Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (Chromolaena odorata L.)

Author(s) Name: Karlina Amir Tahir;Erwin Hafid;Muh. Fitrah;Ahmad Lalo;Nurul Fadilah;Syamsuri Syakri;Syatirah Jalaluddin;Katsuyoshi Matsunami

Author(s) Email: karlina.amir@uin-alauddin.ac.id;erwin.hafid@uin-alauddin.ac.id;muh.fitrah@uin-alauddin.ac.id;ahmadlalo_pom@yahoo.co.id;syamsurisyakri@gmail.com;syatirahjalaluddin@gmail.com;

Author(s) Address: H.M Yasin Limpo No.36, Gowa, South of Sulawesi, Indonesia

Journal: Research Journal of Pharmacy and Technology

Submitted By: karlina amir tahir

Email ID: nilalfath@gmail.com

REVIEWER INFORMATION

	First Reviewer	Second Reviewer
Name:	SUNANDAR IHSAN	PROF. DR. KUMARASWAMY GANDLA
Email ID:	sunandarihsan@uho.ac.id	drkumaraswamygandla@gmail.com
Mobile No:	081342987228	0919000973789
Address:	Universitas Hala Oleo, Kendari, Indonesia	Assoc. editor of RJPT

PAYMENT DETAILS

Order No.: 2024211219014432030539092

Order Date: February 01, 2024, 12:02

Amount INR: 3540.00

OR Amount USD: 210.00

Payment Date: February 01, 2024

Transaction No.: pay_NVWeQiODHeeiGO

Payment Status: PAYMENT VERIFIED

PAPER PUBLICATION / PROCESSING STATUS

- ▶▶ 06/May/2023, 03:44:57 AM --- Article submitted by the author.
- ▶▶ 10/May/2023, 11:32:54 AM --- Article sent back to author for minor corrections.
- ▶▶ 10/May/2023, 11:32:54 AM --- New comments from editorial board.
- ▶▶ 14/May/2023, 07:01:16 AM --- Article resubmitted by author after correction.
- ▶▶ 26/Jul/2023, 06:55:42 PM --- Article is sent to reviewers.
- ▶▶ 13/Sep/2023, 04:53:18 PM --- Article sent back to author for minor corrections.
- ▶▶ 13/Sep/2023, 04:53:28 PM --- New comments from editorial board.
- ▶▶ 20/Sep/2023, 04:36:22 AM --- Article resubmitted by author after correction.
- ▶▶ 20/Sep/2023, 04:36:26 AM --- Article resubmitted by author after correction.
- ▶▶ 23/Oct/2023, 06:02:46 PM --- Article sent back to author for minor corrections.
- ▶▶ 23/Oct/2023, 06:05:25 PM --- New comments from editorial board.
- ▶▶ 31/Oct/2023, 12:32:10 PM --- Article resubmitted by author after correction.
- ▶▶ 31/Oct/2023, 12:32:16 PM --- Article resubmitted by author after correction.
- ▶▶ 31/Oct/2023, 12:32:23 PM --- Article resubmitted by author after correction.
- ▶▶ 31/Oct/2023, 12:32:31 PM --- Article resubmitted by author after correction.
- ▶▶ 25/Dec/2023, 03:00:03 PM --- Article sent back to author for minor corrections.
- ▶▶ 25/Dec/2023, 03:00:14 PM --- New comments from editorial board.
- ▶▶ 04/Jan/2024, 03:49:54 AM --- Article resubmitted by author after correction.
- ▶▶ 25/Jan/2024, 05:54:10 PM --- Final version of article is required.
- ▶▶ 27/Jan/2024, 04:46:09 AM --- Final version of article submitted by author.
- ▶▶ 27/Jan/2024, 04:46:09 AM --- Article is accepted by publisher.
- ▶▶ 27/Jan/2024, 04:46:13 AM --- Final version of article submitted by author.
- ▶▶ 27/Jan/2024, 04:46:13 AM --- Article is accepted by publisher.

<https://www.anvpublication.org/ArticleStatus.aspx?PID=2356034457024539>

6/25/24, 6:32 PM

Article Status

- ▶▶ 01/Feb/2024, 12:19:02 PM --- Payment proof required.
- ▶▶ 01/Feb/2024, 12:49:17 PM --- Payment proof uploaded by author.
- ▶▶ 03/Feb/2024, 05:14:22 PM --- Payment details verified by publisher.
- ▶▶ 05/Feb/2024, 04:10:48 PM --- Payment details verified by publisher.

Gmail interface showing an email from Research Journal of Pharmacy and Technology. The email content includes a thank you message to the author and contact information for the editor.

thank you

Research Journal of Pharmacy and Technology <editor.rjpt@gmail.com>
to me

Oct 18, 2023, 2:34 PM

Dear Author,
Your article has been Sent to Reviewers Comments awaited.
thanks
editor

--
Editor,
Research Journal of Pharmacy and Technology,
RJPT House, Lokmanya Githi Nirman Society,
Rohanipuram, In-front of Sector- 1,
Pt. Deendayal Upadhyay Nagar,
Raipur 492 010. (CG) India
Phone No. +919406051618

www.ajronline.org
www.anypublication.org
www.rjptonline.org

Type of Manuscript:

Research

**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT)
Cells from Siam Weeds (*Chromolaena odorata* L.)**

ABSTRACT

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. *Chromolaena odorata* was extracted with methanol, then partitioned with 5 dissolutions: n-Hexane, Ethyl Acetate, Acetonitrilic, n-Buthanol, Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocytes cells using the bioassay method. After incubation, the result obtained from microplate reader. Each extract was divided into three concentrations, it is 100, 50, 20 ($\mu\text{g}/\text{mL}$). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10 ($\mu\text{g}/\text{mL}$). After being analyzed with the results of the microplate reader, the IC50 of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100 $\mu\text{g}/\text{mL}$. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μl of medium/well.

KEYWORDS: *Chromolaena odorata* L., Skin Cancer, HacaT cell line.

Commented [S11]: The research objectives must be stated clearly before the method statement

Commented [S12]: Results and conclusions must be clearly stated

INTRODUCTION

Skin cancer is one of the most common forms of cancer worldwide, the impenetrable topical barrier is compromised slightly in pathologies such as skin cancer. Another disease that has seen growth in local therapeutic applications is skin cancer which is the most common cancer with one in every three cancer diagnoses being skin cancer according to World Health Organization (WHO)¹. Skin cancer is broadly classified into non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinomas and melanomas².

Cancer development usually occurs when DNA is damaged, and our bodies are unable to repair this damage. These damaged cells begin to grow and divide uncontrollably. Such changes in the skin lead to skin cancer. When damage cell multiply, they form tumors².

Skin cancer develops in various layers of the skin. The tendency to develop skin cancer varies for different skin types. Occurrence varies according to skin color and skin type. In addition, heredity also contributes to the development of skin cancer. The most inherited type of cancer is Melanoma².

Skin cancer can be identified through symptoms. Early identification and early detection of medicine and cure. Sometimes symptoms appear at an advanced stage in other types of cancer. Basal cell cancer appears as a lump on the skin anywhere on the body exposed to sunlight. They can be brown bumps or white bumps on our face, hands, neck, and other exposed body parts. Squamous cell carcinoma on the other hand can appear as ulcerated lumps and patches on the skin whereas melanoma can cause unusual growths on the body especially in existing moles. It can be anywhere in our body³.

The American Cancer Society (ACS) has reported that, there were 87,110 newly diagnosed melanomas, also reported that 9,730 people had an estimated melanoma death in 2017. Skin pigmentation is caused by accumulation of melanin granules which are synthesized in melanocytes and transferred to keratinocytes⁴. Keratinocytes secrete cytokines and chemokines on exposure to UV light which affect melanin production⁵. Melanoma develops at any part of the skin due to defects in skin pigmentation but is most susceptible to the trunk in men and the legs in women⁶.

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. Before the body can recover, these deleterious mutations occur in the genes and the damage becomes irreparable. Moreover, to environmental factors, the patient's genetic history can also be a cause of developing cancer⁷.

UV rays consist of UV-A and UV-B radiation. Although UV-B is considered more dangerous, UV-A also has side effects. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-A can also cause DNA damage by generating free radicals and damaging cell membranes which in turn causes skin cancer. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. UV-B skin also causes oxidative damage to the skin. UV-B radiation further interferes with regulatory proteins, anti-apoptotic relational proteins, proapoptotic proteins, transcription factors, protein kinase, inflammatory enzymes, and growth factors which in turn disrupts the cell cycle⁸. This further causes reasonable mutations in various tumor suppressor genes and oncogenes. These effects together lead to uncontrolled keratinocytes mitosis and initiation of skin cancer upon UV exposure⁷.

It is impossible to completely eradicate cancer, but precautions can be taken to prevent cancer. The following steps should be taken; there should be a reduction in sun exposure, use of tobacco should be avoided, sunscreen should be used to prevent exposure of UV rays, and skin cancer cells can also be removed by biopsy⁸.

The Laruna leaf has been used empirically as a traditional medicine by the people of South Sulawesi. Traditional use of *Chromolaena odorata* (L) as a source of medicine has been widely used in West African countries, Asian countries⁹, and Indonesia as wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, anti-immunosuppressive, anticancer, and other activities^{10,11,12}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals (various factors in plant foods), such as carotenoids, antioxidant vitamins, phenolic compounds, terpenoids, steroids, indole, and fiber, etc¹³. Crude drugs are the effective elements considered responsible for reducing cancer risk¹⁴.

Chromolaena odorata has bioactive compounds with a significant therapeutics' effects on some therapeutic effects on some pathological conditions^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40 -trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihidrokaempferide, Apigenin-7,40 -dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin- 40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

Commented [SI3]: data in Indonesia would be better if it was conveyed

Commented [SI4]: The research objectives must be stated clearly before the method statement

MATERIALS AND METHODS

Materials:

Chromolaena odorata collection is in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, the 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitrilic, N-Buthanol, Ethanol.

Commented [S15]: Composition, collection, storge of sample would be better stated clearly

Table 1. Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77.621	1.55

Table 2. Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68.45
	Ethyl Acetate	245.89
	Acetonitrilic	85.22
	n-Buthanol	147.97
	Ethanol	53.22

Cell Culture

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10 mg of each of extract were dissolved in 1 mL of DMSO. The second stock solution (5 mg/mL) was generated by mixing 0.5 mL of this stock solution with 0.5 mL of DMSO, which was mixed well. The third stock solution (2 mg/mL) was mixed by 0,2 mL of DMSO. To make the final serial concentrations, 1 µL aliquots of the sample stock solutions in two-fold dilution (10, 5, 2 mg/mL) were added to the corresponding well (the final volume became 100 µL) to obtain the final concentrations (100, 50, 20 µg/mL)

Cytotoxicity Investigation

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1 µL of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10 µg/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = \left[\frac{(A_{\text{test}} - A_{\text{blank}})}{(A_{\text{control}} - A_{\text{blank}})} \right] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis

Table 3. % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4. Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test (([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100 µg/mL.




Figure 1. HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic,
D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate (([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹⁶. IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISCUSSION:

The aim of this study was to investigate the potential of natural products using HaCaT cells as a reliable in vitro model for evaluation.

Keratinocytes are the main cells of the human epidermis. In addition to providing an epidermal barrier function, keratinocytes play a role in the inflammatory response of the skin¹⁷, and the process of wound repair in the re-epithelialization step¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) eliminated the problems observed with primary keratinocytes such as their need for growth supplementation, culture differentiation, donor-related variability, and limited lifespan. HaCaT cells have been used for many skin model studies without altering keratinocyte function, cellular response, or differentiation capacity¹⁹. Monolayer HaCaT cell cultures are used for many applications such as cellular toxicity and in vitro analysis of wound healing. However monolayer culture systems cannot completely replicate the true epidermal structure. Whereas the generation of air-liquid interface in keratinocytes cell cultures might result in a more realistic model of the epidermis²⁰.

Keratinocytes are active in epidermal repair and skin immune defense through secretion of growth factors, cytokines, and chemokines. To standardizing in vitro studies, investigated the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, and in relation to different culture conditions and degrees of differentiation¹⁷.

HaCaT cells are a long-lived, spontaneously immortalized human line that exhibits basal cell properties and displays substantial changes in response to two well-established in vitro prodifferentiating agents: increased cell density and as a function of time in culture¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 µl of medium/well¹⁷.

HaCaT cells were generally maintained in normal DMEM supplemented with 10% FBS, both of which contain sufficient calcium to induce differentiation. Dedifferentiation to the basal phenotype requires prolonged growth in low calcium media²².

Statistical evidence has revealed that most deaths from skin cancer are the result of melanoma. Further investigations have shown that the survival rate in patients depends on the stage of the cancer; Early detection and intervention of melanoma involves a higher chance of cure²³.

Topical therapies are delivered through skin & they reach the systemic circulation by crossing the skin layers. There are some commonly used topical agent which includes some corticosteroids is given as: 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoids, Mechlorethamine, Carmustine, Bexarotene, Tazarotene photodynamic^{24,25}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals, such as antioxidant. *Chromolaena odorata* leaf extract contain various antioxidant, terpenoid, phenolic and

Commented [SI6]: an incomplete sentence stating the purpose of the study

Commented [SI7]: Differences with other research can be clarified by comparing it with previous studies on similar topics.

flavonoid compounds²⁶. In addition, the anti-mitotic activity of the results of this investigation shows that silver green nanoparticles synthesized from *Chromolaena odorata* have a potential effect on cell division²⁷. The phytochemical profile of the phenolic extract of *Chromolaena odorata* has great potential for the identification of new compounds with pharmacotherapeutic value²⁸. As a comparison to *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48%, research on *Helicteres isora* extract at a concentration of 300 µg/ml had an activity equivalent to the activity of the standard drug doxorubicin²⁹.

In addition, regarding skin cancer testing with various extracts, for example, Ethosomal gel which produces controlled release of curcumin, Curcumin persists in deeper layers of the skin for a longer time, as well as maximum deposition. Thus, eliminating melanoma cells completely³⁰. Furthermore, on the carcinogenic anti-cancer potential of diosmin in vivo and in vitro, it can be a flavonoid in the treatment of Hepatocellular Carcinoma due to its therapeutic and pharmacological properties. Although this research requires further research on other types of cancer³¹.

In-depth study of melanoma biology has been influential in considering melanoma as a therapy for one of the most resistant malignancies. In the last 30 years, there have been few options for treating metastatic melanoma, so some more thorough clinical studies have been conducted, such as hormonal therapy, immunotherapy, single and combination agent chemotherapy, biochemotherapy, targeted agent therapy, and combination regimens, show less than optimal results³².

CONCLUSION:

Ethyl acetate extract of *Chromolaena odorata* has anti-skin cancer potential regarding to the cytotoxicity test.

CONFLICT OF INTEREST:

There are no conflicts of interest declared by the authors of this study.

REFERENCES

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashed D, Weller RB, Castellsagué X. Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)*. 2017; 7(Suppl 1):5-19. doi: 10.1007/s13555-016-0165-y
3. Akila Victor, Muhammad Rukunuddin Ghalib. Detection of skin cancer cells – a review. *Research J.Pharm and Tech*; 2017; 10(11):4093-4098. doi: 10.5958/0974-360X.2017.00742.9
4. Scott G. Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Invest Dermatol*. 2012; 132(4):1073-4. doi: 10.1038/jid.2012.20
5. Nasti TH, Timares L, MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*. 2015; 91(1):188-200. doi: 10.1111/php.12335
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan. Protein-protein interaction of mutated agouti signaling protein (asip) to melanocortin receptor 1 (mcr1) in melanoma skin cancer: an insilico study. *Research J. Pharm. and Tech* 2018; 11(9): 3913-3917. doi: 10.5958/0974-360X.2018.00718.7
7. Min Jeong Kim 1, Seo Yeon Shin 1, Nu Ri Song 1, Sunoh Kim 2, Sang Ouk Sun 1 and Kyung Mok Park 1. Bioassay-guided characterization, antioxidant, anti-melanogenic and anti-photoaging activities of pueraria thunbergiana l. leaf extracts in human epidermal keratinocytes (hacat) cells. *MDPI* 10. 2022; 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y. New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*. 2014; 2014:243452. doi: 10.1155/2014/243452
9. Qadir MI. Skin cancer: etiology and management. *Pak J Pharm Sci*. 2016; 29(3):999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med*. 12. 2015; 301–306. <https://doi.org/10.1515/jcim-2014-0033>
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed. Biological activities of *chromolaena odorata*: A mechanistic review. Elsevier. 2021; <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S. Flavonoid glycosides from *chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull*. 59. 2011; 129–131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B. Pharmacological importance of *chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal Vol 9*. 2021; 8–11. Doi:10.47957/ijpda.v9i1.452

14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging, and antimicrobial potential of chromolaena odorata leaf extracts against pathogenic bacterium in wound infections a multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. 20. <https://doi.org/10.1016/j.bcab.2018.05.014>
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J. Kaempferide, the most active among the four flavonoids isolated and characterized from chromolaena odorata, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5. 2015; 100912– 100922. <https://doi.org/10.1039/C5RA19199H>
16. Mkhobongo B, Chandran R, Abrahamse H. In vitro photodynamic treatment modality for a375 melanoma cell line using a sulphonated aluminum phthalocyanine chloride-photosensitizer-gold nanoparticle conjugate. *Pharmaceutics.* 2022; 16;14(11):2474. doi: 10.3390/pharmaceutics14112474
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all. Hacat cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediator of Inflammation.* 2017; <https://doi.org/10.1155/2017/7435621>
18. Pastar I, Stojadinovic O, Yin NC, Ramirez H, Nusbaum AG, Sawaya A, Patel SB, Khalid L, Isseroff RR, Tomic-Canic M. Epithelialization in wound healing: a comprehensive review. *Adv Wound Care (New Rochelle).* 2014; 3(7):445-464. doi: 10.1089/wound.2013.0473
19. Micallef L, Belaubre F, Pinon A, Jayat-Vignoles C, Delage C, Charveron M, Simon A. Effects of extracellular calcium on the growth-differentiation switch in immortalized keratinocyte hacat cells compared with normal human keratinocytes. *Exp Dermatol.* 2009; 18(2):143-51. doi: 10.1111/j.1600-0625.2008.00775.x
20. Pedrosa TDN, Catarino CM, Pennacchi PC, Assis SR, Gimenes F, Consolaro MEL, Barros SBM, Maria-Engler SS. A new reconstructed human epidermis for in vitro skin irritation testing. *Toxicol In Vitro.* 2017; 42:31-37. doi: 10.1016/j.tiv.2017.03.010
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R. Anti-inflammatory, and anti-proliferative properties of Chromolaena odorata leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants.* 20. 2014; 359–371. doi:10.1080/10496475.2013.876698
22. Wilson VG. Growth and differentiation of hacat keratinocytes. *Methods Mol Biol.* 2014; 1195:33-41. doi: 10.1007/7651_2013_42
23. Tammineni Sreelatha, M. V. Subramanyam, M. N. Giri Prasad. A survey work on early detection methods of melanoma skin cancer. *Research J. Pharm. and Tech.* 2019; 12(5):2589-2596. doi: 10.5958/0974-360X.2019.00435.9
24. Micali G, Lacarubba F, Nasca MR, Ferraro S, Schwartz RA. Topical pharmacotherapy for skin cancer: part ii. Clinical applications. *Journal of the American Academy of Dermatology.* 2014; 70(6):979.e1-12; quiz 9912. doi: 10.1016/j.jaad.2013.12.037.
25. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kanya Goyal, Shabby Jindal. Treatment of skin cancer by topical drug delivery of nanoparticles: a review. *Research Journal of Pharmacy and Technology.* 2021; 14(10):5589-8. doi: 10.52711/0974-360X.2021.00973
26. Zainab, Nurlailah, Muhammad Ikhwan Rizki. Identification of active compound and antibacterial activity against gram-positive and gram-negative bacteria of chromolaena odorata leaf extract. *Research Journal of Pharmacy and Technology.* 2022; 15(10):4720-6. doi: 10.52711/0974-360X.2022.00793
27. Nishita Gogia, Anandhi D, Kanaga G, Revathi K. In vitro evaluation of anti-mitotic activity of green synthesized silver nanoparticles from ethanolic extract of chromolaena odorata, caesalpinia coriaria (bark and leaves) using allium cepa roots. *Research Journal of Pharmacy and Technology.* 2021; 14(8):4107-0. doi: 10.52711/0974-360X.2021.00711
28. Eze FN, Jayeoye TJ. Chromolaena odorata (Siam weed): A natural reservoir of bioactive compounds with potent anti-fibrillogenic, antioxidative, and cytocompatible properties. *Biomed Pharmacother.* 2021;141:111811. doi: 10.1016/j.biopha.2021.111811.
29. Madhulika Pradhan, S Sribhuaneswari, D Karthikeyan, Sunita Minz, Pavani Sure, Atul N Chandu, Umesh Mishra, K Kamalakannan, A Saravanankumar, T Sivakumar. In-vitro cytoprotection activity of foeniculum vulgare and helicteres isora in cultured human blood lymphocytes and antitumour activity against b16f10 melanoma cell line. *Research J. Pharm. and Tech.* 2018; 1(4):450-452
30. Radha Krishna Kollipara, Vyshnavi Tallapaneni, Bharat Kumar Reddy Sanapalli, Vinoth Kumar G, Veera Venkata Satyanarayana Reddy Karri. Curcumin loaded ethosomal vesicular drug delivery system for the treatment of melanoma skin cancer. *Research Journal of Pharmacy and Technology.* 2019; 12(4):1783-1792. doi: 10.5958/0974-360X.2019.00298.1
31. Perumal S, Langeswaran K. Diosmin anti-tumour efficacious against hepatocellular carcinoma. *Research J. Pharm. and Tech.* 2020; 13(4):1707-1714. doi: 10.5958/0974-360X.2020.00308.X
32. Ghadge Ojaswi, Nadar Divya, Parmar Digna. Melanoma and its drug targets. *Research J. Pharm. and Tech.* 2016; 9(5): 562-570. doi: 10.5958/0974-360X.2016.00107.4



Date: January 27, 2024

Research Journal of Pharmacy and Technology

Paper ID:2356034457024539

Author's Name:Karlina Amir Tahir, Erwin Hafid, Muh. Fitrah, Ahmad Lalo, Nurul Fadilah, Syamsuri Syakri, Syatirah Jalaluddin, Katsuyoshi Matsunami

Paper Title: Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (Chromolaena odorata L.)

Acceptance of Manuscript

With reference to your article titled '**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (Chromolaena odorata L.)**' Author by **Karlina Amir Tahir, Erwin Hafid, Muh. Fitrah, Ahmad Lalo, Nurul Fadilah, Syamsuri Syakri, Syatirah Jalaluddin, Katsuyoshi Matsunami**. We wish to bring to your kind notice the following:

- We acknowledge the receipt of the above mentioned article.
- The above mentioned article has been sent to the reviewer of expert comments.
- The above mentioned article have been accepted for publication in Year : 2024, Vol: 17, Issue:5 of the journal.

Thanking you for submission of manuscript.

Yours sincerely,



Dr. Baharwal

Editor in Chief

Note: -

1. To verify the originality of this certificate please visit the following link:

http://www.anvpublication.org/Download_Acceptance.aspx?PID=2356034457024539

PRINT