

turnitin turnitin

DOC-20240508-WA0025..pdf

 NO REPOSITORY PAPER CHECK

 Chceker

Document Details

Submission ID

trn:oid::1:3009447510

Submission Date

Sep 16, 2024, 9:47 AM GMT+7

Download Date

Sep 16, 2024, 9:48 AM GMT+7

File Name

DOC-20240508-WA0025..pdf

File Size

161.1 KB

5 Pages

3,831 Words

21,679 Characters





15% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.




Filtered from the Report

- Bibliography

Match Groups

-  **36 Not Cited or Quoted 14%**
Matches with neither in-text citation nor quotation marks
-  **3 Missing Quotations 1%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 11%  Internet sources
- 8%  Publications
- 5%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- **36 Not Cited or Quoted 14%**
Matches with neither in-text citation nor quotation marks
- **3 Missing Quotations 1%**
Matches that are still very similar to source material
- **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 11% Internet sources
- 8% Publications
- 5% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

| | | | |
|----|-------------|--|----|
| 1 | Publication | Ludovic Micallef. "Effects of extracellular calcium on the growth-differentiation s... | 1% |
| 2 | Internet | repository.unair.ac.id | 1% |
| 3 | Publication | A. Sakunpak, K. Matsunami, H. Otsuka, P. Panichayupakaranant. "Isolation of ne... | 1% |
| 4 | Internet | escholarship.org | 1% |
| 5 | Publication | Vinu Krishnan, Samir Mitragotri. "Nanoparticles for topical drug delivery: Potenti... | 1% |
| 6 | Internet | www.researchgate.net | 1% |
| 7 | Internet | phcogj.com | 1% |
| 8 | Internet | pubmed.ncbi.nlm.nih.gov | 1% |
| 9 | Publication | Yuya Nomoto, Liva Harinantenaina, Sachiko Sugimoto, Katsuyoshi Matsunami, Hi... | 1% |
| 10 | Internet | www.dovepress.com | 1% |

| | | | |
|----|----------------|--|----|
| 11 | Publication | Carolina Motter Catarino, Tatiana do Nascimento Pedrosa, Paula Comune Pennac... | 1% |
| 12 | Student papers | Mendocino College | 1% |
| 13 | Student papers | University of Hertfordshire | 1% |
| 14 | Student papers | Syiah Kuala University | 0% |
| 15 | Publication | Josiane Aline Monteiro de Oliveira, Darlon Irineu Bernardi, Rodolfo Bento Balbino... | 0% |
| 16 | Internet | jvi.asm.org | 0% |
| 17 | Internet | accp2017.ff.unair.ac.id | 0% |
| 18 | Internet | fjps.springeropen.com | 0% |
| 19 | Internet | www.ack1inhibitor.com | 0% |
| 20 | Internet | www.graphyonline.com | 0% |
| 21 | Internet | api-journal.accscience.com | 0% |
| 22 | Internet | ebin.pub | 0% |
| 23 | Internet | worldwidescience.org | 0% |
| 24 | Internet | 1library.net | 0% |

| | | | |
|----|-------------|--|----|
| 25 | Publication | Maria Gabriela Rolim Silveira Sá, Celso Martins Queiroz-Junior, Paulo Eduardo Ale... | 0% |
| 26 | Internet | core.ac.uk | 0% |
| 27 | Internet | www.pharmaexcipients.com | 0% |
| 28 | Internet | www.techscience.com | 0% |
| 29 | Publication | Hong-Fu Li, Han Feng, Yong Wang, Zhang-Chao Pan et al. "Evaluation of hemostat... | 0% |

ISSN 0974-3618 (Print)
0974-360X (Online)

www.rjptonline.org



RESEARCH ARTICLE

Inhibition of Skin Cancer using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)

Karlina Amir Tahir¹, Erwin Hafid², Muh. Fitrah¹, Ahmad Lalo³, Nurul Fadilah¹,
Syamsuri Syakri¹, Syatirah Jalaluddin¹, Katsuyoshi Matsunami⁴

¹Faculty of Medicine and Health Science, UIN Alauddin Makassar, Makassar, Indonesia.

²Master of Islamic Education Management, UIN Alauddin Makassar, Makassar, Indonesia.

³National Agency of Drug and Food Control, Makassar, Indonesia.

⁴Department of Pharmacognosy, Faculty of Biomedical and Health Science at Hiroshima University, Japan.

*Corresponding Author E-mail: karlina.amir@uin-alauddin.ac.id

ABSTRACT:

Cancer can usually develop due to exposure to sunlight. UV radiation from sunlight is known to damage DNA and is bad for the skin. Skin P stem cell carcinogenesis is caused by UV-A rays that penetrate deep into the dermis layer. UV-B damages cell DNA by being absorbed by proteins in the epidermis. *Chromolaena odorata* was extracted using methanol solvent, then partitioned into 5 solutions in the form of n-Hexane, Ethyl Acetate, Acetonitrile, n-Butanol, and Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocyte cells using the bioassay method. Results obtained from the microplate reader after incubation. Each extract was divided into three concentrations, it is 100, 50, 20(µg/mL). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10(µg/mL). After being analyzed with the results of the microplate reader, the IC₅₀ of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100µg/mL. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100µl of medium/well.

KEYWORDS: *Chromolaena odorata* L, Skin Cancer, HacaT cell line.

INTRODUCTION:

One of the most common types of cancer in the world is skin cancer, which is characterized by the breakdown of the airtight topical barrier. According to the World Health Organization (WHO), skin cancer is a cancer that often occurs in one in three cancer diagnoses, and the application of local treatment for this disease is experiencing increasing development¹. Based on the classification, skin cancer consists of melanoma and non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinoma².

Cancer generally develops due to DNA damage that our body cannot repair. Uncontrolled, these injured cells begin to grow and divide. When these changes occur in the skin, they can cause skin cancer. The proliferation of damaged cells can cause tumors².

Skin cancer can arise in any of the skin's layers. Different skin types have differing propensities to develop skin cancer. The frequency varies according to the kind and color of the skin. Furthermore, genetics plays a role in the development of skin cancer. Melanoma is the most common hereditary kind of cancer².

The symptoms that appear can be used to diagnose skin cancer. Identify and detect the drugs and therapies used early. In other forms of cancer, the appearance of symptoms occurs in the advanced stages of the disease.

Received on 06.05.2023 Modified on 21.10.2023
Accepted on 27.01.2024 © RJPT All right reserved
Research J. Pharm. and Tech 2024; 17(5):1951-1955.
DOI: 10.52711/0974-360X.2024.00309

On any part of the body exposed to sunlight, basal cell carcinoma manifests as a lump on the skin. On the hands, face, neck, and other exposed areas of the body, they may appear as white or brown bumps. Melanoma can cause unusual developments on the body, especially on existing moles, while squamous cell carcinoma can manifest as lumps and ulcerated patches on the skin. This can happen anywhere in our bodies³.

According to data from the American Cancer Society (ACS), 9,730 persons lost their lives to melanoma in 2017 and there were 87,110 newly diagnosed melanomas. Melanin granules, which are produced in melanocytes and transported to keratinocytes, accumulate and result in skin pigmentation⁴. When exposed to UV radiation, keratinocytes release cytokines and chemokines that influence the formation of melanin⁵. Skin pigmentation abnormalities can cause melanoma to grow on any area of the skin, although in men and women it most commonly affects the trunk and legs⁶.

One of the biggest factors that causes cancer to develop is excessive exposure to sunlight. It is known that UV radiation from sunlight can injure the skin and cause DNA to be damaged. These dangerous genetic changes occur before the body has time to recover, causing permanent damage. Apart from that, cancer can also be caused by environmental factors and the patient's genetic background⁷.

UV radiation is divided into UV-A and UV-B photons. Although UV-B is considered more dangerous, UV-A can also have negative consequences. Skin P stem cell carcinogenesis is caused by UV-A rays that penetrate deep into the dermis. By producing free radicals and damaging cell membranes, UV-A can also damage DNA, which can lead to skin cancer. UV-B damages cell DNA by being absorbed by proteins in the epidermis. Skin exposed to UV-B rays also experiences oxidative damage. The cell cycle is further disrupted by UV-B radiation interference with growth factors, transcription factors, anti-apoptotic relational proteins, pro-apoptotic proteins, inflammatory enzymes, and regulatory proteins⁸. It also causes mutations in a number of oncogenes and tumor suppressor genes. This can lead to uncontrolled keratinocyte mitosis and initiation of skin cancer upon exposure to UV light⁷.

Although cancer cannot be completely eradicated, there are steps you can take to reduce the risk. The actions that must be taken are minimizing exposure to sunlight, not using tobacco, using sunscreen to protect the skin from UV radiation, and removing skin cancer cells through biopsy⁸.

The people of South Sulawesi have empirically used Laruna leaves as traditional medicine. *Chromolaena odorata* (L) has long been used in traditional West African, Asian and Indonesian medicine for wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, antiimmunosuppressive and anticancer, as well as other uses^{10,11,12}.

The benefits of raw medicines are related to plant phytochemicals, which include phenolic compounds, carotenoids, antioxidant vitamins, terpenoids, steroids, indole, fiber, etc. The beneficial components thought to be responsible for lowering the risk of cancer are crude drugs¹⁴.

The bioactive compounds found in *Chromolaena odorata* have important therapeutic benefits in a number of clinical disorders^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40-trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihydrokaempferide, Apigenin-7,40-dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin-40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

MATERIALS AND METHODS:

Materials:

Chromolaena odorata collection is in Pangkep reGENCY, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, the 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions:

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitril, N-Buthanol, Ethanol.

Table 1: Extraction weight of *Chromolaena odorata*

| Sample | Solvent | Extract Weight (gr) | % Rendamen |
|---------------------------------|---------|---------------------|------------|
| <i>Chromolaena odorata</i> 5 kg | Metanol | 77.621 | 1.55 |

Table 2: Partition weight of *Chromolaena odorata*

| Sample | Solvent | Extract Weight |
|--|---------------|----------------|
| Methanol extract of <i>Chromolaena odorata</i> | n-Hexane | 68.45 |
| | Ethyl Acetate | 245.89 |
| | Acetonitrilic | 85.22 |
| | n-Buthanol | 147.97 |
| | Ethanol | 53.22 |

Cell Culture:

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10mg of each of extract were dissolved in 1mL of DMSO. The second stock solution (5mg/mL) was generated by mixing 0.5mL of this stock solution with 0.5mL of DMSO, which was mixed well. The third stock solution (2mg/mL) was mixed by 0,2mL of DMSO. To make the final serial concentrations, 1µL aliquots of the sample stock solutions in two-fold dilution (10, 5, 2mg/mL) were added to the corresponding well (the final volume became 100µL) to obtain the final concentrations (100, 50, 20µg/mL)

Cytotoxicity Investigation:

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1µL of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10µg/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = [1 - (A_{\text{test}} - A_{\text{blank}}) / (A_{\text{control}} - A_{\text{blank}})] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis:

Table 3: % inhibition

| Sample | Concentration | | |
|---------------|---------------|-------|-------|
| | 100 | 50 | 20 |
| Hexane | 16.56 | 5.54 | 8.36 |
| Ethyl Acetate | 48.00 | 5.12 | 12.20 |
| Acetonitrilic | 39.62 | 29.59 | -1.97 |
| Buthanol | 7.21 | 14.21 | 14.65 |
| Ethanol | -2.53 | 1.30 | 19.96 |

Table 4: Control positive (Etoposide)

| Control | Concentration | | | |
|-----------|---------------|-------|-------|-------|
| | 100 | 50 | 20 | 10 |
| Etoposide | 79.82 | 78.91 | 69.30 | 73.41 |

RESULTS:

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100µg/mL.




Figure 1: HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic, D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate ([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹⁶. IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISSCUSSION:

This research aims to identify the potential of natural ingredients by using HaCaT cells as a reliable in vitro model for evaluation.

The main cells of the human epidermis are called keratinocytes. Apart from their role in maintaining the

epidermal barrier, keratinocytes also have a role in the inflammatory response of the skin¹⁷ and the re-epithelialization stage of wound healing¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) addresses the problems observed with primary keratinocytes in the form of their need for growth supplementation, culture differentiation, donor-related variability, and limited lifespan. Many skin model investigations have included HaCaT cells without compromising keratinocyte function, cellular response, or differentiation potential¹⁹. There are many uses for monolayer culture of HaCaT cells, including analysis of in vitro wound healing and cellular toxicity. However, the actual structure of the epidermis cannot be completely replicated by monolayer culture methods. On the other hand, keratinocyte cell cultures that provide an air-liquid interface can produce a more accurate representation of the epidermis²⁰.

In the process of repairing the epidermis and maintaining the skin's immune defenses, keratinocytes have an active role by secreting growth factors, cytokines, and chemokines. For in vitro standardization studies, investigate the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, as well as their association with different culture states and different degrees of differentiation¹⁷.

The human line HaCaT cells are long-lived and spontaneously immortalized. They possess basal cell characteristics and respond significantly to two well-known in vitro prodifferentiating agents: increasing cell density and culture duration¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μ l of medium/well¹⁷.

The standard DMEM supplemented with 10% FBS, both of which contain enough calcium to trigger differentiation, was used to sustain HaCaT cells. It takes longer to dedifferentiate to the basal phenotype in low-calcium medium²².

According to statistical data, melanoma is the main cause of death due to skin cancer. Further research revealed that a patient's chances of survival depended on the stage of the cancer. Early detection and treatment of melanoma can increase the chances of recovery²³.

Topical medications are absorbed through the skin and pass through the skin layers to enter the systemic circulation. There are several topical agents that are often used including several corticosteroids, namely 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoid, Mechlorethamine, Carmustine, Bexarotene, Tazarotene photodynamic^{24,25}.

Plant phytochemicals, such as antioxidants, are thought to have beneficial effects over crude drugs. *Chromolaena odorata* leaf extract contains many antioxidants, flavonoids, phenolics, terpenoids²⁶. In addition, the results of the investigation show that the anti-mitotic activity of green silver nanoparticles made from *Chromolaena odorata* has a potential effect on cell division²⁷. The phytochemical profile of the phenolic extract of *Chromolaena odorata* has great potential in the discovery of new compounds with pharmacotherapeutic value²⁸. Research on *Chromolaena odorata* ethyl acetate extract at a concentration of 100 μ g/mL had the highest percentage, namely 48%. Meanwhile, *Helicteres isora* extract at a concentration of 300 μ g/ml has activity equivalent to the standard drug doxorubicin²⁹.

Additionally, when performing skin cancer testing using extracts such as ethosomal gel, which results in controlled release of curcumin, curcumin also has a longer half-life and maximum deposition in the deeper layers of the skin. Thus, it is able to completely eliminate melanoma cells³⁰. In addition, due to its pharmacological and therapeutic qualities, diosmin has the potential to be a flavonoid in the treatment of hepatocellular carcinoma, both in vivo and in vitro, in addition to its carcinogenic anticancer potential. Even so, further research into various types of cancer is still needed³¹.

In-depth studies of melanoma biology have had an influence on the consideration of melanoma as a therapy for one of the most resistant malignancies. There have been several options for treating metastatic melanoma in the last 30 years, so more extensive clinical studies have been conducted. These studies show that combination regimens, hormonal therapy, immunotherapy, single and combination agent chemotherapy, biochemotherapy, and targeted agent therapy do not always provide the best results²³.

CONCLUSION:

Based on cytotoxicity tests, *Chromolaena odorata* ethyl acetate extract has anti-skin cancer potential.

CONFLICT OF INTEREST:

There are no conflicts of interest declared by the authors of this study.

REFERENCES:

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashan D, Weller RB, Castellsagué X. Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)*. 2017; 7(Suppl 1):5-19. doi: 10.1007/s13555-016-0165-y
3. Akila Victor, Muhammad Rukunuddin Ghalib. Detection of skin cancer cells – a review. *Research J.Pharm and Tech*; 2017; 10(11):4093-4098. doi: 10.5958/0974-360X.2017.00742.9
4. Scott G. Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Invest Dermatol*. 2012; 132(4):1073-4. doi: 10.1038/jid.2012.20
5. Nasti TH, Timares L. MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*. 2015; 91(1):188-200. doi: 10.1111/php.12335
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan. Protein-protein interaction of mutated agouti signaling protein (asip) to melanocortin receptor 1 (mc1r) in melanoma skin cancer: an insilico study. *Research J. Pharm. and Tech* 2018; 11(9): 3913-3917. doi: 10.5958/0974-360X.2018.00718.7
7. Min Jeong Kim I, Seo Yeon Shin I, Nu Ri Song I, Sunoh Kim 2, Sang Ouk Sun I and Kyung Mok Park I. Bioassay-guided characterization, antioxidant, anti-melanogenic and anti-photoaging activities of pueraria thunbergiana l. leaf extracts in human epidermal keratinocytes (hacat) cells. *MDPI* 10. 2022; 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y. New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*. 2014; 2014:243452. doi: 10.1155/2014/243452
9. Qadir MI. Skin cancer: etiology and management. *Pak J Pharm Sci*. 2016; 29(3):999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med*. 2015; 301-306. <https://doi.org/10.1515/jcim-2014-0033>
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed. Biological activities of *chromolaena odorata*: A mechanistic review. *Elsevier*. 2021; <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S. Flavonoid glycosides from *chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull.* 59. 2011; 129-131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B. Pharmacological importance of *chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal Vol* 9. 2021; 8-11. Doi:10.47957/ijpda.v9i1.452
14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging, and antimicrobial potential of *chromolaena odorata* leaf extracts against pathogenic bacterium in wound infectionsa multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103-112. 20. <https://doi.org/10.1016/j.bcab.2018.05.014>
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J. Kaempferide, the most active among the four flavonoids isolated and characterized from *chromolaena odorata*, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5. 2015; 100912-100922. <https://doi.org/10.1039/C5RA19199H>
16. Mkhobongo B, Chandran R, Abrahamse H. In vitro photodynamic treatment modality for a375 melanoma cell line using a sulphonated aluminum phthalocyanine chloride-photosensitizer-gold nanoparticle conjugate. *Pharmaceutics*. 2022; 16;14(11):2474. doi: 10.3390/pharmaceutics14112474
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all. Hacat cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediator of Inflammation*. 2017; <https://doi.org/10.1155?2017/7435621>
18. Pastar I, Stojadinovic O, Yin NC, Ramirez H, Nusbaum AG, Sawaya A, Patel SB, Khalid L, Isseroff RR, Tomic-Canic M. Epithelialization in wound healing: a comprehensive review. *Adv Wound Care (New Rochelle)*. 2014; 3(7):445-464. doi: 10.1089/wound.2013.0473
19. Micallef L, Belaubre F, Pinon A, Jayat-Vignoles C, Delage C, Charveron M, Simon A. Effects of extracellular calcium on the growth-differentiation switch in immortalized keratinocyte hacat cells compared with normal human keratinocytes. *Exp Dermatol*. 2009; 18(2):143-51. doi: 10.1111/j.1600-0625.2008.00775.x
20. Pedrosa TDN, Catarino CM, Pennacchi PC, Assis SR, Gimenes F, Consolaro MEL, Barros SBM, Maria-Engler SS. A new reconstructed human epidermis for in vitro skin irritation testing. *Toxicol In Vitro*. 2017; 42:31-37. doi: 10.1016/j.tiv.2017.03.010
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R. Anti-inflammatory, and anti-proliferative properties of *Chromolaena odorata* leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants* 20. 2014; 359-371. doi:10.1080/10496475.2013.876698
22. Wilson VG. Growth and differentiation of hacat keratinocytes. *Methods Mol Biol*. 2014; 1195:33-41. doi: 10.1007/7651_2013_42
23. Tammineni Sreelatha, M. V. Subramanyam, M. N. Giri Prasad. A survey work on early detection methods of melanoma skin cancer. *Research J. Pharm. and Tech*. 2019; 12(5):2589-2596. doi: 10.5958/0974-360X.2019.00435.9
24. Micali G, Lacarubba F, Nasca MR, Ferraro S, Schwartz RA. Topical pharmacotherapy for skin cancer: part ii. Clinical applications. *Journal of the American Academy of Dermatology*. 2014; 70(6):979.e1-12; quiz 9912. doi: 10.1016/j.jaad.2013.12.037.
25. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kamya Goyal, Shammy Jindal. Treatment of skin cancer by topical drug delivery of nanoparticles: a review. *Research Journal of Pharmacy and Technology*. 2021; 14(10):5589-8. doi: 10.52711/0974-360X.2021.00973
26. Zainab, Nurlailah, Muhammad Ikhwan Rizki. Identification of active compound and antibacterial activity against gram-positive and gram-negative bacteria of *chromolaena odorata* leaf extract. *Research Journal of Pharmacy and Technology*. 2022; 15(10):4720-6. doi: 10.52711/0974-360X.2022.00793
27. Nishita Gogia, Anandhi D, Kanaga G, Revathi K. In vitro evaluation of anti-mitotic activity of green synthesized silver nanoparticles from ethanolic extract of *chromolaena odorata*, *caesalpinia coriaria* (bark and leaves) using *allium cepa* roots. *Research Journal of Pharmacy and Technology*. 2021; 14(8):4107-0. doi: 10.52711/0974-360X.2021.00711
28. Eze FN, Jayeoye TJ. *Chromolaena odorata* (Siam weed): A natural reservoir of bioactive compounds with potent anti-fibrillogenic, antioxidative, and cytocompatible properties. *Biomed Pharmacother*. 2021;141:111811. doi: 10.1016/j.biopha.2021.111811.
29. Madhulika Pradhan, S Sribhuaneswari, D Karthikeyan, Sunita Minz, Pavani Sure, Atul N Chandu, Umesh Mishra, K Kamalakannan, A Saravanankumar, T Sivakumar. In-vitro cytoprotection activity of *foeniculum vulgare* and *helicteres isora* in cultured human blood lymphocytes and antitumour activity against b16f10 melanoma cell line. *Research J. Pharm. and Tech*. 2018; 1(4):450-452
30. Radha Krishna Kollipara, Vyshnavi Tallapaneni, Bharat Kumar Reddy Sanapalli, Vinoth Kumar G, Veera Venkata Satyanarayana Reddy Karri. Curcumin loaded ethosomal vesicular drug delivery system for the treatment of melanoma skin cancer. *Research Journal of Pharmacy and Technology*. 2019; 12(4):1783-1792. doi: 10.5958/0974-360X.2019.00298.1
31. Perumal S, Langeswaran K. Diosmin anti-tumour efficacious against hepatocellular carcinoma. *Research J. Pharm. and Tech*. 2020; 13(4):1707-1714. doi: 10.5958/0974-360X.2020.00308.X
32. Ghadge Ojaswi, Nadar Divya, Parmar Digna. Melanoma and its drug targets. *Research J. Pharm. and Tech*. 2016; 9(5): 562-570. doi: 10.5958/0974-360X.2016.00107.4