

Manuskrip versi submit

Type of Manuscript :

Research

**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT)
Cells from Siam Weeds (*Chromolaena odorata* L.)**

ABSTRACT

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultra violet rays which are harmful to the skin and damage DNA. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. *Chromolaena odorata* was extracted with methanol, then partitioned with 5 dissolutions; n-Hexane, Ethyl Acetate, Acetonitrilic, n-Buthanol, Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocytes cells using the bioassay method. After incubation, the result obtained from microplate reader. Each extract was divided into three concentrations, it is 100, 50, 20 ($\mu\text{g}/\text{mL}$). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10 ($\mu\text{g}/\text{mL}$). After being analyzed with the results of the microplate reader, the IC₅₀ of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100 $\mu\text{g}/\text{mL}$. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μl of medium/well.

KEYWORDS: *Chromolaena odorata* L, Skin Cancer, HacaT cell line.

INTRODUCTION

Skin cancer is one of the most common forms of cancer worldwide, the impenetrable topical barrier is compromised slightly in pathologies such as skin cancer. Another disease that has seen growth in local therapeutic applications is skin cancer which is the most common cancer with one in every three cancer diagnoses being skin cancer according to WHO¹. Skin cancer is broadly classified into non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinomas and melanomas².

Cancer development usually occurs when DNA is damaged and our bodies are unable to repair this damage. These damaged cells begins to grow and divide uncontrollably. Such changes in the skin lead to skin cancer. When damage cell multiply, they form tumors².

Skin cancer develops in various layers of the skin. The tendency to develop skin cancer varies for different skin types. Occurrence varies according to skin color and skin type. In addition, heredity also contributes to the development of skin cancer. The most inherited type of cancer is Melanoma².

UV rays consist of UV-A and UV-B radiation. Although UV-B is considered more dangerous, UV-A also has side effects. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-A can also cause DNA damage by generating free radicals and damaging cell membranes which in turn causes skin cancer. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. UV-B skin also causes oxidative damage to the skin. UV-B radiation further interferes with regulatory proteins, anti-apoptotic relational proteins, proapoptotic proteins, transcription factors, protein kinase, inflammatory enzymes and growth factors which in turn disrupts the cell cycle³. This further causes reasonable mutations in various tumor suppressor genes and oncogenes. These effects together lead to uncontrolled keratinocytes mitosis and initiation of skin cancer upon UV exposure⁴.

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultra violet rays which are harmful to the skin and damage DNA. Before the body can recover, these deleterious mutations occur in the genes and the damage becomes irreparable. Moreover to environmental factors, the patient's genetic history can also be a cause of developing cancer⁴.

It is impossible to completely eradicate cancer, but precautions can be taken to prevent cancer. The following steps should be taken; there should be a reduction in sun exposure, use of tobacco should be avoided, sunscreen should be used to prevent exposure of UV rays, and skin cancer cells can also be removed by biopsy⁵.

The Laruna leaf has been used empirically as a traditional medicine by the people of South Sulawesi. Traditional use of *Chromolaena odorata* (L) as a source of medicine has been widely used in West African countries, Asian countries⁵, and Indonesia as wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, anti-immunosuppressive, anticancer and other activities^{6,7,8}.

Chromolaena odorata is has bioactive compounds with a significant therapeutics effects on some therapeutic effects on some pathological conditions^{9,10}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40 -trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihydrokaempferide, Apigenin-7,40 -dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin- 40-methyl ether, Aromadendrin-7,40-dimethyl ether^{8,11}.

MATERIALS AND METHODS

Materials:

Chromolaena odorata collection is located in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, The 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitrilic, N-Buthanol, Ethanol.

Table 1. Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77,621	1,55

Table 2. Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68,45
	Ethyl Acetate	245,89
	Acetonitrilic	85,22
	n-Buthanol	147,97
	Ethanol	53,22

Cell Culture

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10 mg of each of extract were dissolved in 1 mL of DMSO. The second stock solution (5 mg/mL) was generated by mixing 0.5 mL of this stock solution with 0.5 mL of DMSO, which was mixed well. The third stock solution (2 mg/mL) was mixed by 0,2 mL of DMSO. To make the final serial concentrations, 1 µL aliquots of the sample stock solutions in two-fold dilution (10, 5, 2 mg/mL) were added to the corresponding well (the final volume became 100 µL) to obtain the final concentrations (100, 50, 20 µg/mL)

Cytotoxicity Investigation

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1 µL of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10 µg/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = \left[1 - \frac{(A_{\text{test}} - A_{\text{blank}})}{(A_{\text{control}} - A_{\text{blank}})} \right] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis

Table 3. % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4. Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100 µg/mL.




Figure 1. HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic,
D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate ([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹². IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISCUSSION:

The aim of this study was to investigate the potential of natural products using HaCaT cells as a reliable in vitro model for evaluation.

Keratinocytes are the main cells of the human epidermis. In addition to providing an epidermal barrier function, keratinocytes play a role in the inflammatory response of the skin¹³, and the process of wound repair in the re-epithelialization step¹⁴. Establishment of an immortalized human keratinocyte cell line (HaCaT) eliminated the problems observed with primary keratinocytes such as their need for growth supplementation, culture differentiation, donor-related variability and limited lifespan. HaCaT cells have been used for many skin model studies without altering keratinocyte function, cellular response or differentiation capacity¹⁵. Monolayer HaCaT cell cultures are used for many applications such as cellular toxicity and in vitro analysis of wound healing. However monolayer culture systems cannot completely replicate the true epidermal structure. Whereas, the generation of air-liquid interface in keratinocytes cell cultures might result in a more realistic model of the epidermis¹⁶.

Keratinocytes are active in epidermal repair and skin immune defense through secretion of growth factors, cytokines, and chemokines. To standardizing in vitro studies, investigated the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ¹⁷, and in relation to different culture conditions and degrees of differentiation¹³.

HaCaT cells are a long-lived, spontaneously immortalized human line that exhibits basal cell properties and displays substantial changes in response to two well-established in vitro prodifferentiating agents: increased cell density and as a function of time in culture¹³.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 µl of medium/well¹³.

HaCaT cells were generally maintained in normal DMEM supplemented with 10% FBS, both of which contain sufficient calcium to induce differentiation. Dedifferentiation to the basal phenotype requires prolonged growth in low-calcium media¹⁸.

CONCLUSION:

Ethyl acetate extract of *Chromolaena odorata* has anti-skin cancer potential regarding to the cytotoxicity test.

CONFLICT OF INTEREST :

There are no conflicts of interest declared by the authors of this study.

REFERENCES

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashan D, Weller RB et al (2017) Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)* 7(Suppl 1):5–19
3. Singh M, Suman S, Shukla Y (2014) New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int* 2014
4. Min Jeong Kim 1, Seo Yeon Shin 1, Nu Ri Song 1, Sunoh Kim 2, Sang Ouk Sun 1 and Kyung Mok Park 1 (2022) Bioassay-Guided Characterization, Antioxidant, Anti-Melanogenic and Anti-Photoaging Activities of *Pueraria thunbergiana* L. Leaf Extracts in Human Epidermal Keratinocytes (HaCaT) Cells. *MDPI* 10, 2156. <https://doi.org/10.3390/pr10102156>
5. Qadir, M. Imran (2016) Skin Cancer: Etiology and Management. *J. Pharm. Sci*, Vol.29 No.3 pp.999-1003
6. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U., 2015. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med.* 12, 301–306. <https://doi.org/10.1515/jcim-2014-0033>.
7. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed (2021) Biological activities of *Chromolaena odorata*: A mechanistic review. Elsevier <https://doi.org/10.1016/j.sajb.2021.09.001>
8. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S., 2011. Flavonoid glycosides from *Chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull.* 59, 129–131. <https://doi.org/10.1248/cpb.59.129>.
9. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B., 2021. Pharmacological importance of *Chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal.* 8–11.
10. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A., 2018. Phytochemical screening, free radical scavenging and antimicrobial potential of *chromolaena odorata* leaf extracts against pathogenic bacterium in wound infectionsa multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. Vijayaraghavan, K, Rajkumar, J, Seyed, M.A., 20
11. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J., 2015. Kaempferide, the most active among the four flavonoids isolated and characterized from *Chromolaena odorata*, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5, 100912– 100922.
12. Bridgette Mkhobongo, Rahul Chandran * and Heidi Abrahamse (2022) In Vitro Photodynamic Treatment Modality for A375 Melanoma Cell Line Using a Sulphonated Aluminum Phthalocyanine Chloride-Photosensitizer-Gold Nanoparticle Conjugate. *MDPI* 2, 14, 2474. <https://doi.org/10.3390/pharmaceutics14112474>
13. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all (2017) HaCaT Cells as a Reliable in Vitro Differentiation Model to Dissect the Inflammatory/Repair Response of Human Keratinocytes. *Mediator of Inflammation*, <https://doi.org/10.1155/2017/7435621>.
14. Pastar, I., Stojadinovic, O., Yin, N.C., Ramirez, H., Nusbaum, A.G., Sawaya, A., Patel, S.B., Khalid, L., Isseroff, R.R. & Tomic-Canic, M. 2014. Ephetelialization in Wound Healing: a Comprehensive Review. *Advances in Wound Care*, 3(7): 445-464
15. Micallef, L., Belaubre, F., Pinon, A., Jayat-Vignoles, C., Delage, C., Charveron, M. & Simon, A. 2009. Effects of Extracellular Calcium on the Growth-Differentiation Switch in Immortalized Keratinocytes HaCaT Cells Compared with Normal Human Keratinocytes. *Experimental Dermatology*, 18(2): 143-151.
16. Pedrosa do Nascimento, Catarino C.M, Pennacchi, P.C, de Assis, S.R., Gimenes, F., Consolaro, M.E.L., de Moraes Barros, S.B & Maria-Engler, S.S. 2017. A new Reconstructed Human Epidermis for in Vitro Skin Irritation Testing. *Toxicology in Vitro*, 42:31-37
17. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R (2014) Anti-inflammatory and anti-proliferative properties of *Chromolaena odorata* leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants* 20, 359–371.
18. Wilson Van G (2013) Growth and Differentiation of HaCaT Keratinocytes. *Methods Molecular Biology* DOI 10.1007/7651_2013_42.

Certificate of Conflict of Interest:

The article entitled **Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)** is herewith submitted for publication in Research Journal of Pharmacy and Technology. It has not been published before, and it is not under consideration for publication in any other journal (s).

I/We certify that I/We have obtained written permission for the use of text, tables, and/or illustrations from any copyrighted source(s), and I/We declare no conflict of interest.

Signature of author(s)

Name(s) and designation



Karlina Amir Tahir

UIN Alauddin Makassar

Transfer of copyright agreement:

The article entitled **Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata L.*)** is herewith submitted for publication in Research Journal of Pharmacy and Technology. It has not been published before, and it is not under consideration for publication in any other journal (s). It contains no matter that is scandalous, obscene, libelous, or otherwise contrary to law. I/We agree that copies made under these circumstances will continue to carry the copyright notice that appeared in the original published work. I/We certify that I/We have obtained written permission for the use of text, tables, and/or illustrations from any copyrighted source(s), and I/We agree to supply such written permission(s) to Research Journal of Pharmacy and Technology upon request.

When the article is accepted for publication, I/We, as author/authors, hereby agree to transfer(s), assign(s), or otherwise convey(s) all copyright ownership, including any and all rights incidental thereto, exclusively to the Journal, in the event that such work is published by the Journal. The Journal shall own the work, including 1) copyright; 2) the right to grant permission to republish the article in whole or in part, with or without fee; 3) the right to produce preprints or reprints and translate into languages other than English for sale or free distribution; and 4) the right to republish the work in a collection of articles in any other mechanical or electronic format.

The article will be published under the terms of the latest Creative Commons Attribution-NonCommercial- ShareAlike License, unless the journal notifies the author otherwise in writing.

Signature of author(s)

Name(s) and designation



Karlina Amir Tahir

UIN Alauddin Makassar

KORESPONDENSI JURNAL

Submit tanggal 6 mei 2023



A AND V PUBLICATIONS

Report Generated On: 10 Sep, 2024 01:57 PM

Article's Basic Information:

Paper ID : 2356034457024539

Paper Title : **Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (Chromolaena odorata L.)**

Authors : **Karlina Amir Tahir;Erwin Hafid;Muh. Fitrah;Ahmad Lalo;Nurul Fadilah;Syamsuri Syakri;Syatirah Jalaluddin;Katsuyoshi Matsunami**

Author's Email : **karlina.amir@uin-alauddin.ac.id;erwin.hafid@uin-alauddin.ac.id;muhfitrah@uin-alauddin.ac.id;ahmadlalo_pom@yahoo.co.id;syamsurisyakri@gmail.com;syatirahjalaluddin@gmail.com;**

Submitted to Journal : **Research Journal of Pharmacy and Technology**

Submitted By : **karlina amir tahir (nilalfath@gmail.com)**

Date of Submission : **06 May, 2023**

 **Resubmission of Article**

Commented On: 23 Oct, 2023 06:05 PM

Dear Author,
Pl Resubmit the article after making following corrections.

- The Ref no. are not mentioned serially in text.
- The Ref no. are not match with text reference.

Thanks
Editor

 **Resubmission of Article**

Commented On: 13 Sep, 2023 04:53 PM

Dear Author,
Pl Resubmit the article after making corrections suggested by Editor.

The Ref no. are not as per instruction to author.
(Model Journal reference) Template

1. Ballesteros B, Tobias G, Shao L, Pellicer E, Nogués J, Mendoza E, Green ML et al Steam Purification for the Removal of Graphitic Shells Coating Catalytic Particles and the Shortening of Single-Walled Carbon Nanotubes. *Small*. 2008; 4(9):1501-6. doi.org/10.1021/ja061680u

2. Singh B, Lohan S, Sandhu PS, Jain A, Mehta SK. Functionalized carbon nanotubes and their promising applications in therapeutics and diagnostics. In *Nanobiomaterials in Medical Imaging 2016*; 455-478, William Andrew Publishing. <https://doi.org/10.1021/nn700040t>

Thanks
Editor

 **Resubmission of Article**

Commented On: 13 Sep, 2023 04:53 PM

Dear Author,
Pl Resubmit the article after making corrections suggested by Editor.

The Ref no. are not as per instruction to author.
(Model Journal reference) Template

1. Ballesteros B, Tobias G, Shao L, Pellicer E, Nogués J, Mendoza E, Green ML et al Steam Purification for the Removal of Graphitic Shells Coating Catalytic Particles and the Shortening of Single-Walled Carbon Nanotubes. *Small*. 2008; 4(9):1501-6. doi.org/10.1021/ja061680u

2. Singh B, Lohan S, Sandhu PS, Jain A, Mehta SK. Functionalized carbon nanotubes and their promising applications in therapeutics and diagnostics. In *Nanobiomaterials in Medical Imaging 2016*; 455-478, William Andrew Publishing. <https://doi.org/10.1021/nn700040t>

Thanks
Editor

 **Editorial Board Team**

Commented On: 10 May, 2023 11:32 AM

In order to improve the quality of the paper you just go through the below link and cite atleast 10 articles from the list, that have similar works as of yours. The link is http://www.anvpublication.org/Universal_Search.aspx?What=CitationList &tmp_PID=2356034457024539.

Draft artikel revisi 1

Type of Manuscript :

Research

**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT)
Cells from Siam Weeds (*Chromolaena odorata* L.)**

ABSTRACT

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultra violet rays which are harmful to the skin and damage DNA. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. *Chromolaena odorata* was extracted with methanol, then partitioned with 5 dissolutions; n-Hexane, Ethyl Acetate, Acetonitrilic, n-Buthanol, Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocytes cells using the bioassay method. After incubation, the result obtained from microplate reader. Each extract was divided into three concentrations, it is 100, 50, 20 ($\mu\text{g}/\text{mL}$). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10 ($\mu\text{g}/\text{mL}$). After being analyzed with the results of the microplate reader, the IC₅₀ of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100 $\mu\text{g}/\text{mL}$. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μl of medium/well.

KEYWORDS: *Chromolaena odorata* L, Skin Cancer, HacaT cell line.

INTRODUCTION

Skin cancer is one of the most common forms of cancer worldwide, the impenetrable topical barrier is compromised slightly in pathologies such as skin cancer. Another disease that has seen growth in local therapeutic applications is skin cancer which is the most common cancer with one in every three cancer diagnoses being skin cancer according to WHO¹. Skin cancer is broadly classified into non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinomas and melanomas².

Cancer development usually occurs when DNA is damaged and our bodies are unable to repair this damage. These damaged cells begins to grow and divide uncontrollably. Such changes in the skin lead to skin cancer. When damage cell multiply, they form tumors².

Skin cancer develops in various layers of the skin. The tendency to develop skin cancer varies for different skin types. Occurrence varies according to skin color and skin type. In addition, heredity also contributes to the development of skin cancer. The most inherited type of cancer is Melanoma².

Skin cancer can be identified through symptoms. Early identification and early detection of medicine and cure. Sometimes symptoms appear at an advanced stage in other types of cancer. Basal cell cancer appears as a lump on the skin anywhere on the body exposed to sunlight. They can be brown bumps or white bumps on our face, hands, neck and other exposed body parts. Squamous cell carcinoma on the other hand can appear as ulcerated lumps and patches on the skin whereas melanoma can cause unusual growths on the body especially in existing moles. It can be anywhere in our body³.

The American Cancer Society (ACS) has reported that, there were 87,110 newly diagnosed melanomas, also reported that 9,730 people had an estimated melanoma death in 2017. Skin pigmentation is caused by accumulation of melanin granules which are synthesized in melanocytes and transferred to keratinocytes⁴. Keratinocytes secrete cytokines and chemokines on exposure to UV light which affect melanin production⁵. Melanoma develops at any part of the skin due to defects in skin pigmentation, but is most susceptible to the trunk in men and the legs in women⁶.

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultra violet rays which are harmful to the skin and damage DNA. Before the body can recover, these deleterious mutations occur in the genes and the damage becomes irreparable. Moreover to environmental factors, the patient's genetic history can also be a cause of developing cancer⁷.

UV rays consist of UV-A and UV-B radiation. Although UV-B is considered more dangerous, UV-A also has side effects. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-A can also cause DNA damage by generating free radicals and damaging cell membranes which in turn causes skin cancer. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. UV-B skin also causes oxidative damage to the skin. UV-B radiation further interferes with regulatory proteins, anti-apoptotic relational proteins, proapoptotic proteins, transcription factors, protein kinase, inflammatory enzymes and growth factors which in turn disrupts the cell cycle⁸. This further causes reasonable mutations in various tumor suppressor genes and oncogenes. These effects together lead to uncontrolled keratinocytes mitosis and initiation of skin cancer upon UV exposure⁷.

It is impossible to completely eradicate cancer, but precautions can be taken to prevent cancer. The following steps should be taken; there should be a reduction in sun exposure, use of tobacco should be avoided, sunscreen should be used to prevent exposure of UV rays, and skin cancer cells can also be removed by biopsy⁸.

The Laruna leaf has been used empirically as a traditional medicine by the people of South Sulawesi. Traditional use of *Chromolaena odorata* (L) as a source of medicine has been widely used in West African countries, Asian countries⁹, and Indonesia as wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, anti-immunosuppressive, anticancer and other activities^{10,11,12}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals (various factors in plant foods), such as carotenoids, antioxidant vitamins, phenolic compounds, terpenoids, steroids, indole, and fiber, etc¹³. Crude drugs are the effective elements considered responsible for reducing cancer risk¹⁴.

Chromolaena odorata is has bioactive compounds with a significant therapeutics effects on some therapeutic effects on some pathological conditions^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40 -trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihydrokaempferide, Apigenin-7,40 -dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin- 40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

MATERIALS AND METHODS

Materials:

Chromolaena odorata collection is located in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, The 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitrilic, N-Buthanol, Ethanol.

Table 1. Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77,621	1,55

Table 2. Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68,45
	Ethyl Acetate	245,89
	Acetonitrilic	85,22
	n-Buthanol	147,97
	Ethanol	53,22

Cell Culture

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10 mg of each of extract were dissolved in 1 mL of DMSO. The second stock solution (5 mg/mL) was generated by mixing 0.5 mL of this stock solution with 0.5 mL of DMSO, which was mixed well. The third stock solution (2 mg/mL) was mixed by 0,2 mL of DMSO. To make the final serial concentrations, 1 μ L aliquots of the sample stock solutions in two-fold dilution (10, 5, 2 mg/mL) were added to the corresponding well (the final volume became 100 μ L) to obtain the final concentrations (100, 50, 20 μ g/mL)

Cytotoxicity Investigation

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1 μ L of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10 μ g/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = \left[\frac{1 - (A_{\text{test}} - A_{\text{blank}})}{(A_{\text{control}} - A_{\text{blank}})} \right] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis

Table 3. % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4. Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100 µg/mL.




Figure 1. HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic,
D = n-Butanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate ([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹². IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISSCUSSION:

The aim of this study was to investigate the potential of natural products using HaCaT cells as a reliable in vitro model for evaluation.

Keratinocytes are the main cells of the human epidermis. In addition to providing an epidermal barrier function, keratinocytes play a role in the inflammatory response of the skin¹⁷, and the process of wound repair in the re-epithelialization step¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) eliminated the problems observed with primary keratinocytes such as their need for growth supplementation, culture differentiation, donor-related variability and limited lifespan. HaCaT cells have been used for many skin model studies without altering keratinocyte function, cellular response or differentiation capacity¹⁹. Monolayer HaCaT cell cultures are used for many applications such as cellular toxicity and in vitro analysis of wound healing. However monolayer culture systems cannot completely replicate the true epidermal structure. Whereas, the generation of air-liquid interface in keratinocytes cell cultures might result in a more realistic model of the epidermis²⁰.

Keratinocytes are active in epidermal repair and skin immune defense through secretion of growth factors, cytokines, and chemokines. To standardizing in vitro studies, investigated the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, and in relation to different culture conditions and degrees of differentiation¹⁷.

HaCaT cells are a long-lived, spontaneously immortalized human line that exhibits basal cell properties and displays substantial changes in response to two well-established in vitro prodifferentiating agents: increased cell density and as a function of time in culture¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 µl of medium/well¹⁷.

HaCaT cells were generally maintained in normal DMEM supplemented with 10% FBS, both of which contain sufficient calcium to induce differentiation. Dedifferentiation to the basal phenotype requires prolonged growth in low-calcium media²².

Statistical evidence has revealed that the majority of deaths from skin cancer are the result of melanoma. Further investigations have shown that the survival rate in patients depends on the stage of the cancer; Early detection and intervention of melanoma involves a higher chance of cure.

Topical therapies are delivered through skin & they reach the systemic circulation by crossing the skin layers.^{41,42} There are some commonly used topical agent which includes some corticosteroids is given as: 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoids, Mechlorethamine, Carmustine, Bexarotene, Tazarotene photodynamic

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals, such as antioxidant. *Chromolaena odorata* leaf extract contain various antioxidant, terpenoid, phenolic and flavonoid compounds. In addition, the anti-mitotic activity of the results of this investigation shows that silver green nanoparticles synthesized from *C. odorata* have a potential effect on cell division.

CONCLUSION:

Ethyl acetate extract of *Chromolaena odorata* has anti-skin cancer potential regarding to the cytotoxicity test.

CONFLICT OF INTEREST :

There are no conflicts of interest declared by the authors of this study.

REFERENCES

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashan D, Weller RB et al (2017) Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)* 7(Suppl 1):5–19
3. Akila Victor, Muhammad Rukunuddin Ghalib (2017), Detection of Skin Cancer Cells – A Review. *Research J.Pharm and Tech*; 10(11):4093-4098. DOI: 10.5958/0974-360X.2017.00742.9
4. Scott G (2012). Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Investig Dermatol*; 132:1073–1074
5. Nasti TH, Timares L (2015). MC1R, Eumelanin and Pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*;91:188-200
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan (2018). Protein-Protein Interaction of Mutated Agouti Signaling Protein (ASIP) to Melanocortin Receptor 1 (MC1R) in Melanoma Skin Cancer: An Insilico Study. *Research J. Pharm. and Tech*; 11(9): 3913-3917
7. Min Jeong Kim 1, Seo Yeon Shin 1, Nu Ri Song 1, Sunoh Kim 2 , Sang Ouk Sun 1 and Kyung Mok Park 1 (2022) Bioassay-Guided Characterization, Antioxidant, Anti-Melanogenic and Anti-Photoaging Activities of Pueraria thunbergiana L. Leaf Extracts in Human Epidermal Keratinocytes (HaCaT) Cells. *MDPI* 10, 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y (2014) New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*
9. Qadir, M. Imran (2016) Skin Cancer: Etiology and Management. *J. Pharm. Sci*, Vol.29 No.3 pp.999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U (2015). Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med.* 12, 301–306. <https://doi.org/10.1515/jcim-2014-0033>.
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed (2021) Biological activities of *Chromolaena odorata*: A mechanistic review. Elsevier <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S (2011). Flavonoid glycosides from *Chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull.* 59, 129–131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B., (2021). Pharmacological importance of *Chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal.* 8–11
14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging and antimicrobial potential of *chromolaena odorata* leaf extracts against pathogenic bacterium in wound infectionsa multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. Vijayaraghavan, K, Rajkumar, J., Seyed, M.A., 20
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J (2015). Kaempferide, the most active among the four flavonoids isolated and characterized from *Chromolaena odorata*, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5, 100912– 100922
16. Bridgette Mkhobongo, Rahul Chandran and Heidi Abrahamse (2022) In Vitro Photodynamic Treatment Modality for A375 Melanoma Cell Line Using a Sulphonated Aluminum Phthalocyanine Chloride-Photosensitizer-Gold Nanoparticle Conjugate. *MDPI* 2, 14, 2474. <https://doi.org/10.3390/pharmaceutics14112474>
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all (2017) HaCaT Cells as a Reliable in Vitro Differentiation Model to Dissect the Inflammatory/Repair Response of Human Keratinocytes. *Mediator of Inflammation*, <https://doi.org/10.1155?2017/7435621>
18. Pastar, I., Stojadinovic, O., Yin, N.C, Ramirez, H., Nusbaum, A.G., Sawaya, A., Patel, S.B., Khalid, L., Isseroff, R.R. & Tomic-Canic, M (2014). Epithelialization in Wound Healing: a Comprehensive Review. *Advances in Wound Care*, 3(7): 445-464
19. Micallef, L., Belaubre, F., Pinon, A., Jayat-Vignoles, C., Delage, C., Charveron, M. & Simon, A (2009). Effects of Extracellular Calcium on the Growth-Differentiation Switch in Immortalized Keratinocytes HaCaT Cells Compared with Normal Human Keratinocytes. *Experimental Dermatology*, 18(2): 143-151
20. Pedrosa do Nascimento, Catarino C.M, Pennacchi, P.C, de Assis, S.R., Gimenes, F., Consolaro, M.E.L., de Moraes Barros, S.B & Maria-Engler, S.S (2017). A new Reconstructed Human Epidermis for in Vitro Skin Irritation Testing. *Toxicology in Vitro*, 42:31-37
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R (2014) Anti-inflammatory and anti-proliferative properties of *Chromolaena odorata* leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants* 20, 359–371
22. Wilson Van G (2013) Growth and Differentiation of HaCaT Keratinocytes. *Methods Molecular Biology* DOI 10.1007/7651_2013_42

23. Tammineni Sreelatha, M. V. Subramanyam, M. N. Giri Prasad (2019). A Survey work on Early Detection methods of Melanoma Skin Cancer. *Research J. Pharm. and Tech.* 12(5):2589-2596
24. McGillis ST, Fein H (2004). Topical treatment strategies for non-melanoma skin cancer & precursor lesions. In *Seminars in cutaneous medicine & surgery*. Vol. 23, No. 3, pp. 174-183
25. Micali G, Lacarrubba F, Nasca MR, Ferraro S, Schwartz RA (2014). Topical pharmacotherapy for skin cancer: part II. Clinical applications. *Journal of the American Academy of Dermatology*. 70(6):979-e1)
26. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kanya Goyal, Shammy Jindal (2021). Treatment of Skin Cancer by Topical Drug Delivery of Nanoparticles: A Review. *Research Journal of Pharmacy and Technology*. 14(10):5589-8
27. Zainab, Nurlailah, Muhammad Ikhwan Rizki (2022). Identification of active compound and Antibacterial activity against gram-positive and gram-negative bacteria of *Chromolaena odorata* leaf extract. *Research Journal of Pharmacy and Technology*. 15(10):4720-6.)
28. Nishita Gogia, Anandhi D, Kanaga G, Revathi K (2021). In Vitro Evaluation of Anti-Mitotic Activity of Green Synthesized Silver Nanoparticles from Ethanolic Extract of *Chromolaena odorata*, *Caesalpinia coriaria* (Bark and Leaves) using *Allium cepa* Roots. *Research Journal of Pharmacy and Technology*. 14(8):4107-0

Draft artikel versi revisi ke 2

Type of Manuscript:

Research

**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT)
Cells from Siam Weeds (*Chromolaena odorata* L.)**

ABSTRACT

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. *Chromolaena odorata* was extracted with methanol, then partitioned with 5 dissolutions: n-Hexane, Ethyl Acetate, Acetonitrilic, n-Buthanol, Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocytes cells using the bioassay method. After incubation, the result obtained from microplate reader. Each extract was divided into three concentrations, it is 100, 50, 20 ($\mu\text{g}/\text{mL}$). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10 ($\mu\text{g}/\text{mL}$). After being analyzed with the results of the microplate reader, the IC₅₀ of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100 $\mu\text{g}/\text{mL}$. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μl of medium/well.

KEYWORDS: *Chromolaena odorata* L., Skin Cancer, HacaT cell line.

INTRODUCTION

Skin cancer is one of the most common forms of cancer worldwide, the impenetrable topical barrier is compromised slightly in pathologies such as skin cancer. Another disease that has seen growth in local therapeutic applications is skin cancer which is the most common cancer with one in every three cancer diagnoses being skin cancer according to World Health Organization (WHO)¹. Skin cancer is broadly classified into non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinomas and melanomas².

Cancer development usually occurs when DNA is damaged, and our bodies are unable to repair this damage. These damaged cells begin to grow and divide uncontrollably. Such changes in the skin lead to skin cancer. When damage cell multiply, they form tumors².

Skin cancer develops in various layers of the skin. The tendency to develop skin cancer varies for different skin types. Occurrence varies according to skin color and skin type. In addition, heredity also contributes to the development of skin cancer. The most inherited type of cancer is Melanoma².

Skin cancer can be identified through symptoms. Early identification and early detection of medicine and cure. Sometimes symptoms appear at an advanced stage in other types of cancer. Basal cell cancer appears as a lump on the skin anywhere on the body exposed to sunlight. They can be brown bumps or white bumps on our face, hands, neck, and other exposed body parts. Squamous cell carcinoma on the other hand can appear as ulcerated lumps and patches on the skin whereas melanoma can cause unusual growths on the body especially in existing moles. It can be anywhere in our body³.

The American Cancer Society (ACS) has reported that, there were 87,110 newly diagnosed melanomas, also reported that 9,730 people had an estimated melanoma death in 2017. Skin pigmentation is caused by accumulation of melanin granules which are synthesized in melanocytes and transferred to keratinocytes⁴. Keratinocytes secrete cytokines and chemokines on exposure to UV light which affect melanin production⁵. Melanoma develops at any part of the skin due to defects in skin pigmentation but is most susceptible to the trunk in men and the legs in women⁶.

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. Before the body can recover, these deleterious mutations occur in the genes and the damage becomes irreparable. Moreover, to environmental factors, the patient's genetic history can also be a cause of developing cancer⁷.

UV rays consist of UV-A and UV-B radiation. Although UV-B is considered more dangerous, UV-A also has side effects. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-A can also cause DNA damage by generating free radicals and damaging cell membranes which in turn causes skin cancer. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. UV-B skin also causes oxidative damage to the skin. UV-B radiation further interferes with regulatory proteins, anti-apoptotic relational proteins, proapoptotic proteins, transcription factors, protein kinase, inflammatory enzymes, and growth factors which in turn disrupts the cell cycle⁸. This further causes reasonable mutations in various tumor suppressor genes and oncogenes. These effects together lead to uncontrolled keratinocytes mitosis and initiation of skin cancer upon UV exposure⁷.

It is impossible to completely eradicate cancer, but precautions can be taken to prevent cancer. The following steps should be taken; there should be a reduction in sun exposure, use of tobacco should be avoided, sunscreen should be used to prevent exposure of UV rays, and skin cancer cells can also be removed by biopsy⁸.

The Laruna leaf has been used empirically as a traditional medicine by the people of South Sulawesi. Traditional use of *Chromolaena odorata* (L) as a source of medicine has been widely used in West African countries, Asian countries⁹, and Indonesia as wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, anti-immunosuppressive, anticancer, and other activities^{10,11,12}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals (various factors in plant foods), such as carotenoids, antioxidant vitamins, phenolic compounds, terpenoids, steroids, indole, and fiber, etc¹³. Crude drugs are the effective elements considered responsible for reducing cancer risk¹⁴.

Chromolaena odorata has bioactive compounds with a significant therapeutics' effects on some therapeutic effects on some pathological conditions^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40 -trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihydrokaempferide, Apigenin-7,40 -dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin- 40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

MATERIALS AND METHODS

Materials:

Chromolaena odorata collection is in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, the 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitrilic, N-Buthanol, Ethanol.

Table 1. Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77.621	1.55

Table 2. Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68.45
	Ethyl Acetate	245.89
	Acetonitrilic	85.22
	n-Buthanol	147.97
	Ethanol	53.22

Cell Culture

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10 mg of each of extract were dissolved in 1 mL of DMSO. The second stock solution (5 mg/mL) was generated by mixing 0.5 mL of this stock solution with 0.5 mL of DMSO, which was mixed well. The third stock solution (2 mg/mL) was mixed by 0,2 mL of DMSO. To make the final serial concentrations, 1 μ L aliquots of the sample stock solutions in two-fold dilution (10, 5, 2 mg/mL) were added to the corresponding well (the final volume became 100 μ L) to obtain the final concentrations (100, 50, 20 μ g/mL)

Cytotoxicity Investigation

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1 μ L of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10 μ g/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = \left[\frac{1 - (A_{\text{test}} - A_{\text{blank}})}{(A_{\text{control}} - A_{\text{blank}})} \right] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis

Table 3. % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4. Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100 µg/mL.




Figure 1. HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic, D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate ([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹⁶. IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISCUSSION:

The aim of this study was to investigate the potential of natural products using HaCaT cells as a reliable in vitro model for evaluation.

Keratinocytes are the main cells of the human epidermis. In addition to providing an epidermal barrier function, keratinocytes play a role in the inflammatory response of the skin¹⁷, and the process of wound repair in the re-epithelialization step¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) eliminated the problems observed with primary keratinocytes such as their need for growth supplementation, culture differentiation, donor-related variability, and limited lifespan. HaCaT cells have been used for many skin model studies without altering keratinocyte function, cellular response, or differentiation capacity¹⁹. Monolayer HaCaT cell cultures are used for many applications such as cellular toxicity and in vitro analysis of wound healing. However monolayer culture systems cannot completely replicate the true epidermal structure. Whereas the generation of air-liquid interface in keratinocytes cell cultures might result in a more realistic model of the epidermis²⁰.

Keratinocytes are active in epidermal repair and skin immune defense through secretion of growth factors, cytokines, and chemokines. To standardizing in vitro studies, investigated the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, and in relation to different culture conditions and degrees of differentiation¹⁷.

HaCaT cells are a long-lived, spontaneously immortalized human line that exhibits basal cell properties and displays substantial changes in response to two well-established in vitro prodifferentiating agents: increased cell density and as a function of time in culture¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 µl of medium/well¹⁷.

HaCaT cells were generally maintained in normal DMEM supplemented with 10% FBS, both of which contain sufficient calcium to induce differentiation. Dedifferentiation to the basal phenotype requires prolonged growth in low calcium media²².

Statistical evidence has revealed that most deaths from skin cancer are the result of melanoma. Further investigations have shown that the survival rate in patients depends on the stage of the cancer; Early detection and intervention of melanoma involves a higher chance of cure²³.

Topical therapies are delivered through skin & they reach the systemic circulation by crossing the skin layers. There are some commonly used topical agent which includes some corticosteroids is given as: 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoids, Mechlorethamine, Carmustine, Bexarotene, Tazarotene photodynamic^{24, 25}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals, such as antioxidant. *Chromolaena odorata* leaf extract contain various antioxidant, terpenoid, phenolic and

flavonoid compounds²⁶. In addition, the anti-mitotic activity of the results of this investigation shows that silver green nanoparticles synthesized from *Chromolaena odorata* have a potential effect on cell division²⁷. The phytochemical profile of the phenolic extract of *Chromolaena odorata* has great potential for the identification of new compounds with pharmacotherapeutic value²⁸. As a comparison to *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48%, research on *Helicteres isora* extract at a concentration of 300 µg/ml had an activity equivalent to the activity of the standard drug doxorubicin²⁹.

In addition, regarding skin cancer testing with various extracts, for example, Ethosomal gel which produces controlled release of curcumin, Curcumin persists in deeper layers of the skin for a longer time, as well as maximum deposition. Thus, eliminating melanoma cells completely³⁰. Furthermore, on the carcinogenic anti-cancer potential of diosmin in vivo and in vitro, it can be a flavonoid in the treatment of Hepatocellular Carcinoma due to its therapeutic and pharmacological properties. Although this research requires further research on other types of cancer³¹.

In-depth study of melanoma biology has been influential in considering melanoma as a therapy for one of the most resistant malignancies. In the last 30 years, there have been few options for treating metastatic melanoma, so some more thorough clinical studies have been conducted, such as hormonal therapy, immunotherapy, single and combination agent chemotherapy, biochemotherapy, targeted agent therapy, and combination regimens, show less than optimal results³².

CONCLUSION:

Ethyl acetate extract of *Chromolaena odorata* has anti-skin cancer potential regarding to the cytotoxicity test.

CONFLICT OF INTEREST:


There are no conflicts of interest declared by the authors of this study.

REFERENCES

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashed D, Weller RB, Castellsagué X. Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)*. 2017; 7(Suppl 1):5-19. doi: 10.1007/s13555-016-0165-y
3. Akila Victor, Muhammad Rukunuddin Ghalib. Detection of skin cancer cells – a review. *Research J.Pharm and Tech*; 2017; 10(11):4093-4098. doi: 10.5958/0974-360X.2017.00742.9
4. Scott G. Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Invest Dermatol*. 2012; 132(4):1073-4. doi: 10.1038/jid.2012.20
5. Nasti TH, Timares L. MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*. 2015; 91(1):188-200. doi: 10.1111/php.12335
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan. Protein-protein interaction of mutated agouti signaling protein (asip) to melanocortin receptor 1 (mc1r) in melanoma skin cancer: an insilico study. *Research J. Pharm. and Tech* 2018; 11(9): 3913-3917. doi: 10.5958/0974-360X.2018.00718.7
7. Min Jeong Kim 1, Seo Yeon Shin 1, Nu Ri Song 1, Sunoh Kim 2, Sang Ouk Sun 1 and Kyung Mok Park 1. Bioassay-guided characterization, antioxidant, anti-melanogenic and anti-photoaging activities of pueraria thunbergiana l. leaf extracts in human epidermal keratinocytes (hacat) cells. *MDPI* 10. 2022; 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y. New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*. 2014; 2014:243452. doi: 10.1155/2014/243452
9. Qadir MI. Skin cancer: etiology and management. *Pak J Pharm Sci*. 2016; 29(3):999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med*. 12. 2015; 301–306. <https://doi.org/10.1515/jcim-2014-0033>
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed. Biological activities of *chromolaena odorata*: A mechanistic review. *Elsevier*. 2021; <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S. Flavonoid glycosides from *chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull*. 59. 2011; 129–131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B. Pharmacological importance of *chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal Vol* 9. 2021; 8–11. Doi:10.47957/ijpda.v9i1.452

14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging, and antimicrobial potential of chromolaena odorata leaf extracts against pathogenic bacterium in wound infections a multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. 20. <https://doi.org/10.1016/j.bcab.2018.05.014>
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J. Kaempferide, the most active among the four flavonoids isolated and characterized from chromolaena odorata, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5. 2015; 100912– 100922. <https://doi.org/10.1039/C5RA19199H>
16. Mkhobongo B, Chandran R, Abrahamse H. In vitro photodynamic treatment modality for a375 melanoma cell line using a sulfonated aluminum phthalocyanine chloride-photosensitizer-gold nanoparticle conjugate. *Pharmaceutics.* 2022; 16;14(11):2474. doi: 10.3390/pharmaceutics14112474
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all. Hacat cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediator of Inflammation.* 2017; <https://doi.org/10.1155/2017/7435621>
18. Pastar I, Stojadinovic O, Yin NC, Ramirez H, Nusbaum AG, Sawaya A, Patel SB, Khalid L, Isseroff RR, Tomic-Canic M. Epithelialization in wound healing: a comprehensive review. *Adv Wound Care (New Rochelle).* 2014; 3(7):445-464. doi: 10.1089/wound.2013.0473
19. Micallef L, Belaube F, Pinon A, Jayat-Vignoles C, Delage C, Charveron M, Simon A. Effects of extracellular calcium on the growth-differentiation switch in immortalized keratinocyte hacat cells compared with normal human keratinocytes. *Exp Dermatol.* 2009; 18(2):143-51. doi: 10.1111/j.1600-0625.2008.00775.x
20. Pedrosa TDN, Catarino CM, Pennacchi PC, Assis SR, Gimenes F, Consolaro MEL, Barros SBM, Maria-Engler SS. A new reconstructed human epidermis for in vitro skin irritation testing. *Toxicol In Vitro.* 2017; 42:31-37. doi: 10.1016/j.tiv.2017.03.010
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R. Anti-inflammatory, and anti-proliferative properties of Chromolaena odorata leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants.* 20. 2014; 359–371. doi:10.1080/10496475.2013.876698
22. Wilson VG. Growth and differentiation of hacat keratinocytes. *Methods Mol Biol.* 2014; 1195:33-41. doi: 10.1007/7651_2013_42
23. Tammini Sreelatha, M. V. Subramanyam, M. N. Giri Prasad. A survey work on early detection methods of melanoma skin cancer. *Research J. Pharm. and Tech.* 2019; 12(5):2589-2596. doi: 10.5958/0974-360X.2019.00435.9
24. Micali G, Lacarubba F, Nasca MR, Ferraro S, Schwartz RA. Topical pharmacotherapy for skin cancer: part ii. Clinical applications. *Journal of the American Academy of Dermatology.* 2014; 70(6):979.e1-12; quiz 9912. doi: 10.1016/j.jaad.2013.12.037.
25. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kamy Goyal, Shammy Jindal. Treatment of skin cancer by topical drug delivery of nanoparticles: a review. *Research Journal of Pharmacy and Technology.* 2021; 14(10):5589-8. doi: 10.52711/0974-360X.2021.00973
26. Zainab, Nurlailah, Muhammad Ikhwan Rizki. Identification of active compound and antibacterial activity against gram-positive and gram-negative bacteria of chromolaena odorata leaf extract. *Research Journal of Pharmacy and Technology.* 2022; 15(10):4720-6. doi: 10.52711/0974-360X.2022.00793
27. Nishita Gogia, Anandhi D, Kanaga G, Revathi K. In vitro evaluation of anti-mitotic activity of green synthesized silver nanoparticles from ethanolic extract of chromolaena odorata, caesalpinia coriaria (bark and leaves) using allium cepa roots. *Research Journal of Pharmacy and Technology.* 2021; 14(8):4107-0. doi: 10.52711/0974-360X.2021.00711
28. Eze FN, Jayeoye TJ. Chromolaena odorata (Siam weed): A natural reservoir of bioactive compounds with potent anti-fibrillogenic, antioxidative, and cytocompatible properties. *Biomed Pharmacother.* 2021;141:111811. doi: 10.1016/j.biopha.2021.111811.
29. Madhulika Pradhan, S Sribhuwaneswari, D Karthikeyan, Sunita Minz, Pavani Sure, Atul N Chandu, Umesh Mishra, K Kamalakannan, A Saravanankumar, T Sivakumar. In-vitro cytoprotection activity of foeniculum vulgare and helicteres isora in cultured human blood lymphocytes and antitumour activity against b16f10 melanoma cell line. *Research J. Pharm. and Tech.* 2018; 1(4):450-452
30. Radha Krishna Kollipara, Vyshnavi Tallapaneni, Bharat Kumar Reddy Sanapalli, Vinoth Kumar G, Veera Venkata Satyanarayana Reddy Karri. Curcumin loaded ethosomal vesicular drug delivery system for the treatment of melanoma skin cancer. *Research Journal of Pharmacy and Technology.* 2019; 12(4):1783-1792. doi: 10.5958/0974-360X.2019.00298.1
31. Perumal S, Langeswaran K. Diosmin anti-tumour efficacious against hepatocellular carcinoma. *Research J. Pharm. and Tech.* 2020; 13(4):1707-1714. doi: 10.5958/0974-360X.2020.00308.X
32. Ghadge Ojaswi, Nadar Divya, Parmar Digna. Melanoma and its drug targets. *Research J. Pharm. and Tech.* 2016; 9(5): 562-570. doi: 10.5958/0974-360X.2016.00107.4

KOMENTAR EDITOR SETELAH REVISI DRAFT ARTIKEL UNTUK PROSES REVIEW OLEH REVIEWER



REVIEW DARI REVIEWER

Type of Manuscript:

Research

**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT)
Cells from Siam Weeds (*Chromolaena odorata* L.)**

ABSTRACT

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. *Chromolaena odorata* was extracted with methanol, then partitioned with 5 dissolutions: n-Hexane, Ethyl Acetate, Acetonitrilic, n-Buthanol, Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocytes cells using the bioassay method. After incubation, the result obtained from microplate reader. Each extract was divided into three concentrations, it is 100, 50, 20 ($\mu\text{g}/\text{mL}$). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10 ($\mu\text{g}/\text{mL}$). After being analyzed with the results of the microplate reader, the IC50 of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100 $\mu\text{g}/\text{mL}$. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μl of medium/well.

KEYWORDS: *Chromolaena odorata* L., Skin Cancer, HacaT cell line.

Commented [S11]: The research objectives must be stated clearly before the method statement

Commented [S12]: Results and conclusions must be clearly stated

INTRODUCTION

Skin cancer is one of the most common forms of cancer worldwide, the impenetrable topical barrier is compromised slightly in pathologies such as skin cancer. Another disease that has seen growth in local therapeutic applications is skin cancer which is the most common cancer with one in every three cancer diagnoses being skin cancer according to World Health Organization (WHO)¹. Skin cancer is broadly classified into non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinomas and melanomas².

Cancer development usually occurs when DNA is damaged, and our bodies are unable to repair this damage. These damaged cells begin to grow and divide uncontrollably. Such changes in the skin lead to skin cancer. When damage cell multiply, they form tumors².

Skin cancer develops in various layers of the skin. The tendency to develop skin cancer varies for different skin types. Occurrence varies according to skin color and skin type. In addition, heredity also contributes to the development of skin cancer. The most inherited type of cancer is Melanoma².

Skin cancer can be identified through symptoms. Early identification and early detection of medicine and cure. Sometimes symptoms appear at an advanced stage in other types of cancer. Basal cell cancer appears as a lump on the skin anywhere on the body exposed to sunlight. They can be brown bumps or white bumps on our face, hands, neck, and other exposed body parts. Squamous cell carcinoma on the other hand can appear as ulcerated lumps and patches on the skin whereas melanoma can cause unusual growths on the body especially in existing moles. It can be anywhere in our body³.

The American Cancer Society (ACS) has reported that, there were 87,110 newly diagnosed melanomas, also reported that 9,730 people had an estimated melanoma death in 2017. Skin pigmentation is caused by accumulation of melanin granules which are synthesized in melanocytes and transferred to keratinocytes⁴. Keratinocytes secrete cytokines and chemokines on exposure to UV light which affect melanin production⁵. Melanoma develops at any part of the skin due to defects in skin pigmentation but is most susceptible to the trunk in men and the legs in women⁶.

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. Before the body can recover, these deleterious mutations occur in the genes and the damage becomes irreparable. Moreover, to environmental factors, the patient's genetic history can also be a cause of developing cancer⁷.

UV rays consist of UV-A and UV-B radiation. Although UV-B is considered more dangerous, UV-A also has side effects. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-A can also cause DNA damage by generating free radicals and damaging cell membranes which in turn causes skin cancer. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. UV-B skin also causes oxidative damage to the skin. UV-B radiation further interferes with regulatory proteins, anti-apoptotic relational proteins, proapoptotic proteins, transcription factors, protein kinase, inflammatory enzymes, and growth factors which in turn disrupts the cell cycle⁸. This further causes reasonable mutations in various tumor suppressor genes and oncogenes. These effects together lead to uncontrolled keratinocytes mitosis and initiation of skin cancer upon UV exposure⁷.

It is impossible to completely eradicate cancer, but precautions can be taken to prevent cancer. The following steps should be taken; there should be a reduction in sun exposure, use of tobacco should be avoided, sunscreen should be used to prevent exposure of UV rays, and skin cancer cells can also be removed by biopsy⁸.

The Laruna leaf has been used empirically as a traditional medicine by the people of South Sulawesi. Traditional use of *Chromolaena odorata* (L) as a source of medicine has been widely used in West African countries, Asian countries⁹, and Indonesia as wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, anti-immunosuppressive, anticancer, and other activities^{10,11,12}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals (various factors in plant foods), such as carotenoids, antioxidant vitamins, phenolic compounds, terpenoids, steroids, indole, and fiber, etc¹³. Crude drugs are the effective elements considered responsible for reducing cancer risk¹⁴.

Chromolaena odorata has bioactive compounds with a significant therapeutics' effects on some therapeutic effects on some pathological conditions^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40 -trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihidrokaempferide, Apigenin-7,40 -dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin- 40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

Commented [SI3]: data in Indonesia would be better if it was conveyed

Commented [SI4]: The research objectives must be stated clearly before the method statement

MATERIALS AND METHODS

Materials:

Chromolaena odorata collection is in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, the 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitrilic, N-Buthanol, Ethanol.

Commented [S15]: Composition, collection, storge of sample would be better stated clearly

Table 1. Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77.621	1.55

Table 2. Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68.45
	Ethyl Acetate	245.89
	Acetonitrilic	85.22
	n-Buthanol	147.97
	Ethanol	53.22

Cell Culture

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10 mg of each of extract were dissolved in 1 mL of DMSO. The second stock solution (5 mg/mL) was generated by mixing 0.5 mL of this stock solution with 0.5 mL of DMSO, which was mixed well. The third stock solution (2 mg/mL) was mixed by 0,2 mL of DMSO. To make the final serial concentrations, 1 µL aliquots of the sample stock solutions in two-fold dilution (10, 5, 2 mg/mL) were added to the corresponding well (the final volume became 100 µL) to obtain the final concentrations (100, 50, 20 µg/mL)

Cytotoxicity Investigation

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1 µL of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10 µg/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = \left[\frac{(A_{\text{test}} - A_{\text{blank}})}{(A_{\text{control}} - A_{\text{blank}})} \right] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis

Table 3. % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4. Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test (([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100 µg/mL.




Figure 1. HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic, D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate (([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹⁶. IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISCUSSION:

The aim of this study was to investigate the potential of natural products using HaCaT cells as a reliable in vitro model for evaluation.

Keratinocytes are the main cells of the human epidermis. In addition to providing an epidermal barrier function, keratinocytes play a role in the inflammatory response of the skin¹⁷, and the process of wound repair in the re-epithelialization step¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) eliminated the problems observed with primary keratinocytes such as their need for growth supplementation, culture differentiation, donor-related variability, and limited lifespan. HaCaT cells have been used for many skin model studies without altering keratinocyte function, cellular response, or differentiation capacity¹⁹. Monolayer HaCaT cell cultures are used for many applications such as cellular toxicity and in vitro analysis of wound healing. However monolayer culture systems cannot completely replicate the true epidermal structure. Whereas the generation of air-liquid interface in keratinocytes cell cultures might result in a more realistic model of the epidermis²⁰.

Keratinocytes are active in epidermal repair and skin immune defense through secretion of growth factors, cytokines, and chemokines. To standardizing in vitro studies, investigated the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, and in relation to different culture conditions and degrees of differentiation¹⁷.

HaCaT cells are a long-lived, spontaneously immortalized human line that exhibits basal cell properties and displays substantial changes in response to two well-established in vitro prodifferentiating agents: increased cell density and as a function of time in culture¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 µl of medium/well¹⁷.

HaCaT cells were generally maintained in normal DMEM supplemented with 10% FBS, both of which contain sufficient calcium to induce differentiation. Dedifferentiation to the basal phenotype requires prolonged growth in low calcium media²².

Statistical evidence has revealed that most deaths from skin cancer are the result of melanoma. Further investigations have shown that the survival rate in patients depends on the stage of the cancer; Early detection and intervention of melanoma involves a higher chance of cure²³.

Topical therapies are delivered through skin & they reach the systemic circulation by crossing the skin layers. There are some commonly used topical agent which includes some corticosteroids is given as: 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoids, Mechlorethamine, Carmustine, Bexarotene, Tazarotene photodynamic^{24,25}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals, such as antioxidant. *Chromolaena odorata* leaf extract contain various antioxidant, terpenoid, phenolic and

Commented [SI6]: an incomplete sentence stating the purpose of the study

Commented [SI7]: Differences with other research can be clarified by comparing it with previous studies on similar topics.

flavonoid compounds²⁶. In addition, the anti-mitotic activity of the results of this investigation shows that silver green nanoparticles synthesized from *Chromolaena odorata* have a potential effect on cell division²⁷. The phytochemical profile of the phenolic extract of *Chromolaena odorata* has great potential for the identification of new compounds with pharmacotherapeutic value²⁸. As a comparison to *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48%, research on *Helicteres isora* extract at a concentration of 300 µg/ml had an activity equivalent to the activity of the standard drug doxorubicin²⁹.

In addition, regarding skin cancer testing with various extracts, for example, Ethosomal gel which produces controlled release of curcumin, Curcumin persists in deeper layers of the skin for a longer time, as well as maximum deposition. Thus, eliminating melanoma cells completely³⁰. Furthermore, on the carcinogenic anti-cancer potential of diosmin in vivo and in vitro, it can be a flavonoid in the treatment of Hepatocellular Carcinoma due to its therapeutic and pharmacological properties. Although this research requires further research on other types of cancer³¹.

In-depth study of melanoma biology has been influential in considering melanoma as a therapy for one of the most resistant malignancies. In the last 30 years, there have been few options for treating metastatic melanoma, so some more thorough clinical studies have been conducted, such as hormonal therapy, immunotherapy, single and combination agent chemotherapy, biochemotherapy, targeted agent therapy, and combination regimens, show less than optimal results³².

CONCLUSION:

Ethyl acetate extract of *Chromolaena odorata* has anti-skin cancer potential regarding to the cytotoxicity test.

CONFLICT OF INTEREST:

There are no conflicts of interest declared by the authors of this study.

REFERENCES

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashed D, Weller RB, Castellsagué X. Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)*. 2017; 7(Suppl 1):5-19. doi: 10.1007/s13555-016-0165-y
3. Akila Victor, Muhammad Rukunuddin Ghalib. Detection of skin cancer cells – a review. *Research J.Pharm and Tech*; 2017; 10(11):4093-4098. doi: 10.5958/0974-360X.2017.00742.9
4. Scott G. Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Invest Dermatol*. 2012; 132(4):1073-4. doi: 10.1038/jid.2012.20
5. Nasti TH, Timares L, MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*. 2015; 91(1):188-200. doi: 10.1111/php.12335
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan. Protein-protein interaction of mutated agouti signaling protein (asip) to melanocortin receptor 1 (mcr1) in melanoma skin cancer: an insilico study. *Research J. Pharm. and Tech* 2018; 11(9): 3913-3917. doi: 10.5958/0974-360X.2018.00718.7
7. Min Jeong Kim 1, Seo Yeon Shin 1, Nu Ri Song 1, Sunoh Kim 2, Sang Ouk Sun 1 and Kyung Mok Park 1. Bioassay-guided characterization, antioxidant, anti-melanogenic and anti-photoaging activities of pueraria thunbergiana l. leaf extracts in human epidermal keratinocytes (hacat) cells. *MDPI* 10. 2022; 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y. New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*. 2014; 2014:243452. doi: 10.1155/2014/243452
9. Qadir MI. Skin cancer: etiology and management. *Pak J Pharm Sci*. 2016; 29(3):999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med*. 12. 2015; 301–306. <https://doi.org/10.1515/jcim-2014-0033>
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed. Biological activities of *chromolaena odorata*: A mechanistic review. Elsevier. 2021; <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S. Flavonoid glycosides from *chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull*. 59. 2011; 129–131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B. Pharmacological importance of *chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal Vol* 9. 2021; 8–11. Doi:10.47957/ijpda.v9i1.452

14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging, and antimicrobial potential of chromolaena odorata leaf extracts against pathogenic bacterium in wound infections a multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. 20. <https://doi.org/10.1016/j.bcab.2018.05.014>
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J. Kaempferide, the most active among the four flavonoids isolated and characterized from chromolaena odorata, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5. 2015; 100912– 100922. <https://doi.org/10.1039/C5RA19199H>
16. Mkhobongo B, Chandran R, Abrahamse H. In vitro photodynamic treatment modality for a375 melanoma cell line using a sulphonated aluminum phthalocyanine chloride-photosensitizer-gold nanoparticle conjugate. *Pharmaceutics.* 2022; 16;14(11):2474. doi: 10.3390/pharmaceutics14112474
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all. Hacat cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediator of Inflammation.* 2017; <https://doi.org/10.1155/2017/7435621>
18. Pastar I, Stojadinovic O, Yin NC, Ramirez H, Nusbaum AG, Sawaya A, Patel SB, Khalid L, Isseroff RR, Tomic-Canic M. Epithelialization in wound healing: a comprehensive review. *Adv Wound Care (New Rochelle).* 2014; 3(7):445-464. doi: 10.1089/wound.2013.0473
19. Micallef L, Belaubre F, Pinon A, Jayat-Vignoles C, Delage C, Charveron M, Simon A. Effects of extracellular calcium on the growth-differentiation switch in immortalized keratinocyte hacat cells compared with normal human keratinocytes. *Exp Dermatol.* 2009; 18(2):143-51. doi: 10.1111/j.1600-0625.2008.00775.x
20. Pedrosa TDN, Catarino CM, Pennacchi PC, Assis SR, Gimenes F, Consolaro MEL, Barros SBM, Maria-Engler SS. A new reconstructed human epidermis for in vitro skin irritation testing. *Toxicol In Vitro.* 2017; 42:31-37. doi: 10.1016/j.tiv.2017.03.010
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R. Anti-inflammatory, and anti-proliferative properties of Chromolaena odorata leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants.* 20. 2014; 359–371. doi:10.1080/10496475.2013.876698
22. Wilson VG. Growth and differentiation of hacat keratinocytes. *Methods Mol Biol.* 2014; 1195:33-41. doi: 10.1007/7651_2013_42
23. Tammineni Sreelatha, M. V. Subramanyam, M. N. Giri Prasad. A survey work on early detection methods of melanoma skin cancer. *Research J. Pharm. and Tech.* 2019; 12(5):2589-2596. doi: 10.5958/0974-360X.2019.00435.9
24. Micali G, Lacarubba F, Nasca MR, Ferraro S, Schwartz RA. Topical pharmacotherapy for skin cancer: part ii. Clinical applications. *Journal of the American Academy of Dermatology.* 2014; 70(6):979.e1-12; quiz 9912. doi: 10.1016/j.jaad.2013.12.037.
25. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kanya Goyal, Shabby Jindal. Treatment of skin cancer by topical drug delivery of nanoparticles: a review. *Research Journal of Pharmacy and Technology.* 2021; 14(10):5589-8. doi: 10.52711/0974-360X.2021.00973
26. Zainab, Nurlailah, Muhammad Ikhwan Rizki. Identification of active compound and antibacterial activity against gram-positive and gram-negative bacteria of chromolaena odorata leaf extract. *Research Journal of Pharmacy and Technology.* 2022; 15(10):4720-6. doi: 10.52711/0974-360X.2022.00793
27. Nishita Gogia, Anandhi D, Kanaga G, Revathi K. In vitro evaluation of anti-mitotic activity of green synthesized silver nanoparticles from ethanolic extract of chromolaena odorata, caesalpinia coriaria (bark and leaves) using allium cepa roots. *Research Journal of Pharmacy and Technology.* 2021; 14(8):4107-0. doi: 10.52711/0974-360X.2021.00711
28. Eze FN, Jayeoye TJ. Chromolaena odorata (Siam weed): A natural reservoir of bioactive compounds with potent anti-fibrillogenic, antioxidative, and cytocompatible properties. *Biomed Pharmacother.* 2021;141:111811. doi: 10.1016/j.biopha.2021.111811.
29. Madhulika Pradhan, S Sribhuaneswari, D Karthikeyan, Sunita Minz, Pavani Sure, Atul N Chandu, Umesh Mishra, K Kamalakannan, A Saravanankumar, T Sivakumar. In-vitro cytoprotection activity of foeniculum vulgare and helicteres isora in cultured human blood lymphocytes and antitumour activity against b16f10 melanoma cell line. *Research J. Pharm. and Tech.* 2018; 1(4):450-452
30. Radha Krishna Kollipara, Vyshnavi Tallapaneni, Bharat Kumar Reddy Sanapalli, Vinoth Kumar G, Veera Venkata Satyanarayana Reddy Karri. Curcumin loaded ethosomal vesicular drug delivery system for the treatment of melanoma skin cancer. *Research Journal of Pharmacy and Technology.* 2019; 12(4):1783-1792. doi: 10.5958/0974-360X.2019.00298.1
31. Perumal S, Langeswaran K. Diosmin anti-tumour efficacious against hepatocellular carcinoma. *Research J. Pharm. and Tech.* 2020; 13(4):1707-1714. doi: 10.5958/0974-360X.2020.00308.X
32. Ghadge Ojaswi, Nadar Divya, Parmar Digna. Melanoma and its drug targets. *Research J. Pharm. and Tech.* 2016; 9(5): 562-570. doi: 10.5958/0974-360X.2016.00107.4

PERBAIKAN DARI HASIL REVIEW OLEH REVIEWER

Type of Manuscript:

Research

**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT)
Cells from Siam Weeds (*Chromolaena odorata* L.)**

ABSTRACT

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. *Chromolaena odorata* was extracted with methanol, then partitioned with 5 dissolutions: n-Hexane, Ethyl Acetate, Acetonitrilic, n-Butanol, Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocytes cells using the bioassay method. After incubation, the result obtained from microplate reader. Each extract was divided into three concentrations, it is 100, 50, 20 ($\mu\text{g}/\text{mL}$). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10 ($\mu\text{g}/\text{mL}$). After being analyzed with the results of the microplate reader, the IC50 of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100 $\mu\text{g}/\text{mL}$. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μl of medium/well.

KEYWORDS: *Chromolaena odorata* L., Skin Cancer, HacaT cell line.

INTRODUCTION

Skin cancer is one of the most common forms of cancer worldwide, the impenetrable topical barrier is compromised slightly in pathologies such as skin cancer. Another disease that has seen growth in local therapeutic applications is skin cancer which is the most common cancer with one in every three cancer diagnoses being skin cancer according to World Health Organization (WHO)¹. Skin cancer is broadly classified into non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinomas and melanomas².

Cancer development usually occurs when DNA is damaged, and our bodies are unable to repair this damage. These damaged cells begin to grow and divide uncontrollably. Such changes in the skin lead to skin cancer. When damage cell multiply, they form tumors².

Skin cancer develops in various layers of the skin. The tendency to develop skin cancer varies for different skin types. Occurrence varies according to skin color and skin type. In addition, heredity also contributes to the development of skin cancer. The most inherited type of cancer is Melanoma².

Skin cancer can be identified through symptoms. Early identification and early detection of medicine and cure. Sometimes symptoms appear at an advanced stage in other types of cancer. Basal cell cancer appears as a lump on the skin anywhere on the body exposed to sunlight. They can be brown bumps or white bumps on our face, hands, neck, and other exposed body parts. Squamous cell carcinoma on the other hand can appear as ulcerated lumps and patches on the skin whereas melanoma can cause unusual growths on the body especially in existing moles. It can be anywhere in our body³.

The American Cancer Society (ACS) has reported that, there were 87,110 newly diagnosed melanomas, also reported that 9,730 people had an estimated melanoma death in 2017. Skin pigmentation is caused by accumulation of melanin granules which are synthesized in melanocytes and transferred to keratinocytes⁴. Keratinocytes secrete cytokines and chemokines on exposure to UV light which affect melanin production⁵. Melanoma develops at any part of the skin due to defects in skin pigmentation but is most susceptible to the trunk in men and the legs in women⁶.

Most cancers develop due to excessive sun exposure. It is known that sunlight consists of ultraviolet rays which are harmful to the skin and damage DNA. Before the body can recover, these deleterious mutations occur in the genes and the damage becomes irreparable. Moreover, to environmental factors, the patient's genetic history can also be a cause of developing cancer⁷.

UV rays consist of UV-A and UV-B radiation. Although UV-B is considered more dangerous, UV-A also has side effects. UV-A reaches deep into the dermis, causing skin P stem cell carcinogenesis. UV-A can also cause DNA damage by generating free radicals and damaging cell membranes which in turn causes skin cancer. UV-B is absorbed by proteins in the epidermis and causes DNA damage to cells. UV-B skin also causes oxidative damage to the skin. UV-B radiation further interferes with regulatory proteins, anti-apoptotic relational proteins, proapoptotic proteins, transcription factors, protein kinase, inflammatory enzymes, and growth factors which in turn disrupts the cell cycle⁸. This further causes reasonable mutations in various tumor suppressor genes and oncogenes. These effects together lead to uncontrolled keratinocytes mitosis and initiation of skin cancer upon UV exposure⁷.

It is impossible to completely eradicate cancer, but precautions can be taken to prevent cancer. The following steps should be taken; there should be a reduction in sun exposure, use of tobacco should be avoided, sunscreen should be used to prevent exposure of UV rays, and skin cancer cells can also be removed by biopsy⁸.

The Laruna leaf has been used empirically as a traditional medicine by the people of South Sulawesi. Traditional use of *Chromolaena odorata* (L) as a source of medicine has been widely used in West African countries, Asian countries⁹, and Indonesia as wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, anti-immunosuppressive, anticancer, and other activities^{10,11,12}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals (various factors in plant foods), such as carotenoids, antioxidant vitamins, phenolic compounds, terpenoids, steroids, indole, and fiber, etc¹³. Crude drugs are the effective elements considered responsible for reducing cancer risk¹⁴.

Chromolaena odorata is has bioactive compounds with a significant therapeutics' effects on some therapeutic effects on some pathological conditions^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40 -trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihydrokaempferide, Apigenin-7,40 -dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin- 40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

MATERIALS AND METHODS

Materials:

Chromolaena odorata collection is in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, the 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitrilic, N-Buthanol, Ethanol.

Table 1. Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77.621	1.55

Table 2. Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68.45
	Ethyl Acetate	245.89
	Acetonitrilic	85.22
	n-Buthanol	147.97
	Ethanol	53.22

Cell Culture

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10 mg of each of extract were dissolved in 1 mL of DMSO. The second stock solution (5 mg/mL) was generated by mixing 0.5 mL of this stock solution with 0.5 mL of DMSO, which was mixed well. The third stock solution (2 mg/mL) was mixed by 0,2 mL of DMSO. To make the final serial concentrations, 1 μ L aliquots of the sample stock solutions in two-fold dilution (10, 5, 2 mg/mL) were added to the corresponding well (the final volume became 100 μ L) to obtain the final concentrations (100, 50, 20 μ g/mL)

Cytotoxicity Investigation

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1 μ L of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10 μ g/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = \left[\frac{1 - (A_{\text{test}} - A_{\text{blank}})}{(A_{\text{control}} - A_{\text{blank}})} \right] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis

Table 3. % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4. Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100 µg/mL.




Figure 1. HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic,
D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate ([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹⁶. IC₅₀ value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISCUSSION:

The aim of this study was to investigate the potential of natural products using HaCaT cells as a reliable in vitro model for evaluation.

Keratinocytes are the main cells of the human epidermis. In addition to providing an epidermal barrier function, keratinocytes play a role in the inflammatory response of the skin¹⁷, and the process of wound repair in the re-epithelialization step¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) eliminated the problems observed with primary keratinocytes such as their need for growth supplementation, culture differentiation, donor-related variability, and limited lifespan. HaCaT cells have been used for many skin model studies without altering keratinocyte function, cellular response, or differentiation capacity¹⁹. Monolayer HaCaT cell cultures are used for many applications such as cellular toxicity and in vitro analysis of wound healing. However monolayer culture systems cannot completely replicate the true epidermal structure. Whereas the generation of air-liquid interface in keratinocytes cell cultures might result in a more realistic model of the epidermis²⁰.

Keratinocytes are active in epidermal repair and skin immune defense through secretion of growth factors, cytokines, and chemokines. To standardizing in vitro studies, investigated the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, and in relation to different culture conditions and degrees of differentiation¹⁷.

HaCaT cells are a long-lived, spontaneously immortalized human line that exhibits basal cell properties and displays substantial changes in response to two well-established in vitro prodifferentiating agents: increased cell density and as a function of time in culture¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 µl of medium/well¹⁷.

HaCaT cells were generally maintained in normal DMEM supplemented with 10% FBS, both of which contain sufficient calcium to induce differentiation. Dedifferentiation to the basal phenotype requires prolonged growth in low calcium media²².

Statistical evidence has revealed that most deaths from skin cancer are the result of melanoma. Further investigations have shown that the survival rate in patients depends on the stage of the cancer; Early detection and intervention of melanoma involves a higher chance of cure²³.

Topical therapies are delivered through skin & they reach the systemic circulation by crossing the skin layers. There are some commonly used topical agent which includes some corticosteroids is given as: 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoids, Mechlorethamine, Carmustine, Bexarotene, Tazarotene photodynamic^{24, 25}.

The beneficial effects of crude drugs are believed to be attributed to plant phytochemicals, such as antioxidant. *Chromolaena odorata* leaf extract contain various antioxidant, terpenoid, phenolic and

flavonoid compounds²⁶. In addition, the anti-mitotic activity of the results of this investigation shows that silver green nanoparticles synthesized from *Chromolaena odorata* have a potential effect on cell division²⁷. The phytochemical profile of the phenolic extract of *Chromolaena odorata* has great potential for the identification of new compounds with pharmacotherapeutic value²⁸. As a comparison to *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48%, research on *Helicteres isora* extract at a concentration of 300 µg/ml had an activity equivalent to the activity of the standard drug doxorubicin²⁹.

In addition, regarding skin cancer testing with various extracts, for example, Ethosomal gel which produces controlled release of curcumin, Curcumin persists in deeper layers of the skin for a longer time, as well as maximum deposition. Thus, eliminating melanoma cells completely³⁰. Furthermore, on the carcinogenic anti-cancer potential of diosmin in vivo and in vitro, it can be a flavonoid in the treatment of Hepatocellular Carcinoma due to its therapeutic and pharmacological properties. Although this research requires further research on other types of cancer³¹.

In-depth study of melanoma biology has been influential in considering melanoma as a therapy for one of the most resistant malignancies. In the last 30 years, there have been few options for treating metastatic melanoma, so some more thorough clinical studies have been conducted, such as hormonal therapy, immunotherapy, single and combination agent chemotherapy, biochemotherapy, targeted agent therapy, and combination regimens, show less than optimal results³².

CONCLUSION:

Ethyl acetate extract of *Chromolaena odorata* has anti-skin cancer potential regarding to the cytotoxicity test.

CONFLICT OF INTEREST:


There are no conflicts of interest declared by the authors of this study.

REFERENCES

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashan D, Weller RB, Castellsagué X. Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)*. 2017; 7(Suppl 1):5-19. doi: 10.1007/s13555-016-0165-y
3. Akila Victor, Muhammad Rukunuddin Ghalib. Detection of skin cancer cells – a review. *Research J.Pharm and Tech*; 2017; 10(11):4093-4098. doi: 10.5958/0974-360X.2017.00742.9
4. Scott G. Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Invest Dermatol*. 2012; 132(4):1073-4. doi: 10.1038/jid.2012.20
5. Nasti TH, Timares L. MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*. 2015; 91(1):188-200. doi: 10.1111/php.12335
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan. Protein-protein interaction of mutated agouti signaling protein (asip) to melanocortin receptor 1 (mc1r) in melanoma skin cancer: an insilico study. *Research J. Pharm. and Tech* 2018; 11(9): 3913-3917. doi: 10.5958/0974-360X.2018.00718.7
7. Min Jeong Kim 1, Seo Yeon Shin 1, Nu Ri Song 1, Sunoh Kim 2, Sang Ouk Sun 1 and Kyung Mok Park 1. Bioassay-guided characterization, antioxidant, anti-melanogenic and anti-photoaging activities of pueraria thunbergiana l. leaf extracts in human epidermal keratinocytes (hacat) cells. *MDPI* 10. 2022; 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y. New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*. 2014; 2014:243452. doi: 10.1155/2014/243452
9. Qadir MI. Skin cancer: etiology and management. *Pak J Pharm Sci*. 2016; 29(3):999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med*. 12. 2015; 301–306. <https://doi.org/10.1515/jcim-2014-0033>
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed. Biological activities of *chromolaena odorata*: A mechanistic review. *Elsevier*. 2021; <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S. Flavonoid glycosides from *chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull*. 59. 2011; 129–131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B. Pharmacological importance of *chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal Vol* 9. 2021; 8–11. Doi:10.47957/ijpda.v9i1.452

14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging, and antimicrobial potential of chromolaena odorata leaf extracts against pathogenic bacterium in wound infections a multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. 20. <https://doi.org/10.1016/j.bcab.2018.05.014>
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J. Kaempferide, the most active among the four flavonoids isolated and characterized from chromolaena odorata, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5. 2015; 100912–100922. <https://doi.org/10.1039/C5RA19199H>
16. Mkhobongo B, Chandran R, Abrahamse H. In vitro photodynamic treatment modality for a375 melanoma cell line using a sulphonated aluminum phthalocyanine chloride-photosensitizer-gold nanoparticle conjugate. *Pharmaceutics.* 2022; 16;14(11):2474. doi: 10.3390/pharmaceutics14112474
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all. Hacat cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediator of Inflammation.* 2017; <https://doi.org/10.1155?2017/7435621>
18. Pastar I, Stojadinovic O, Yin NC, Ramirez H, Nusbaum AG, Sawaya A, Patel SB, Khalid L, Isseroff RR, Tomic-Canic M. Epithelialization in wound healing: a comprehensive review. *Adv Wound Care (New Rochelle).* 2014; 3(7):445-464. doi: 10.1089/wound.2013.0473
19. Micallef L, Belaubre F, Pinon A, Jayat-Vignoles C, Delage C, Charveron M, Simon A. Effects of extracellular calcium on the growth-differentiation switch in immortalized keratinocyte hacat cells compared with normal human keratinocytes. *Exp Dermatol.* 2009; 18(2):143-51. doi: 10.1111/j.1600-0625.2008.00775.x
20. Pedrosa TDN, Catarino CM, Pennacchi PC, Assis SR, Gimenes F, Consolaro MEL, Barros SBM, Maria-Engler SS. A new reconstructed human epidermis for in vitro skin irritation testing. *Toxicol In Vitro.* 2017; 42:31-37. doi: 10.1016/j.tiv.2017.03.010
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R. Anti-inflammatory, and anti-proliferative properties of Chromolaena odorata leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants* 20. 2014; 359–371. doi:10.1080/10496475.2013.876698
22. Wilson VG. Growth and differentiation of hacat keratinocytes. *Methods Mol Biol.* 2014; 1195:33-41. doi: 10.1007/7651_2013_42
23. Tammineni Sreelatha, M. V. Subramanyam, M. N. Giri Prasad. A survey work on early detection methods of melanoma skin cancer. *Research J. Pharm. and Tech.* 2019; 12(5):2589-2596. doi: 10.5958/0974-360X.2019.00435.9
24. Micali G, Lacarubba F, Nasca MR, Ferraro S, Schwartz RA. Topical pharmacotherapy for skin cancer: part ii. Clinical applications. *Journal of the American Academy of Dermatology.* 2014; 70(6):979.e1-12; quiz 9912. doi: 10.1016/j.jaad.2013.12.037.
25. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kamyra Goyal, Shamy Jindal. Treatment of skin cancer by topical drug delivery of nanoparticles: a review. *Research Journal of Pharmacy and Technology.* 2021; 14(10):5589-8. doi: 10.52711/0974-360X.2021.00973
26. Zainab, Nurlailah, Muhammad Ikhwan Rizki. Identification of active compound and antibacterial activity against gram-positive and gram-negative bacteria of chromolaena odorata leaf extract. *Research Journal of Pharmacy and Technology.* 2022; 15(10):4720-6. doi: 10.52711/0974-360X.2022.00793
27. Nishita Gogia, Anandhi D, Kanaga G, Revathi K. In vitro evaluation of anti-mitotic activity of green synthesized silver nanoparticles from ethanolic extract of chromolaena odorata, caesalpinia coriaria (bark and leaves) using allium cepa roots. *Research Journal of Pharmacy and Technology.* 2021; 14(8):4107-0. doi: 10.52711/0974-360X.2021.00711
28. Eze FN, Jayeoye TJ. Chromolaena odorata (Siam weed): A natural reservoir of bioactive compounds with potent anti-fibrillogenic, antioxidative, and cytocompatible properties. *Biomed Pharmacother.* 2021;141:111811. doi: 10.1016/j.biopha.2021.111811.
29. Madhulika Pradhan, S Sribhuwaneswari, D Karthikeyan, Sunita Minz, Pavani Sure, Atul N Chandu, Umesh Mishra, K Kamalakannan, A Saravanankumar, T Sivakumar. In-vitro cytoprotection activity of foeniculum vulgare and helicteres isora in cultured human blood lymphocytes and antitumour activity against b16f10 melanoma cell line. *Research J. Pharm. and Tech.* 2018; 1(4):450-452
30. Radha Krishna Kollipara, Vyshnavi Tallapaneni, Bharat Kumar Reddy Sanapalli, Vinoth Kumar G, Veera Venkata Satyanarayana Reddy Karri. Curcumin loaded ethosomal vesicular drug delivery system for the treatment of melanoma skin cancer. *Research Journal of Pharmacy and Technology.* 2019; 12(4):1783-1792. doi: 10.5958/0974-360X.2019.00298.1
31. Perumal S, Langeswaran K. Diosmin anti-tumour efficacious against hepatocellular carcinoma. *Research J. Pharm. and Tech.* 2020; 13(4):1707-1714. doi: 10.5958/0974-360X.2020.00308.X
32. Ghadge Ojaswi, Nadar Divya, Parmar Digna. Melanoma and its drug targets. *Research J. Pharm. and Tech.* 2016; 9(5): 562-570. doi: 10.5958/0974-360X.2016.00107.4

Informasi payment setelah perbaikan diterima



A AND V PUBLICATIONS

Logout

SEARCH
DASHBOARD
PAPER SUBMISSION
SUBSCRIPTION
BOOKS
BOOK ADVERTISEMENT
MY CART (0)
MY ORDERS

Paper ID: 2356034457024539

Title: Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)

Authors: Karlina Amir Tahir;Erwin Hafid;Muh. Fitrah;Ahmad Lalo;Nurul Fadilah;Syamsuri Syakri;Syatirah Jalaluddin;Katsuyoshi Matsunami
(karlina.amir@uin-alauddin.ac.id;erwin.hafid@uin-alauddin.ac.id;muhfitrah@uin-alauddin.ac.id;ahmadlalo_pom@yahoo.co.id;syamsurisyakri@gmail.com;syatirahjalaluddin@gmail.com)

Submitted To: Research Journal of Pharmacy and Technology

Payment Transaction No.: pay_NVWeQjODHeeiGO

Paid On: 2/1/2024 12:00:00 AM

INR: 3540.00


Or USD: 210.00

PAYMENT VERIFIED (03-Feb-2024)

- [Editorial Comments](#)
- [Reviewer Comments](#)
- [Download Acceptance Letter](#)
- [Print Report](#)

6/25/24, 6:32 PM

Article Status



A and V Publications

Report Generated on: 6/25/2024 4:56:55 PM

BASIC INFORMATION

Paper ID: 2356034457024539	Submission Date: May 06, 2023
Paper Title: Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (<i>Chromolaena odorata</i> L.)	
Author(s) Name: Karlina Amir Tahir;Erwin Hafid;Muh. Fitrah;Ahmad Lalo;Nurul Fadilah;Syamsuri Syakri;Syatirah Jalaluddin;Katsuyoshi Matsunami	
Author(s) Email: karlina.amir@uin-alsuddin.ac.id;erwin.hafid@uin-alauddin.ac.id;muhfitrah@uin-alauddin.ac.id;ahmadlalo_pom@yahoo.co.id;syamsurisyakri@gmail.com;syatirahjalaluddin@gmail.com;	
Author(s) Address: H.M Yasin Limpo No.36, Gowa, South of Sulawesi, Indonesia	
Journal: Research Journal of Pharmacy and Technology	
Submitted By: karlina amir tahir	Email ID: nilalfath@gmail.com

REVIEWER INFORMATION

	First Reviewer	Second Reviewer
Name:	SUNANDAR IHSAN	PROF. DR. KUMARASWAMY GANDLA
Email ID:	sunandarihsan@uho.ac.id	drkumaraswamygandla@gmail.com
Mobile No:	081342987228	0919000973789
Address:	Universitas Halu Oleo, Kendari, Indonesia	Assoc. editor of RJPT

PAYMENT DETAILS

Order No.: 2024211219014432030539092	Order Date: February 01, 2024, 12:02
Amount INR: 3540.00	OR Amount USD: 210.00
Payment Date: February 01, 2024	Transaction No.: pay_NVWeQjODHeeiGO
Payment Status: PAYMENT VERIFIED	

Status : mulai tanggal 6 mei 2023 sd januari 2024 artikel diterima oleh publisher untuk terbit di bulan Mei 2024

PAPER PUBLICATION / PROCESSING STATUS

- ▶ 06/May/2023, 03:44:57 AM --- Article submitted by the author.
- ▶ 10/May/2023, 11:32:54 AM --- Article sent back to author for minor corrections.
- ▶ 10/May/2023, 11:32:54 AM --- New comments from editorial board.
- ▶ 14/May/2023, 07:01:16 AM --- Article resubmitted by author after correction.
- ▶ 26/Jul/2023, 06:55:42 PM --- Article is sent to reviewers.
- ▶ 13/Sep/2023, 04:53:18 PM --- Article sent back to author for minor corrections.
- ▶ 13/Sep/2023, 04:53:28 PM --- New comments from editorial board.
- ▶ 20/Sep/2023, 04:36:22 AM --- Article resubmitted by author after correction.
- ▶ 20/Sep/2023, 04:36:26 AM --- Article resubmitted by author after correction.
- ▶ 23/Oct/2023, 06:02:46 PM --- Article sent back to author for minor corrections.
- ▶ 23/Oct/2023, 06:05:25 PM --- New comments from editorial board.
- ▶ 31/Oct/2023, 12:32:10 PM --- Article resubmitted by author after correction.
- ▶ 31/Oct/2023, 12:32:16 PM --- Article resubmitted by author after correction.
- ▶ 31/Oct/2023, 12:32:23 PM --- Article resubmitted by author after correction.
- ▶ 31/Oct/2023, 12:32:31 PM --- Article resubmitted by author after correction.
- ▶ 25/Dec/2023, 03:00:03 PM --- Article sent back to author for minor corrections.
- ▶ 25/Dec/2023, 03:00:14 PM --- New comments from editorial board.
- ▶ 04/Jan/2024, 03:49:54 AM --- Article resubmitted by author after correction.
- ▶ 25/Jan/2024, 05:54:10 PM --- Final version of article is required.
- ▶ 27/Jan/2024, 04:46:09 AM --- Final version of article submitted by author.
- ▶ 27/Jan/2024, 04:46:09 AM --- Article is accepted by publisher.
- ▶ 27/Jan/2024, 04:46:13 AM --- Final version of article submitted by author.
- ▶ 27/Jan/2024, 04:46:13 AM --- Article is accepted by publisher.


<https://www.anvpublication.org/ArticleStatus.aspx?PID=2356034457024539>

6/25/24, 6:32 PM

Article Status

- ▶ 01/Feb/2024, 12:19:02 PM --- Payment proof required.
- ▶ 01/Feb/2024, 12:49:17 PM --- Payment proof uploaded by author.
- ▶ 03/Feb/2024, 05:14:22 PM --- Payment details verified by publisher.
- ▶ 05/Feb/2024, 04:10:48 PM --- Payment details verified by publisher.

Status Galley proof sebelum terbit :



Filter

Reset

Q2


Inhibition of Skin Cancer using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)

Creator : Tahir K.A.


Research Journal of Pharmacy and Technology

Journal
publish at
2024

Documents cited by public policy (Overton)



Documents related to SDGs (UN)



Research Journal of Pharmacy and Technology

Q2

Pharmacology, Toxicology and Pharmaceutics...
best quartile

SJR 2023
0.27

powered by scimagajr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagajr.com">
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.



Date: January 27, 2024

Research Journal of Pharmacy and Technology

Paper ID:2356034457024539

Author's Name:Karlina Amir Tahir, Erwin Hafid, Muh. Fitrah, Ahmad Lalo, Nurul Fadilah,

Syamsuri Syakri, Syatirah Jalaluddin, Katsuyoshi Matsunami

Paper Title: Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)



Acceptance of Manuscript

With reference to your article titled '**Inhibition of Skin Cancer Using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)**' Author by **Karlina Amir Tahir, Erwin Hafid, Muh. Fitrah, Ahmad Lalo, Nurul Fadilah, Syamsuri Syakri, Syatirah Jalaluddin, Katsuyoshi Matsunami**. We wish to bring to your kind notice the following:

- We acknowledge the receipt of the above mentioned article.
- The above mentioned article has been sent to the reviewer of expert comments.
- The above mentioned article have been accepted for publication in Year : 2024, Vol: 17, Issue:5 of the journal.

Thanking you for submission of manuscript.

Yours sincerely,



Sr. Baharwal

Editor in Chief


Note: -

1. To verify the originality of this certificate please visit the following link:


http://www.anvpublication.org/Download_Acceptance.aspx?PID=2356034457024539

PRINT

KOMENTAR EDITOR UNTUK GALLEY PROOF SEBELUM TERBIT



Komentar editor untuk artikel/jurnal yang terbit dibulan Mei berdasarkan hasil koreksi di galley proof



RESEARCH ARTICLE

Inhibition of Skin Cancer using Human Epidermal Keratinocytes (HaCaT) Cells from Siam Weeds (*Chromolaena odorata* L.)

Karlina Amir Tahir¹, Erwin Hafid², Muh. Fitrah¹, Ahmad Lalo³, Nurul Fadilah¹, Syamsuri Syakri¹, Syatirah Jalaluddin¹, Katsuyoshi Matsunami⁴

¹Faculty of Medicine and Health Science, UIN Alauddin Makassar, Makassar, Indonesia.

²Master of Islamic Education Management, UIN Alauddin Makassar, Makassar, Indonesia.

³National Agency of Drug and Food Control, Makassar, Indonesia.

⁴Department of Pharmacognosy, Faculty of Biomedical and Health Science at Hiroshima University, Japan.

*Corresponding Author E-mail: karlina.amir@uin-alauddin.ac.id

ABSTRACT:

Cancer can usually develop due to exposure to sunlight. UV radiation from sunlight is known to damage DNA and is bad for the skin. Skin P stem cell carcinogenesis is caused by UV-A rays that penetrate deep into the dermis layer. UV-B damages cell DNA by being absorbed by proteins in the epidermis. *Chromolaena odorata* was extracted using methanol solvent, then partitioned into 5 solutions in the form of n-Hexane, Ethyl Acetate, Acetonitrile, n-Butanol, and Ethanol. The five extracts obtained were tested with Human Epidermal Keratinocyte cells using the bioassay method. Results obtained from the microplate reader after incubation. Each extract was divided into three concentrations, it is 100, 50, 20(µg/mL). Then in the positive control (Etoposide), it was divided into four concentrations, 100, 50, 20, 10(µg/mL). After being analyzed with the results of the microplate reader, the IC₅₀ of *Chromolaena odorata* was 48% in the ethyl acetate extract with a concentration of 100µg/mL. HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100µl of medium/well.

KEYWORDS: *Chromolaena odorata* L, Skin Cancer, HacaT cell line.

INTRODUCTION:

One of the most common types of cancer in the world is skin cancer, which is characterized by the breakdown of the airtight topical barrier. According to the World Health Organization (WHO), skin cancer is a cancer that often occurs in one in three cancer diagnoses, and the application of local treatment for this disease is experiencing increasing development¹. Based on the classification, skin cancer consists of melanoma and non-melanoma skin cancer which includes basal cell carcinoma and squamous cell carcinoma².

Cancer generally develops due to DNA damage that our body cannot repair. Uncontrolled, these injured cells begin to grow and divide. When these changes occur in the skin, they can cause skin cancer. The proliferation of damaged cells can cause tumors².

Skin cancer can arise in any of the skin's layers. Different skin types have differing propensities to develop skin cancer. The frequency varies according to the kind and color of the skin. Furthermore, genetics plays a role in the development of skin cancer. Melanoma is the most common hereditary kind of cancer².

The symptoms that appear can be used to diagnose skin cancer. Identify and detect the drugs and therapies used early. In other forms of cancer, the appearance of symptoms occurs in the advanced stages of the disease.

On any part of the body exposed to sunlight, basal cell carcinoma manifests as a lump on the skin. On the hands, face, neck, and other exposed areas of the body, they may appear as white or brown bumps. Melanoma can cause unusual developments on the body, especially on existing moles, while squamous cell carcinoma can manifest as lumps and ulcerated patches on the skin. This can happen anywhere in our bodies³.

According to data from the American Cancer Society (ACS), 9,730 persons lost their lives to melanoma in 2017 and there were 87,110 newly diagnosed melanomas. Melanin granules, which are produced in melanocytes and transported to keratinocytes, accumulate and result in skin pigmentation⁴. When exposed to UV radiation, keratinocytes release cytokines and chemokines that influence the formation of melanin⁵. Skin pigmentation abnormalities can cause melanoma to grow on any area of the skin, although in men and women it most commonly affects the trunk and legs⁶.

One of the biggest factors that causes cancer to develop is excessive exposure to sunlight. It is known that UV radiation from sunlight can injure the skin and cause DNA to be damaged. These dangerous genetic changes occur before the body has time to recover, causing permanent damage. Apart from that, cancer can also be caused by environmental factors and the patient's genetic background⁷.

UV radiation is divided into UV-A and UV-B photons. Although UV-B is considered more dangerous, UV-A can also have negative consequences. Skin P stem cell carcinogenesis is caused by UV-A rays that penetrate deep into the dermis. By producing free radicals and damaging cell membranes, UV-A can also damage DNA, which can lead to skin cancer. UV-B damages cell DNA by being absorbed by proteins in the epidermis. Skin exposed to UV-B rays also experiences oxidative damage. The cell cycle is further disrupted by UV-B radiation interference with growth factors, transcription factors, anti-apoptotic relational proteins, pro-apoptotic proteins, inflammatory enzymes, and regulatory proteins⁸. It also causes mutations in a number of oncogenes and tumor suppressor genes. This can lead to uncontrolled keratinocyte mitosis and initiation of skin cancer upon exposure to UV light⁷.

Although cancer cannot be completely eradicated, there are steps you can take to reduce the risk. The actions that must be taken are minimizing exposure to sunlight, not using tobacco, using sunscreen to protect the skin from UV radiation, and removing skin cancer cells through biopsy⁸.

The people of South Sulawesi have empirically used Laruna leaves as traditional medicine. *Chromolaena odorata* (L) has long been used in traditional West African, Asian and Indonesian medicine for wound healing, antibacterial, antimalarial, anti-inflammatory, antioxidant, antiimmunosuppressive and anticancer, as well as other uses^{10,11,12}.

The benefits of raw medicines are related to plant phytochemicals, which include phenolic compounds, carotenoids, antioxidant vitamins, terpenoids, steroids, indole, fiber, etc. The beneficial components thought to be responsible for lowering the risk of cancer are crude drugs¹⁴.

The bioactive compounds found in *Chromolaena odorata* have important therapeutic benefits in a number of clinical disorders^{13,14}. Odoratin, Persicogenin, Acacetin, Ombuin, 5,7-Dihydroxy-6,40-dimethoxyflavanone, Erio-dictyol-7,30,40-trimethyl ether, Alyssifolinone, Aromadendrin-7-dimethyl ether, Dihydrokaempferide, Apigenin-7,40- dimethyl ether, Isosakuranetin, Sakuranetin, Sakuranetin-40-methyl ether, Aromadendrin-7,40-dimethyl ether^{12,15}.

MATERIALS AND METHODS:

Materials:

Chromolaena odorata collection is in Pangkep regency, and the following ingredients were used; The HacaT cell, Dulbecco's Modified Eagle's Media, dimethyl sulfoxide (DMSO), etoposide, MTT, the 96-well plates, Solvents for the extraction line, obtained from the faculty of Biomedical and Health at Hiroshima University, Japan.

Methods:

This is a quantitative laboratory experimental study with a pure experimental research design (True Experimental): post test only control design. This study was carried out in Department of Pharmacognosy at Hiroshima University in Japan.

Extractions:

Chromolaena odorata extract was dissolved in methanol to obtain a thick methanol extract. Then partitioned into solid liquid using n-Hexane, Ethyl Acetate, Acetonitril, N-Buthanol, Ethanol.

Table 1: Extraction weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight (gr)	% Rendamen
<i>Chromolaena odorata</i> 5 kg	Metanol	77.621	1.55

Table 2: Partition weight of *Chromolaena odorata*

Sample	Solvent	Extract Weight
Methanol extract of <i>Chromolaena odorata</i>	n-Hexane	68.45
	Ethyl Acetate	245.89
	Acetonitrilic	85.22
	n-Buthanol	147.97
	Ethanol	53.22

Cell Culture:

Preparation of the Stock Solutions and the Serial Concentrations To make the initial stock solution (10 mg/mL) of *Chromolaena odorata* samples, 10mg of each of extract were dissolved in 1mL of DMSO. The second stock solution (5mg/mL) was generated by mixing 0.5mL of this stock solution with 0.5mL of DMSO, which was mixed well. The third stock solution (2mg/mL) was mixed by 0,2mL of DMSO. To make the final serial concentrations, 1µL aliquots of the sample stock solutions in two-fold dilution (10, 5, 2mg/mL) were added to the corresponding well (the final volume became 100µL) to obtain the final concentrations (100, 50, 20µg/mL)

Cytotoxicity Investigation:

The colorimetric cell viability MTT assay was used to determine the cytotoxicity against the Human epidermal keratinocytes (HaCaT) cells. DMSO was used as a negative control, and etoposide as a positive control. A total of 1µL of each etoposide stock solution was added to the corresponding wells of a 96 microplate, resulting in final serial concentrations of (100, 50, 20, 10µg/mL).

The cytotoxic activity was calculated from the equation:

$$\% \text{ inhibition} = [1 - (A_{\text{test}} - A_{\text{blank}}) / (A_{\text{control}} - A_{\text{blank}})] \times 100.$$

Where A control is the absorbance of the control (DMSO) well, A test is the absorbance of the test wells, and Ablank is the absorbance of the cell-free wells.

Data Analysis:

Table 3: % inhibition

Sample	Concentration		
	100	50	20
Hexane	16.56	5.54	8.36
Ethyl Acetate	48.00	5.12	12.20
Acetonitrilic	39.62	29.59	-1.97
Buthanol	7.21	14.21	14.65
Ethanol	-2.53	1.30	19.96

Table 4: Control positive (Etoposide)

Control	Concentration			
	100	50	20	10
Etoposide	79.82	78.91	69.30	73.41

RESULTS:

The results of the cytotoxic test are useful for knowing how much the cytotoxic effect and anticancer potential of *Chromolaena odorata* extract. The results are determined by calculating the IC50 value, the concentration value capable of suppressing the growth of 50% of the cancer cell population based on MTT test ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]).

The result of determining the percentage inhibition of the extracts as shown in the graph. The ethyl acetate extract was the most active, with an IC value of 48%, with a concentration of 100µg/mL.




Figure 1: HaCaT cytotoxicity

A = n-Hexane, B = Ethyl Acetate, C = Acetonitrilic, D = n-Buthanol, E = Ethanol.

The cytotoxic parameter used was the ability to convert MTT substrate ([3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]) into purple formazan by the enzyme succinate dehydrogenase in living cells. In the MTT method, the percentage of cell death is the difference between the absorbance of the control and the absorbance of the sample divide by the absorbance of the control multiplied by 100%. The data was processed using probit analysis to obtain IC values for the sample¹⁶. IC50 value indicates the percentage of cell death in culture as much as 50%.

Based on research that has been done, the result of microplate reader on Human Epidermal Keratinocytes cells were treated *Chromolaena odorata* of ethyl acetate extract at concentrations of 100 µg/mL has the highest percentage of 48% among other extracts.

DISCUSSION:

This research aims to identify the potential of natural ingredients by using HaCaT cells as a reliable in vitro model for evaluation.

The main cells of the human epidermis are called keratinocytes. Apart from their role in maintaining the

epidermal barrier, keratinocytes also have a role in the inflammatory response of the skin¹⁷ and the re-epithelialization stage of wound healing¹⁸. Establishment of an immortalized human keratinocyte cell line (HaCaT) addresses the problems observed with primary keratinocytes in the form of their need for growth supplementation, culture differentiation, donor-related variability, and limited lifespan. Many skin model investigations have included HaCaT cells without compromising keratinocyte function, cellular response, or differentiation potential¹⁹. There are many uses for monolayer culture of HaCaT cells, including analysis of in vitro wound healing and cellular toxicity. However, the actual structure of the epidermis cannot be completely replicated by monolayer culture methods. On the other hand, keratinocyte cell cultures that provide an air-liquid interface can produce a more accurate representation of the epidermis²⁰.

In the process of repairing the epidermis and maintaining the skin's immune defenses, keratinocytes have an active role by secreting growth factors, cytokines, and chemokines. For in vitro standardization studies, investigate the use of HaCaT cells as a suitable model to follow the release of inflammatory mediators and skin repair in response to TNF α or IL-1 β ²¹, as well as their association with different culture states and different degrees of differentiation¹⁷.

The human line HaCaT cells are long-lived and spontaneously immortalized. They possess basal cell characteristics and respond significantly to two well-known in vitro prodifferentiating agents: increasing cell density and culture duration¹⁷.

HaCaT cell proliferation was determined at indicated intervals using the MTT colorimetric assay. This assay was based on the ability of live cell succinate dehydrogenase to reduce the yellow salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma-Aldrich, St. Louis, MO, USA) to insoluble purple-blue formazan precipitate. Experiments were carried out on 96-well plates containing a final volume of 100 μ l of medium/well¹⁷.

The standard DMEM supplemented with 10% FBS, both of which contain enough calcium to trigger differentiation, was used to sustain HaCaT cells. It takes longer to dedifferentiate to the basal phenotype in low-calcium medium²².

According to statistical data, melanoma is the main cause of death due to skin cancer. Further research revealed that a patient's chances of survival depended on the stage of the cancer. Early detection and treatment of melanoma can increase the chances of recovery²³.

Topical medications are absorbed through the skin and pass through the skin layers to enter the systemic circulation. There are several topical agents that are often used including several corticosteroids, namely 5-fluorouracil (5-FU, Efudex), Imiquimod, Resiquimod, Ingenol, Mebutate, Diclofenac, Retinoid, Mechllorethamine, Carmustine, Bexarotene, Tazarotene photodynamic^{24,25}.

Plant phytochemicals, such as antioxidants, are thought to have beneficial effects over crude drugs. *Chromolaena odorata* leaf extract contains many antioxidants, flavonoids, phenolics, terpenoids²⁶. In addition, the results of the investigation show that the anti-mitotic activity of green silver nanoparticles made from *Chromolaena odorata* has a potential effect on cell division²⁷. The phytochemical profile of the phenolic extract of *Chromolaena odorata* has great potential in the discovery of new compounds with pharmacotherapeutic value²⁸. Research on *Chromolaena odorata* ethyl acetate extract at a concentration of 100 μ g/mL had the highest percentage, namely 48%. Meanwhile, *Helicteres isora* extract at a concentration of 300 μ g/ml has activity equivalent to the standard drug doxorubicin²⁹.

Additionally, when performing skin cancer testing using extracts such as ethosomal gel, which results in controlled release of curcumin, curcumin also has a longer half-life and maximum deposition in the deeper layers of the skin. Thus, it is able to completely eliminate melanoma cells³⁰. In addition, due to its pharmacological and therapeutic qualities, diosmin has the potential to be a flavonoid in the treatment of hepatocellular carcinoma, both in vivo and in vitro, in addition to its carcinogenic anticancer potential. Even so, further research into various types of cancer is still needed³¹.

In-depth studies of melanoma biology have had an influence on the consideration of melanoma as a therapy for one of the most resistant malignancies. There have been several options for treating metastatic melanoma in the last 30 years, so more extensive clinical studies have been conducted. These studies show that combination regimens, hormonal therapy, immunotherapy, single and combination agent chemotherapy, biochemotherapy, and targeted agent therapy do not always provide the best results²³.

CONCLUSION:

Based on cytotoxicity tests, *Chromolaena odorata* ethyl acetate extract has anti-skin cancer potential.

CONFLICT OF INTEREST:

There are no conflicts of interest declared by the authors of this study.

REFERENCES:

1. WHO (2020) Skin cancers. <https://www.who.int/uv/faq/skincancer/en/index1.htm>
2. Apalla Z, Nashan D, Weller RB, Castellsagué X. Skin cancer: epidemiology, disease burden, pathophysiology, diagnosis, and therapeutic approaches. *Dermatol Ther (Heidelb)*. 2017; 7(Suppl 1):5-19. doi: 10.1007/s13555-016-0165-y
3. Akila Victor, Muhammad Rukunuddin Ghalib. Detection of skin cancer cells – a review. *Research J.Pharm and Tech*; 2017; 10(11):4093-4098. doi: 10.5958/0974-360X.2017.00742.9
4. Scott G. Demonstration of melanosome transfer by a shedding microvesicle mechanism. *J Invest Dermatol*. 2012; 132(4):1073-4. doi: 10.1038/jid.2012.20
5. Nasti TH, Timares L. MC1R, eumelanin and pheomelanin: their role in determining the susceptibility to skin cancer. *Photochem Photobiol*. 2015; 91(1):188-200. doi: 10.1111/php.12335
6. Keerthana Karunakaran, Jansi Rani Malaiyappan, Rajini Raja Muniyan. Protein-protein interaction of mutated agouti signaling protein (asip) to melanocortin receptor 1 (mc1r) in melanoma skin cancer: an insilico study. *Research J. Pharm. and Tech* 2018; 11(9): 3913-3917. doi: 10.5958/0974-360X.2018.00718.7
7. Min Jeong Kim I, Seo Yeon Shin I, Nu Ri Song I, Sunoh Kim 2, Sang Ouk Sun I and Kyung Mok Park I. Bioassay-guided characterization, antioxidant, anti-melanogenic and anti-photoaging activities of pueraria thunbergiana l. leaf extracts in human epidermal keratinocytes (hacat) cells. *MDPI* 10. 2022; 2156. <https://doi.org/10.3390/pr10102156>
8. Singh M, Suman S, Shukla Y. New enlightenment of skin cancer chemoprevention through phytochemicals: in vitro and in vivo studies and the underlying mechanisms. *Biomed Res Int*. 2014; 2014:243452. doi: 10.1155/2014/243452
9. Qadir MI. Skin cancer: etiology and management. *Pak J Pharm Sci*. 2016; 29(3):999-1003
10. Aba, P.E., Joshua, P.E., Ezeonuogu, F.C., Ezeja, M.I., Omoja, V.U., Umeakuana, P.U. Possible anti-diarrhoeal potential of ethanol leaf extract of *Chromolaena odorata* in castor oil-induced rats. *J. Complement. Integr. Med*. 2015; 301–306. <https://doi.org/10.1515/jcim-2014-0033>
11. Femi Olawalea,b, Kolawole Olofinsanc, Opeyemi Iwaloyed. Biological activities of *chromolaena odorata*: A mechanistic review. *Elsevier*. 2021; <https://doi.org/10.1016/j.sajb.2021.09.001>
12. Hung, T.M., Cuong, T.D., Dang, N.H., Zhu, S., Long, P.Q., Komatsu, K., Min, B.S. Flavonoid glycosides from *chromolaena odorata* leaves and their in vitro cytotoxic activity. *Chem. Pharm. Bull.* 59. 2011; 129–131. <https://doi.org/10.1248/cpb.59.129>
13. Ajay, A., Kumar, R., Badhusha, S., Abhishek, K., Gowda, S.K., Ramesh, B. Pharmacological importance of *chromolaena odorata*: a review. *Int. J. Pharm. Drug Anal Vol* 9. 2021; 8–11. Doi:10.47957/ijpda.v9i1.452
14. Vijayaraghavan, K., Rajkumar, J., Seyed, M.A (2018). Phytochemical screening, free radical scavenging, and antimicrobial potential of *chromolaena odorata* leaf extracts against pathogenic bacterium in wound infectionsa multispectrum perspective. *Biocatal. Agric. Biotechnol.* 15, 103–112. 20. <https://doi.org/10.1016/j.bcab.2018.05.014>
15. Nath, L.R., Gorantla, J.N., Joseph, S.M., Antony, J., Thankachan, S., Menon, D.B., Sankar, S., Lankalapalli, R.S., Anto, R.J. Kaempferide, the most active among the four flavonoids isolated and characterized from *chromolaena odorata*, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv.* 5. 2015; 100912–100922. <https://doi.org/10.1039/C5RA19199H>
16. Mkhobongo B, Chandran R, Abrahamse H. In vitro photodynamic treatment modality for a375 melanoma cell line using a sulphonated aluminum phthalocyanine chloride-photosensitizer-gold nanoparticle conjugate. *Pharmaceutics*. 2022; 16;14(11):2474. doi: 10.3390/pharmaceutics14112474
17. Irma Colombo, Erico Sangiovanni, Roberta Maggio, et all. Hacat cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediator of Inflammation*. 2017; <https://doi.org/10.1155?2017/7435621>
18. Pastar I, Stojadinovic O, Yin NC, Ramirez H, Nusbaum AG, Sawaya A, Patel SB, Khalid L, Isseroff RR, Tomic-Canic M. Epithelialization in wound healing: a comprehensive review. *Adv Wound Care (New Rochelle)*. 2014; 3(7):445-464. doi: 10.1089/wound.2013.0473
19. Micallef L, Belaubre F, Pinon A, Jayat-Vignoles C, Delage C, Charveron M, Simon A. Effects of extracellular calcium on the growth-differentiation switch in immortalized keratinocyte hacat cells compared with normal human keratinocytes. *Exp Dermatol*. 2009; 18(2):143-51. doi: 10.1111/j.1600-0625.2008.00775.x
20. Pedrosa TDN, Catarino CM, Pennacchi PC, Assis SR, Gimenes F, Consolaro MEL, Barros SBM, Maria-Engler SS. A new reconstructed human epidermis for in vitro skin irritation testing. *Toxicol In Vitro*. 2017; 42:31-37. doi: 10.1016/j.tiv.2017.03.010
21. Yajarla, V.N.G., Nimmanapalli, R.P., Parikapandla, S., Gupta, G., Karnati, R. Anti-inflammatory, and anti-proliferative properties of *Chromolaena odorata* leaf extracts in normal and skin-cancer cell lines. *J. Herbs. Spices Med. Plants* 20. 2014; 359–371. doi:10.1080/10496475.2013.876698
22. Wilson VG. Growth and differentiation of hacat keratinocytes. *Methods Mol Biol*. 2014; 1195:33-41. doi: 10.1007/7651_2013_42
23. Tammineni Sreelatha, M. V. Subramanyam, M. N. Giri Prasad. A survey work on early detection methods of melanoma skin cancer. *Research J. Pharm. and Tech*. 2019; 12(5):2589-2596. doi: 10.5958/0974-360X.2019.00435.9
24. Micali G, Lacarrubba F, Nasca MR, Ferraro S, Schwartz RA. Topical pharmacotherapy for skin cancer: part ii. Clinical applications. *Journal of the American Academy of Dermatology*. 2014; 70(6):979.e1-12; quiz 9912. doi: 10.1016/j.jaad.2013.12.037.
25. Navdeep Singh, Shivi Sondhi, Sanyam Sharma, Dheeraj Singh, Vishal Koundal, Kamya Goyal, Shammy Jindal. Treatment of skin cancer by topical drug delivery of nanoparticles: a review. *Research Journal of Pharmacy and Technology*. 2021; 14(10):5589-8. doi: 10.52711/0974-360X.2021.00973
26. Zainab, Nurlailah, Muhammad Ikhwan Rizki. Identification of active compound and antibacterial activity against gram-positive and gram-negative bacteria of *chromolaena odorata* leaf extract. *Research Journal of Pharmacy and Technology*. 2022; 15(10):4720-6. doi: 10.52711/0974-360X.2022.00793
27. Nishita Gogia, Anandhi D, Kanaga G, Revathi K. In vitro evaluation of anti-mitotic activity of green synthesized silver nanoparticles from ethanolic extract of *chromolaena odorata*, *caesalpinia coriaria* (bark and leaves) using *allium cepa* roots. *Research Journal of Pharmacy and Technology*. 2021; 14(8):4107-0. doi: 10.52711/0974-360X.2021.00711
28. Eze FN, Jayeoye TJ. *Chromolaena odorata* (Siam weed): A natural reservoir of bioactive compounds with potent anti-fibrillogenic, antioxidative, and cytocompatible properties. *Biomed Pharmacother*. 2021;141:111811. doi: 10.1016/j.biopha.2021.111811.
29. Madhulika Pradhan, S Sribhuaneswari, D Karthikeyan, Sunita Minz, Pavani Sure, Atul N Chandu, Umesh Mishra, K Kamalakannan, A Saravanankumar, T Sivakumar. In-vitro cytoprotection activity of *foeniculum vulgare* and *helicteres isora* in cultured human blood lymphocytes and antitumour activity against b16f10 melanoma cell line. *Research J. Pharm. and Tech*. 2018; 1(4):450-452
30. Radha Krishna Kollipara, Vyshnavi Tallapaneni, Bharat Kumar Reddy Sanapalli, Vinoth Kumar G, Veera Venkata Satyanarayana Reddy Karri. Curcumin loaded ethosomal vesicular drug delivery system for the treatment of melanoma skin cancer. *Research Journal of Pharmacy and Technology*. 2019; 12(4):1783-1792. doi: 10.5958/0974-360X.2019.00298.1
31. Perumal S, Langeswaran K. Diosmin anti-tumour efficacious against hepatocellular carcinoma. *Research J. Pharm. and Tech*. 2020; 13(4):1707-1714. doi: 10.5958/0974-360X.2020.00308.X
32. Ghadge Ojaswi, Nadar Divya, Parmar Digna. Melanoma and its drug targets. *Research J. Pharm. and Tech*. 2016; 9(5): 562-570. doi: 10.5958/0974-360X.2016.00107.4