

67_RJPT_17_8_2024.

by TURNITIN NO REPOSITORY

Submission date: 24-Sep-2024 05:59PM (UTC+0530)

Submission ID: 2462929380

File name: 67_RJPT_17_8_2024.pdf (136.3K)

Word count: 3502

Character count: 19256

RESEARCH ARTICLE

2
**Growth Differentiation Factor -15 (GDF-15) Levels in Diabetic Conditions:
Animal Model**

Darmawansyih^{1,5*}, Asri Saleh², Rosdiana Natzir³, Ika Yustisia³, Firdaus Hamid⁴

¹Doctoral Program, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

²Chemistry Study Program, Science and Technology Faculty,
Universitas Islam Negeri Alauddin, Makassar, Indonesia.

³Department of Biochemistry, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

⁴Departement of Clinical Mikrobiology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

⁵Department of Medical, Faculty of Medicine and Health Science,
Universitas Islam Negeri Alauddin, Makassar, Indonesia.

*Corresponding Author E-mail: darmawansyih@uin-alauddin.ac.id

ABSTRACT:

Diabetes mellitus (DM) is a disease characterized by increased blood sugar levels, consisting of Diabetes Mellitus type I (DMT1) and Diabetes Mellitus type II (DMT2). Chronic increases in blood glucose levels will cause oxidative stress through the accumulation of Reactive Oxygen Species (ROS). Oxidative stress will induce the secretion of pro- and anti-inflammatory mediators, including Growth Differentiation Factor-15 (GDF-15). GDF-15 levels can be used to predict the risk and progression of DM disease. In exploring diabetes, several studies were carried out on experimental animal models of diabetes, both T1DM and 2DMT models. This research is an experimental study with a Pre and Post-Test Control Group Design approach using rat animals divided into four groups (I=Control group; II=Alloxan group; III=Streptozotocin group and IV=Streptozotocin and high fat diet group) with the aim of to see the levels of Growth Differentiation Factor-15 (GDF-15) in diabetic animal model. Based on this research, there was an increase in levels of Growth Differentiation Factor-15 (GDF-15) before and after diabetic animal model in the Streptozotocin group P value 0.010; and Streptozotocin with a High Fat Diet group with P value of 0.043 (significance value $P < 0.05$).

KEYWORDS: Diabetes mellitus, GDF-15, Oxidative stress.

INTRODUCTION:

Diabetes mellitus (DM) is a disease resulting from chronic metabolic disorders, characterized by blood sugar levels that exceed normal limits (hyperglycemic)¹, which can be caused either by impaired insulin secretion, insulin action, or both. Diabetes mellitus (DM) is disease of global concern². Type I Diabetes Mellitus (T1DM) is caused by impaired insulin secretion due to damage to pancreatic cells, and type II Diabetes Mellitus (T2DM) is caused by impaired insulin action^{3,4}.

In DM, there is an increase in blood glucose levels that exceed normal limits (hyperglycemic) over a long period of time (chronic), which will cause oxidative stress through the accumulation of Reactive Oxygen Species (ROS)⁵. In oxidative stress, there will be an increase in the production of free radicals such as malondialdehyde (MDA), which will react with tissue, cell proteins, and DNA nitrogen bases, which can cause cell injury, enzyme damage, changes in cell membranes, disruption of cell receptors, cell damage, DNA damage, which will ultimately cause disruption of insulin production and work. Diabetes is a defect in the body's ability to convert glucose (sugar) to energy^{3,6}. Oxidative stress in DM will also stimulate the body's inflammatory response in the form of the secretion of pro- and anti-inflammatory mediators, including Growth

Differentiation Factor-15 (GDF-15)⁶⁻⁹.

Growth Differentiation Factor-15 (GDF-15) is a member of the superfamily of Transforming Growth Factor- β (TGF- β) which is also called MIC-1 (macrophage inhibitory cytokine-1)¹⁰. Under normal conditions, GDF 15 has quite low levels in the blood and will increase if cellular stress occurs, such as heart failure, kidney failure, cancer, chronic inflammation, hyperglycemia, and obesity¹¹. GDF-15 levels can be used to predict the risk and progression of DM disease¹².

In exploring diabetes, several studies were carried out on experimental animal models of diabetes, both T1DM and 2DMT models¹³. There are several animal models of diabetes induction, namely the alloxan induction method, the streptozotocin induction method, or the streptozotocin induction method combined with a high-fat diet¹⁴.

MATERIALS AND METHODS:

This research is experimental research with a Pre and Post-Test Control Group Design approach using experimental mice divided into four groups, each consisting of 4-6 mice (The sample of mice used was calculated based on the Resource Equation Method Division of experimental animal groups: group I = Control group; Group II = Alloxan induction (dose 150 mg/KgBW_{i.p}); Group III = Streptozotocin induction (dose 45 mg/KgBW_{i.p}) and Group IV = Streptozotocin induction (dose 35 mg/KgBW) plus a high-fat diet/HFD (calorie source comes from 60% fat for 21 days), mice will be declared diabetic if the TGB level > 200 mg/dl. This study examines the Growth Differentiation Factor-15 (GDF-15) levels in diabetes animal models.

Blood from mice from groups I and II was taken 48 hours after induction, while groups III and IV were taken 72 hours after. Blood from mice was taken via the retrobulbar plexus. Temporary Blood Glucose levels (TBG) were checked using a one-touch brand glucometer, while GDF-15 levels were checked using Double Antibody Sandwich ELISA (Enzyme-Linked Immunosorbent Assay). TBG and GDF-15 level changes analysis in diabetes conditions were analyzed using the T-test (for normally distributed data) and the Wilcoxon test (for not normally distributed data).

RESULTS:

From the research results, it was found that in the control group, there was no difference between TGB levels and GDF 15 levels in the first and second sampling (P value for TGB = 0.186 and P value for GDF 15 = 0.68) with an average TGB pre 135 and average TGB post 120.25 (TGB within normal limits).

Meanwhile, the group of mice that received diabetes induction using three induction methods, namely the streptozotocin injection method with a dose of 45 mg/KgBW, the Alloxan injection method 150 mg/KgBW, and the STZ injection method 30 mg/KgBW + High-fat diet succeeded in increasing TGB levels. > 200 mg/dl and is said to be diabetes. In this study, the alloxan group had an average pre-diabetic TGB of 121.8 mg/dl and an average post-diabetic TGB of 294.2 mg/dl. The STZ group had an average pre-TGB level of 90.6 mg/dl and a post-TGB level of 334.6 mg/dl, while the STZ+HFD group had an average pre-TGB level of 106.8 mg/dl and an average post-TGB level of 307 mg. /dl. From these data, after statistical tests were carried out, it was found that there were differences in pre and post-diabetic TGB levels in all groups with a P value (streptozotocin group) of 0.000; P value (alloxan group) 0.003; P value (STZ+HFD group) 0.043 (significance value <0.05). Meanwhile, for GDF-15 levels in the control group, there was no statistical change in levels; for the diabetes group, the STZ group experienced changes in GDF-15 levels before and after diabetes, namely with a P value of 0.010 (significance value <0.05), while changes in GDF 15 levels in the alloxan group it was not significant with a P value of 0.677 (significance value <0.05) and in the STZ+HFD group there was a significant change in GDF 15 levels before and after diabetes with a P value of 0.043 ((significance value <0.05).

DISCUSSION:

Several induction methods are often used in experimental animal models of diabetes. Several previous studies have used alloxan for diabetes induction. Pei-pei Yin (2018) also used 150 mg/KgBW alloxan injection to induce diabetes in his research and proved that TGB levels could increase with this method¹⁵. Apart from that, Basma and Ramadan (2017) in their study also used alloxan to induce diabetes¹⁶. Administration of alloxan can cause type I diabetes mellitus through the mechanism of oxidative stress by ROS, which will cause damage to pancreatic beta cells. Damage to pancreatic beta cells by ROS has been associated with fragmentation of the cells' DNA, leading to stimulation of poly ADP-ribose polymerase 1, an enzyme that plays a vital role in DNA repair processes.¹⁷ Compared to other induction methods, alloxan is rarely used and is starting to be abandoned because it can cause kidney toxicity with a very narrow effective dose range. Besides, blood glucose levels in this induction model are multiphasic or unstable^{18,19}.

Compared to other induction methods, alloxan is rarely used and is starting to be abandoned because it can cause kidney toxicity with a very narrow effective dose range. Besides, blood glucose levels in this induction

model are multiphasic or unstable β cell damage, which will fail insulin secretion (a characteristic of T1DM). Meanwhile, a single medium dose of STZ (between 40-55 mg/KgBW) will cause mild insulin secretion failure that resembles the end stage of T2DM, while a single light dose of <35 mg/KgBW in normal diet mice does not show signs of hyperglycemia^(9,14). In this study, STZ was used at a dose of 45 mg/KgBW, which will provide a T2DM model.

Many previous studies have also used diabetes induction methods using STZ, including research by Naymul Karim et al. (2018) and SobayirilHidayati (2021)^{19,20}. STZ can cause diabetic conditions in experimental animal models through selective damage to pancreatic β cells, whose effects can be seen 72 hours after administration of STZ or depending on the dose administered. The toxic impact of STZ begins with the uptake of STZ into cells via the low-affinity glucose transporter-2 (GLUT-2) found in the cell membrane of the pancreas, which then causes cell damage. This condition will cause experimental animals to experience insulin deficiency, hyperglycemia, polydipsia, and polyuria^{21,22}.

The group that received low-dose STZ diabetes induction with a high-fat diet (HFD) also showed increased blood glucose levels. The STZ+HFD method is among the many popular T2DM induction methods used in animal experiments. It shows the same metabolic characteristics and natural history development as T2DM and can cause more stable and long-lasting hyperglycemia. The low STZ dose used in this study was 35 mg/KgBW combined with a high-fat diet with a calorie source of 60% fat. A single dose of STZ at a light dose of <35 mg/KgBW in mice on a regular diet did not show signs of hyperglycemia; however, if the light dose was combined with HFD, it was proven to induce diabetes²³. This group's average blood glucose level after induction was 307 mg/dl. This is by the target for successful induction, namely TGB levels > 200 mg/dl²⁴.

A high-fat diet will increase the circulation of free fatty acids (FFA). Elevated FFA may impair insulin sensitivity in muscle by inhibiting IRS/PI3 kinase autophosphorylation, which would decrease GLUT4 translocation to the plasma membrane. FFA can also increase serine phosphorylation of the Irs (Insulin Receptor) protein, thereby interfering with insulin signal transduction. HFD can also cause upregulation of inflammatory genes and the release of pro-inflammatory cytokines. In the liver, HFD will cause hepatic steatosis, a condition associated with insulin resistance in the liver. These conditions will cause the pathomechanism of T2DM²⁵.

GDF 15 is a stress-induced protein, which will increase in conditions of cell injury and if cells experience stress, such as in diabetes. In this study, it was proven that there was a significant increase in GDF 15 levels in diabetes conditions for the STZ group and STZ+HFD group, which represented the T2DM model. In T2DM conditions, there is low-grade inflammation that occurs chronically, causing accumulation of ROS production and giving rise to a complex stress response in the form of oxidative stress, protein glycation, inflammation, and hormonal changes, which can induce GDF 15 expression through the tumor suppressor protein P53 and the Early Growth transcription factor. Response Protein-1 (EGR-1). Apart from that, oxidative stress can also cause stress on the endoplasmic reticulum, which will cause ephosphorylation of eIF2 α (*Eukaryotic Translation Initiation Factor 2 α*) will increase the activation of ATF4 (Activating Transcription Factor 4), where ATF4, P53 and EGR-1 together increase the expression of GDF-15²⁵. Meanwhile, for the alloxan group, which represents the T1DM model, although there was an increase in GDF 15 levels, it was not statistically significant. The value of change in GDF 15 levels, which was not significant in the alloxan group, could be caused by the mechanism of action of alloxan in inducing T1DM, which takes place acutely through the direct destruction of pancreatic β cells. The diabetogenic agent alloxan causes necrosis and degeneration of pancreatic β cells in mice through an oxidative stress mechanism²⁶. In T1DM, damage to pancreatic β cells occurs, which will fail insulin secretion and hyperglycemia. Chronic hyperglycemia conditions will then cause ROS accumulation and a complex stress response. Blood sampling in this group was conducted 48 hours after induction, resulting in the necrosis and degeneration phase of pancreatic β cells. At the same time, the inflammatory reaction had not lasted long, so there had not been an accumulation of ROS, which could cause a complex stress response. Researchers suspect this is the cause of the increase in GDF 15 levels in the alloxan induction group, which was not significant.

The difference in mean TGB and GDF 15 in diabetic conditions between the induction groups also showed significance, with a P value of 0.000 for TGB levels and a P value of 0.003 for GDF 15 levels (significance value $P < 0.05$). Based on the different test results between groups, the post-hoc test was then continued to see which group was most significant for increasing TGB levels. From the results of the analysis, the three induction methods were significant. Meanwhile, the significance of the difference in mean GDF15 levels between groups in diabetic conditions was continued using Post Hoc Kruskal Wallis Analysis, and it was found that the STZ+HFD group had the highest increase

in GDF-15 levels. This is because the STZ+HFD model shows metabolic characteristics and natural history development that are more similar to T2DM than other induction methods so that it can cause more stable and long-lasting hyperglycemia. This condition will cause the accumulation of ROS, which then induces a complex stress response in the form of oxidative stress, protein glycation, inflammation, and hormonal changes, which can induce GDF 15 expression through the tumor suppressor protein P53 and the transcription factor *Early Growth Response Protein-1 (EGR-1)*. Apart from that, oxidative stress can also cause stress on the endoplasmic reticulum, which will cause phosphorylation of *eIF2 α* (*Eukaryotic Translation Initiation Factor 2 α*). *eIF2 α* will increase the activation of ATF4 (*Activating Transcription Factor 4*), where ATF4, P53 and EGR-1 together increase the expression of GDF-15^{27,28}.

In this research, there were several limitations, including the blood glucose level sex amined, where temporary glucose is strongly influenced by many factors, such as therats' irregulareating times. A histological examination of the pancreas was not carried out to see changes in pancreatic cell structure that could explain the progressing phase of DM disease. This study had no data on body weight before and after diabetes, which could be used to discuss changes in *Growth Differentiation Factor-15 (GDF-15)* levels.

CONCLUSIONS:

Changes in Temporary Blood Glucose (TGB) levels before and after the diabetic animal model condition were significant in all induction groups. Meanwhile, changes in *Growth Differentiation Factor-15 (GDF-15)* levels before and after the diabetic animal model were significant in the Streptozotocin induction and Streptozotocin groups with a high-fat diet. Meanwhile, there was no significant change in *Growth Differentiation Factor-15cs (GDF-15)* levels in the alloxan group before and after diabetic conditions

6

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this investigation.

ACKNOWLEDGMENTS:

The researcher expresses her highest gratitude to Doctoral Program Faculty of Medicine Universitas Hasanuddin and Department of Medical, Faculty of Medicine and Health Science, Universitas Islam Negeri Alauddin, Makassar, Indonesia.

REFERENCES:

1. Vinit Kumar Sharma, Madan Kaushik, Deepika Rani. Role of Ayurveda in the Management of Diabetes Mellitus: A Comprehensive Review to explore the awareness. Research

Journal of Pharmacy and Technology. 2023; 16(11): 5528-1. doi: 10.52711/0974-360X.2023.00894

2. Sanusi Sani, ZayyanuShitu, Faatimah Bah, Tijjani Salihu Shinkafi, San San Oo, Zaw Aung, Vidya Bhagat. Overview of Diabetes, Management, and Pharmacotherapy Updates a Review. Research J. Pharm. and Tech. 2019; 12(5): 2597-2602. doi: 10.5958/0974-360X.2019.00436.0
3. M. Sujatha Kumari, M. Kiran Babu, Rehana Sulthana, M. Srinivas, Ch. Prasanthi. Diabetes Mellitus: Present status and Drug therapy Updates. Research J. Pharm. and Tech. 2014; 7(1): 84-94.
4. Samidha Kamtekar, Vrushali Keer. Management of Diabetes: A Review. Research J. Pharm. and Tech. 2014 7(9): 1065-1072.
5. Imad S. Mahmoud, Khalil I. Altaif, Abdulrasool M. Wayyes, Watheq Mohammed Al – Jewari, Iyad A. Hailat, Moeen F. Dababneh. Diabetic Mellitus as a Predisposing factor in enhancing infections by Candida species and their Antifungal susceptibilities. Research Journal of Pharmacy and Technology. 2023; 16(11): 5189-2. doi: 10.52711/0974-360X.2023.00841
6. S.V. Rupashri, Gheena. S. Recent Advances in Diabetes Research. Research J. Pharm. and Tech. 2016; 9(10): 1806-1808. doi: 10.5958/0974-360X.2016.00366.8
7. Chatki PK, Tabassum S. Analytical Methods of Dihydropyridines Based Calcium Channel Blockers-Amlodipine, Lacidipine, Isradipine, Nifedipine, Felodipine, Cilnidipine and its related formulations: A Review. Asian Journal of Research in Chemistry. 2021; 14(3): 221-234. <https://doi.org/10.52711/0974-4150.2021.00039>
8. Suresh Kumar Sutrakar, Drutpal Singh Baghel. Biochemical Parameters Variations in Type-II Diabetes Mellitus: Special Reference in Rewa Region. Asian J. Research Chem. 2014; 7(10): 877-881.
9. Veena Prabavathy J, Sangeetha. R. Stress Induced Type 2 Diabetes Mellitus among Industrial Workers – A Review. Research J. Pharm. and Tech. 2019; 12(1): 396-402.
10. Roaa Hatem Al Gburi, Raid D. Hashim, Hayder Ahmed Kadhim, Şevki Adam, Haider Abdulkareem Almashhadani. Correlation of Angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers with serum GDF-15 in a group of hypertensive Iraqi patients. Research Journal of Pharmacy and Technology. 2022; 15(8): 3466-0. doi: 10.52711/0974-360X.2022.00580
11. Prawitasari DS. Diabetes Melitus dan Antioksidan. 2019; 1(1): 47-51.
12. Al-kuraishy HM, Batiha GE. Metformin and growth differentiation factor 15 (GDF15) in type 2 diabetes mellitus : A hidden treasure. 2022; (November): 806-14.
13. Ozougwu JC. The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. 2017; (January 2013).
14. Mandal M, Varghese A, Gaviraju VK, Talwar SN, Malini SS. Impact of hyperglycaemia on molecular markers of oxidative stress and antioxidants in type 2 diabetes mellitus. 2019; 215-22.
15. Shawki HA, Elzehery R, Shahin M, Abo EM. Evaluation of some oxidative markers in diabetes and diabetic retinopathy. Diabetol Int [Internet]. 2021; 12(1): 108-17. Available from: <https://doi.org/10.1007/s13340-020-00450-w>
16. Bahgat MM, Ibrahim DR. Proinflammatory cytokine polarization in type 2 diabetes. 2020; 170-5.
17. Ahluwalia TS, Kilpeläinen TO, Singh S, Rossing P. Editorial: Novel biomarkers for type 2 diabetes. Front Endocrinol (Lausanne). 2019; 10(SEP): 1-3.
18. Tsalamandris S, Antonopoulos AS, Oikonomou E, Papamikroulis G, Vogiatzi G. Risk Factors and Cardiovascular Disease Prevention The Role of Inflammation in Diabetes : Current Concepts and Future Perspectives. 2019; 50-9.
19. A WN, Z WM. Sample Size Calculation in Animal Studies Using Resource Equation Approach. 2017; 24(5): 101-5.
20. Karim N, Jeenduang N, Tangpong J. Anti-Glycemic and Anti-Hepatotoxic Effects of Mangosteen Vinegar Rind. 2018; 68(2): 163-9.
21. Solihatil Hidayati. Antidiabetic Activity of Peperomia pellucida In

- Streptozotocin-Induced Diabetic Mice. 2021; 7(2): 120–30.
22. Enriched C. Hypoglycemic Effects in Alloxan-Induced Diabetic Rats of the Phenolic Extract from Mongolian Oak. 2018;
 23. Ramadan BK, Schaalán MF, Tolba AM. Hypoglycemic and pancreatic protective effects of *Portulaca oleracea* extract in alloxan induced diabetic rats. *BMC Complement Altern Med* [Internet]. 2017; 1–10. <http://dx.doi.org/10.1186/s12906-016-1530-1>
 24. Macdonald O, Mohammed A. ScienceDirect Alloxan-induced diabetes , a common model for evaluating the glycemic-control potential of therapeutic compounds and plants extracts in experimental studies. *Medicina (B Aires)* [Internet]. 2018; 1–10. <https://doi.org/10.1016/j.medic.2018.02.001>
 25. Furman BL. Streptozotocin-Induced Diabetic Models in Mice and Rats. *Curr Protoc*. 2021;1.
 26. Gheibi S, Kash K, Ghasemi A. *Biomedicine and Pharmacotherapy A practical guide for induction of type-2 diabetes in rat: Incorporating a high-fat diet and streptozotocin*. 2017; 95(24): 605–13.
 27. Baek SJ, Eling T. Growth Differentiation Factor 15 (GDF15): A Survival Protein with Therapeutic Potential in Metabolic Diseases. 2020;1:46–58.
 28. Wt K, Fs C, Ts Q, Ew AN, Nh K. High-fat diet and streptozotocin in the induction of type 2 diabetes mellitus : a new proposal. 2019; 91: 1–14.
 29. Adela R, Banerjee SK. GDF-15 as a target and biomarker for diabetes and cardiovascular diseases: A translational prospective. *J Diabetes Res*. 2015; 2015.
 30. Wang D, Day EA, Townsend LK, Djordjevic D, Jørgensen SB, Steinberg GR. GDF15: emerging biology and therapeutic applications for obesity and cardiometabolic disease. *Nat Rev Endocrinol* [Internet]. 2021; 17(October). Available from: <http://dx.doi.org/10.1038/s41574-021-00529-7>

ORIGINALITY REPORT

14%

SIMILARITY INDEX

11%

INTERNET SOURCES

10%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	www.rjptonline.org Internet Source	2%
2	rjptonline.org Internet Source	2%
3	"Biomarkers in Diabetes", Springer Science and Business Media LLC, 2023 Publication	1%
4	Submitted to Universitas Indonesia Student Paper	1%
5	Osasenaga Macdonald Ighodaro, Abiola Mohammed Adeosun, Oluseyi Adeboye Akinloye. "Alloxan-induced diabetes, a common model for evaluating the glycemetic-control potential of therapeutic compounds and plants extracts in experimental studies", <i>Medicina</i> , 2018 Publication	1%
6	orcid.org Internet Source	1%
7	estd.perpus.untad.ac.id	

Internet Source

1 %

8

www.mdpi.com

Internet Source

1 %

9

www.thieme-connect.com

Internet Source

1 %

10

Submitted to Jawaharlal Nehru Technological University

Student Paper

1 %

11

coek.info

Internet Source

1 %

12

link.springer.com

Internet Source

1 %

13

Muthmainah, Hanik Badriyah Hidayati, Budi Yanti. "Improving Health for Better Future Life: Strengthening from Basic Science to Clinical Research", CRC Press, 2023

Publication

1 %

14

www.sciencegate.app

Internet Source

1 %

Exclude quotes

Off

Exclude matches

< 15 words

Exclude bibliography

On