


**REVIEWER PADA 3 (TIGA) JURNAL  
INTERNASIONAL BEREPUTASI:**

- 1. Journal Plos One - Q1 : SJR 0,84  
United States**
- 2. Journal of health science and medical  
research - Q4 : SJR 0,16  
Thailand**
- 3. Belitung Nursing Journal -  
Q2 : SJR 0,45  
Indonesia**




1.Jurnal Plus One

Q1 : SJR 0,84

United States



Search plos one



99+

Tulis

Mail

Kotak Masuk 1.723

Berbintang

Chat

Ditunda











Terkirim


Meet

Draf 2


Selengkapnya


Label



 Summarize

Invitation to apply to join PLOS ONE Editorial Board Eksternal Kotak Masuk x

 **PLOS Editorial Board Management** <edboardmgmt@plos.org>  
kepada saya

 Menerjemahkan... X

Dear Dr. Risnah

This is Melissa Quinton with *PLOS ONE*. I'd like to take this opportunity to ask whether you are interested in le membership with *PLOS ONE*. We are currently seeking additional Academic Editors in various areas and are r apply to join the *PLOS ONE* Editorial Board.

The role description is on our [website](#) but in summary, you would be responsible for overseeing the peer review assessing whether they should go out for peer review, securing reviewers, and issuing decisions. Workload is ' typically expect Editors to handle two manuscripts per month, though this varies according to subject area, sub commitments.

Candidates that qualify for the Editorial Board:

- Have a strong publication record (10 or more peer-reviewed published articles)
- Are Principal Investigators or oversee research at an equivalent level
- Are affiliated with an academic, governmental, or relevant nonprofit research institution

If you are interested in this opportunity, please follow this [link](#) to provide your information so that we may asses questions, please do not hesitate to reach out.

I look forward to hearing from you.

99+

Tulis

Mail

Chat

Meet

Kotak Masuk

Berbintang

Ditunda

Terkirim

Draf

Selengkapnya

1.723

2

←

📁

🕒

🗑️

📧

🕒

🔄

📎

📄

⋮

✦ Summarize

📧

Invitation to re-review a revised paper - PLOS ONE PONE-D-21-33982 [EMID:21475775a2bff60f]

Eksternal Kotak Masuk x

👤

**PLOS ONE** <em@editorialmanager.com>  
kepada saya

📎

Menerjemahkan...

×

Label

Dear Dr. Risnah,

Thank you for your previous review for *PLOS ONE* submission "Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia." (PONE-D-21-33982R1). Your comments were shared with the authors, who have now submitted a revised submission. We would be grateful if you could review this revised submission and evaluate whether the authors have addressed your concern for publication. At the end of the submission PDF, you will find a response to previous reviewer comments from the author manuscript with tracked changes, if available.

**Manuscript #:** PONE-D-21-33982R1

**Title:** Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.


**Authors:** Addis Eyeberu; Tamirat Getachew; Adera Debela; Bikila Balis; Bajirond Eshetu; Sinetibet Mesfin; Habtamu Birhan; Meron Degefa; Addisu Alemu; Merga Dheressa; Yadeta Dessie

**Article type:** Research Article

The abstract is appended below, along with more detailed information about *PLOS ONE* and its [publication criteria](#). Please refer to [the different article types that we publish](#).



Q plos one



99+

Tulis



✦ Summarize

Mail

Kotak Masuk	1.723
-------------	-------

Berbintang

Chat

Ditunda

## Terkirim

## Meet

**Draf**

2

▼ Selengkapnya

Label

PLOS ONE: Agreement to Review PONE-D-21-33982R1 - [EMID:31de5f59413bafbfdea] (PLOS ONE: Persetujuan untuk Meninjau PONE-D-21-33982R1 - [EMID:31de59413bafbfdea]) Eksternal Kotak Masuk x

Eksternal      Kotak Masuk x



PLOS ONE &lt;em@editorialmanager.com&gt;

kepada saya



Inggris → Indonesia

[Tampilkan versi asli](#)



Jul	Review Due	Ag
<b>18</b>	Waktu Sen, 18 Jul 2022 (WITA)	Sen
Sen	Siapa PLOS ONE*	Tide
	<a href="#">Tambah ke kalender »</a>	Sep
		Tide

**Nomor Naskah:** PONE-D-21-33982R1

**Judul Naskah:** Pemanfaatan Analgesia Persalinan Farmakologis: Survei Penyedia Layanan Obstetri di Etiopia Timur.

Yth. Dr. Risnah,

99+

Tulis

Mail

Kotak Masuk

1.723

Chat

Berbintang

Ditunda

Terkirim

Meet

Draf

2

Selengkapnya

Label

plos one

←

📧

🕒

🗑️

📧

🕒

🔄

📧

📄

⋮

✦ Summarize

Review assignment for PONE-D-21-33982R1 is due soon - [EMID:4a6e1eaa552cc25c] (Tugas peninjauan untuk PONE-D-21-33982R1 segera jatuh tempo - [EMID:4a6e1eaa552cc25c]) Eksternal Kotak Masuk x

**PLOS ONE** <em@editorialmanager.com>  
kepada saya

Inggris → Indonesia  
[Tampilkan versi asli](#)

PONE-D-21-33982R1

Pemanfaatan Analgesia Persalinan Farmakologis: Survei Penyedia Layanan Obstetri di Etiopia Timur.  
PLOS SATU

Yth. Dr. Risnah,

Terima kasih telah setuju untuk mengirimkan tinjauan pada naskah PLOS ONE "Pemanfaatan Analgesia Persalinan Farmakologis: Survei Penyedia Layanan Obstetri di Etiopia Timur..". Sebagai pengingat, tinjauan Anda harus diserahkan paling lambat tanggal 18 Juli 2022 pukul 17.00 WIB ke <https://www.editorialmanager.com/pone/>.

Kami memahami bahwa pandemi COVID-19 dapat memengaruhi kemampuan Anda untuk mengerjakan tugas ini. PLC memberikan perpanjangan waktu untuk tugas pengulas. Silakan kirim email kepada kami di [plosone@plos.org](mailto:plosone@plos.org) jika Anda memerlukan lebih banyak waktu.

Jika Anda tidak dapat mengirimkan komentar Anda pada tanggal jatuh tempo yang disebutkan, harap diperhatikan bahwa kami akan memberikan keputusan berdasarkan evaluasinya sendiri terhadap naskah atau tinjauan yang diterima dalam waktu 14 hari.

<https://mail.google.com/mail/u/0/#search/plos+one/FMfcgzGpGwISzFpJKzFKwGqvFCKdmBjx>

1/1



Q plos one



99+

Tulis



✦ Summarize

Mail

**Kotak Masuk**

1.723

Chat

Berbintang

Ditunda

Terkirim

## Meet

**Draf**

2

▼ Selengkapnya

Label



PLOS ONE &lt;plosjournals@plos.org&gt;

kepada saya



Inggris → Indonesia

[Tampilkan versi asli](#)



Risnah yang terhormat, Risnah

Terima kasih telah menghubungi PLOS ONE. Kami akan membalas pertanyaan Anda sesegera mungkin.

Sementara itu, silakan lihat tautan berikut untuk informasi lebih lanjut tentang proses kami:

Pesan kepada komunitas kami mengenai COVID-19 <https://blogs.plos.org/plos/2020/03/a-message-to-our-community->

Pedoman Pengajuan <http://journals.plos.org/plosone/s/submission-guidelines>

Pedoman Peninjau <http://journals.plos.org/plosone/s/reviewer-guidelines>

Kriteria Publikasi <http://journals.plos.org/plosone/s/criteria-for-publication>

Proses Editorial dan Tinjauan Sejawat <http://journals.plos.org/plosone/s/editorial-and-peer-review-process>

Video Pendek PLOS ONE [https://www.youtube.com/playlist?list=PL\\_O2Hm19V2gEUZoyf7J4nva9W9u1XhJXN](https://www.youtube.com/playlist?list=PL_O2Hm19V2gEUZoyf7J4nva9W9u1XhJXN)

Kami menghargai perhatian Anda dan akan segera menghubungi Anda kembali.



Q plos one



99+

Tulis



✦ Summarize

Mail

**Kotak Masuk**

1.723

Chat

Berbintang

Ditunda

Terkirim

## Meet

**Draf**

2

▼ Selengkapnya

Label



Review Assignment for PONE-D-21-33982R1 is Now Due - [EMID:7d694bd95dbbe18e] (Tugas Peninjauan untuk PONE-D-21-33982R1 Sudah Harus Diserahkan) [EMID:7d694bd95dbbe18e]

Eksternal      Kotak Masuk x

**PLOS ONE** [em@editorialmanager.com](mailto:em@editorialmanager.com)  
kepada saya

 Inggris → Indonesia  
[Tampilkan versi asli](#)



PONE-D-21-33982R1

Pemanfaatan Analgesia Persalinan Farmakologis: Survei Penyedia Layanan Obstetri di Etiopia Timur.

Tuan, Addis Eyeberu

PLOS SATU

Yth. Dr. Risnah,

Terima kasih telah setuju untuk mengirimkan tinjauan naskah PLOS ONE 'Pemanfaatan Analgesia Persalinan Farmak di Etiopia Timur..' Anda setuju untuk meninjau naskah pada 08 Jul 2022 08:47PM dan tinjauan lengkap Anda harus disi 11:59PM EST.

Kami memahami bahwa pandemi COVID-19 dapat memengaruhi kemampuan Anda untuk mengerjakan tugas ini. PLC memberikan perpanjangan waktu untuk tugas pengulas. Silakan kirim email kepada kami di [plosone@plos.org](mailto:plosone@plos.org) jika Anc

Kami mencatat bahwa saat ini status tinjauan Anda adalah 'tersimpan sebagian'. Kami akan berterima kasih jika Anda :





✦ Summarize

Eksternal      Kotak Masuk x

**PLOS ONE** [em@editorialmanager.com](mailto:em@editorialmanager.com)  
kepada saya



Kami memahami bahwa pandemi COVID-19 dapat memengaruhi kemampuan Anda untuk mengerjakan tugas ini. PLC memberikan perpanjangan waktu untuk tugas pengulas. Silakan kirim email kepada kami di [plosone@plos.org](mailto:plosone@plos.org) jika Anc



99+

Tulis

Mail

Kotak Masuk 1.723

Chat

Ber bintang

Ditunda

Terkirim

Meet

Draf 2

Selengkapnya

Label

plos one

<>📧🕒🗑️📧🕒🔄📧📄⋮

✦ Summarize

PLOS ONE

<em@editorialmanager.com>

kepada saya

Inggris → Indonesia

[Tampilkan versi asli](#)

⚙️

PONE-D-21-33982R1

Pemanfaatan Analgesia Persalinan Farmakologis: Survei Penyedia Layanan Obstetri di Etiopia Timur.

Yth. Dr. Risnah,

Terima kasih telah mempertimbangkan undangan untuk meninjau naskah yang dirujuk di atas. Dalam upaya untuk mer kepada para penulis, kami sedang berupaya untuk memajukan naskah ini ke tahap berikutnya dari proses penyuntinga tidak akan diperlukan lagi. Kami mohon maaf atas ketidaknyamanan yang mungkin ditimbulkan dan terima kasih atas v pertanyaan, silakan hubungi kami di [plosone@plos.org](mailto:plosone@plos.org).

Semoga sukses,

PLOS SATU

[alamat email: plosone.org](mailto:plosone@plos.org)



✦ Summarize

Label



Sungguh-sungguh,  
Shein Ei Cho  
Rekan Peneliti Editorial Senior, Layanan Dewan Editorial, PLOS  
Atas nama  
Melissa Quinton  
Manajer Program Perekrutan, PLOS  
Yth. Dr. Risnah ,



✕ 𠄎

▼



✦ Summarize

PLOS ONE: A decision has been made on PONE-D-21-33982R1 - [EMID:c559b0ad9c3f1927] (PLOS ONE: Keputusan telah dibuat pada F 33982R1 - [EMID:c559b0ad9c3f1927]) Eksternal Kotak Masuk x

Eksternal      Kotak Masuk x

2



R



Inggris → Indonesia  
[Tampilkan versi asli](#)



Pemanfaatan Analgesia Persalinan Farmakologis: Survei Penyedia Layanan Obstetri di Etiopia Timur.  
PLOS SATU

Anda juga dapat mengakses komentar ulasan dan surat keputusan dengan masuk ke Editorial Manager sebagai memerlukan bantuan terkait hal ini!



Gmail

Q plos one



99+

Tulis

Mail

**Kotak Masuk**

1.723

Chat

Berbintang

Ditunda

## Terkirim

## Meet

**Draf**

2

▼ Selengkapnya

Label



✦ Summarize

Utilization of pharmacological labour analgesia: a survey of obstetric c  
in eastern Ethiopia has been published (Pemanfaatan analgesia persal  
farmakologis: survei penyedia layanan kebidanan di Ethiopia timur tela  
diterbitkan) Eksternal Kotak Masuk x

Eksternal

Kotak Masuk x



**Web of Science researcher profiles** [noreply@webofscience.com](mailto:noreply@webofscience.com) [lewat](#) [amazonses.com](#)  
kepada saya



Inggris → Indonesia

[Tampilkan versi asli](#)



 [Logo Perpustakaan Umum Ilmu Pengetahuan](#)

Risnah Risnah yang terhormat,

Kami menulis untuk memberi tahu Anda bahwa [Pemanfaatan analgesia persalinan farmakologis: survei penyedia perawatan obstetrik di Ethiopia timur](#) , yang Anda t untuk Plos One, diterbitkan pada tanggal 10 September oleh International Health

# Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.

--Manuscript Draft--

<b>Manuscript Number:</b>	PONE-D-21-33982
<b>Article Type:</b>	Research Article
<b>Full Title:</b>	Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.
<b>Short Title:</b>	Utilization of Pharmacological Labour Analgesia
<b>Corresponding Author:</b>	Addis Eyeberu Haramaya University College of Health Sciences: Haramaya University College of Health and Medical Sciences harar, ETHIOPIA
<b>Keywords:</b>	Utilization; Analgesia; labor pain; pharmacological labor pain management; Harar
<b>Abstract:</b>	<p><b>Background</b></p> <p>Despite a crucial role played by Obstetric care providers (OCPs) in providing pain relief methods during labor, the evidence suggests that there is an ample number of women in Ethiopia failed to have access to pain relief methods. Moreover, to the best knowledge of the investigators concerned, evidence is scarce regarding the utilization of pharmacological labor pain management methods in the study area. Thus, this study was carried out to determine the practice of pharmacological labor pain management methods among obstetric caregivers in Harari regional state health facilities, Ethiopia.</p> <p><b>Method</b></p> <p>A facility-based cross-sectional study was conducted from May 20 to June 10, 2021, in Harari regional state health facilities, Ethiopia. All obstetric caregivers in Harari regional state health facilities were included in the study. Data were collected using a structured questionnaire adapted from the previous studies. The data were entered into Epi-data version 3.1 statistical software. Statistical analysis was carried out by using SPSS for windows version 22. Bivariate and multivariate analyses were employed to determine the association between independent variables and the outcome variable.</p> <p><b>Result</b></p> <p>The overall utilization of pharmacological labor pain relief methods was 50.9% [(95% CI (45.8,55.8)]. The main barriers for laboring mothers receiving pain relief include high patient flow (54.4%), the limited number of staff (51.9%), and lack of knowledge and skill (32.9%). Being female [AOR=2.07, 95% CI (1.31,3.26)], not having training [AOR=1.60, 95%CI (1.02,2.53)], having unfavorable attitude [AOR=9.1, 95%CI: (5.34,15.35)] were the predictors of utilization of pharmacological labor pain management methods.</p> <p><b>Conclusion</b></p> <p>The study revealed that the overall prevalence of utilization of pharmacological labor pain management methods was low. In general, this study identified that variables like sex, training, and attitude of care providers were factors associated with utilization of pharmacological labor pain management methods. The health facilities should give the training to enhance the skill of obstetric care providers and equip themselves with sufficient analgesics drugs.</p>
<b>Order of Authors:</b>	Addis Eyeberu
	Tamirat Getachew
	Adera Debela

	Bikila Balis
	Bajirond Eshetu
	Sinetibet Mesfin
	Habtamu Bekele
	Dawit Tamiru
	Getahun Tiruye
	Meron Degefa
	Addisu Alemu
	Merga Dheressa
	Yadeta Dessie
<b>Additional Information:</b>	
<b>Question</b>	<b>Response</b>
<p><b>Financial Disclosure</b></p> <p>Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the <a href="#">submission guidelines</a> for detailed requirements. View published research articles from <a href="#">PLOS ONE</a> for specific examples.</p> <p>This statement is required for submission and <b>will appear in the published article</b> if the submission is accepted. Please make sure it is accurate.</p> <p><b>Unfunded studies</b> Enter: <i>The author(s) received no specific funding for this work.</i></p> <p><b>Funded studies</b> Enter a statement with the following details:</p> <ul style="list-style-type: none"> <li>• Initials of the authors who received each award</li> <li>• Grant numbers awarded to each author</li> <li>• The full name of each funder</li> <li>• URL of each funder website</li> <li>• Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?</li> <li>• <b>NO</b> - Include this sentence at the end of your statement: <i>The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</i></li> <li>• <b>YES</b> - Specify the role(s) played.</li> </ul>	<p>The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</p>



<p>* typeset</p> <p><b>Competing Interests</b></p> <p>Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any <a href="#">competing interests</a> that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests.</p> <p>This statement is <b>required</b> for submission and <b>will appear in the published article</b> if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.</p> <p>View published research articles from <a href="#">PLOS ONE</a> for specific examples.</p>	<p>The authors have declared that no competing interests exist.</p>
<p><b>NO authors have competing interests</b></p> <p>Enter: <i>The authors have declared that no competing interests exist.</i></p> <p><b>Authors with competing interests</b></p> <p>Enter competing interest details beginning with this statement:</p> <p><i>I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]</i></p>	
<p>* typeset</p> <p><b>Ethics Statement</b></p> <p>Enter an ethics statement for this submission. This statement is required if the study involved:</p> <ul style="list-style-type: none"> <li>• Human participants</li> <li>• Human specimens or tissue</li> </ul>	<p>Ethical clearance was taken from the Institutional Health Research Ethics Review Committee (IHRERC) of the College of Health and Medical Sciences of Haramaya University. In addition, permission to proceed was obtained from all public hospitals. And also, informed voluntary written and signed consent was obtained from each participant.</p>

- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below.

Consult the [submission guidelines](#) for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

### Format for specific study types

#### Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

#### Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

#### Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

#### Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare exceptions to address legal and ethical concerns. See the [PLOS Data Policy](#) and [FAQ](#) for detailed information.

Yes - all data are fully available without restriction

A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and **will be published in the article**, if accepted.

**Important:** Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

**Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.**

- If the data are **held or will be held in a public repository**, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: *All XXX files are available from the XXX database (accession number(s) XXX, XXX).*
- If the data are all contained **within the manuscript and/or Supporting Information files**, enter the following:  
*All relevant data are within the manuscript and its Supporting Information files.*
- If neither of these applies but you are able to provide **details of access elsewhere**, with or without limitations, please do so. For example:

*Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for access to confidential data.*

*The data underlying the results presented in the study are available from (include the name of the third party*

All relevant data are within the manuscript and its Supporting Information files.

<p><i>and contact information or URL).</i></p> <ul style="list-style-type: none"><li>• This text is appropriate if the data are owned by a third party and authors do not have permission to share the data.</li></ul> <p>* typeset</p>	
Additional data availability information:	

# Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.

Addis Eyeberu<sup>1\*</sup>, Tamirat Getachew<sup>1</sup>, Adera Debela<sup>1</sup>, Bikila Balis<sup>1</sup>, Bajirond Eshetu<sup>1</sup>, Sinetibet Mesfin<sup>1</sup>, Habtamu Bekele<sup>1</sup>, Dawit Tamiru<sup>1</sup>, Getahun Tiruye<sup>1</sup>, Meron Degefa<sup>1</sup>, Addisu Alemu<sup>2</sup>, Merga Dheressa<sup>1</sup>, Yadeta Dessie<sup>2</sup>

Author's affiliation

1. School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar.
2. School of Public Health, College of Health and Medical Sciences, Haramaya University, Harar.

Authors email address

AE- [addiseyeberu@gmail.com](mailto:addiseyeberu@gmail.com) , TG- [tamirget@gmail.com](mailto:tamirget@gmail.com) , AD- [aksanadera62@gmail.com](mailto:aksanadera62@gmail.com) , BB- [bik.balis2008@gmail.com](mailto:bik.balis2008@gmail.com) , BE- [bajeshetu@gmail.com](mailto:bajeshetu@gmail.com) , SM- [sine3686@gmail.com](mailto:sine3686@gmail.com) , HB- [habeti.bekele@gmail.com](mailto:habeti.bekele@gmail.com) , DT- [dawittamru5@gmail.com](mailto:dawittamru5@gmail.com) , GT- [gtiruye@gmail.com](mailto:gtiruye@gmail.com) , MD- [merrydegefa1@gmail.com](mailto:merrydegefa1@gmail.com) , AA: [alemuadisu789@gmail.com](mailto:alemuadisu789@gmail.com) , MD- [mderesa@yahoo.com](mailto:mderesa@yahoo.com) , YD: [yad\\_de2005@yahoo.com](mailto:yad_de2005@yahoo.com)

*Corresponding author information.*

- Addis Eyeberu
- [addiseyeberu@gmail.com](mailto:addiseyeberu@gmail.com) : Institutional email: [Addis.Eyeberu@haramaya.edu.et](mailto:Addis.Eyeberu@haramaya.edu.et)
- Haramaya University College of Health and Medical Science, Harar, Ethiopia
- ORCID: <https://orcid.org/0000-0002-3147-3770>; Institutional link: [www.haramaya.edu.et](http://www.haramaya.edu.et)
- P.O. BOX 138 Dire Dawa, Ethiopia

## Abstract

**Background:** Despite a crucial role played by Obstetric care providers (OCPs) in providing pain relief methods during labor, the evidence suggests that there is an ample number of women in Ethiopia failed to have access to pain relief methods. Moreover, to the best knowledge of the investigators concerned, evidence is scarce regarding the utilization of pharmacological labor pain management methods in the study area. Thus, this study was carried out to determine the practice of pharmacological labor pain management methods among obstetric caregivers in Harari regional state health facilities, Ethiopia.

**Method:** A facility-based cross-sectional study was conducted from May 20 to June 10, 2021, in Harari regional state health facilities, Ethiopia. All obstetric caregivers in Harari regional state health facilities were included in the study. Data were collected using a structured questionnaire adapted from the previous studies. The data were entered into Epi-data version 3.1 statistical software. Statistical analysis was carried out by using SPSS for windows version 22. Bivariate and multivariate analyses were employed to determine the association between independent variables and the outcome variable.

**Result:** The overall utilization of pharmacological labor pain relief methods was 50.9% [(95% CI (45.8,55.8)]. The main barriers for laboring mothers receiving pain relief include high patient flow (54.4%), the limited number of staff (51.9%), and lack of knowledge and skill (32.9%). Being female [AOR=2.07, 95% CI (1.31,3.26)], not having training [AOR=1.60, 95%CI (1.02,2.53)], having unfavorable attitude [AOR=9.1, 95%CI: (5.34,15.35)] were the predictors of utilization of pharmacological labor pain management methods.

**Conclusion:** The study revealed that the overall prevalence of utilization of pharmacological labor pain management methods was low. In general, this study identified that variables like sex, training, and attitude of care providers were factors associated with utilization of pharmacological labor pain management methods. The health facilities should give the training to enhance the skill of obstetric care providers and equip themselves with sufficient analgesics drugs.

**Keywords:** Utilization, Analgesia, labor pain, pharmacological labor pain management, Harar

## INTRODUCTION

Pain during labor and childbirth is a unique and the most severe pain that a woman ever has to face. The extent to which a woman feels in control of pain during labor is an indicator of maternal emotive wellbeing in childbirth [1]. Pharmacological labor pain management methods include oral tablets (paracetamol, codeine, or tramadol), inhalations, and injectable analgesics [2].

Pain related to childbirth is present during pregnancy, labor, and post-partum periods [3]. The satisfaction of childbirth depends on the care provided to the parturient during labor and delivery [4]. Evidence showed that 85.1% of the patients want their labor pain to be relieved, but 38.9% did receive analgesia during labor [5].

Even if Obstetric care providers (OCPs) have an imperative role in providing pain relief options during labor, the majority of women in Ethiopia have not accessed pain relief methods. The utilization of pharmacologic obstetric analgesia methods was found to be zero [6, 7]. The common factors that affect the utilization of pharmacological labor pain management methods were inadequate knowledge, negative attitude, unavailability of analgesic drugs in the health facilities, individual factor, and institutional factor [6-8].

Mother with intense labor pain affected by emotional distress, depression, and anxiety. Fear and anxiety are significant influences on pain experiences which is one reason why mothers are accompanied by another person during childbirth [9, 10]. Psychosocial factors have been implicated in the pain experienced during childbirth, which can have both short and long-term consequences on the mother's health and her relationship with her infant [3].

Ethiopia develops and implements Ethiopian Emergency Obstetric and Newborn Care (EmONC) assessment standards to enhance appropriate management of labor pain and management of complications during labor and delivery [11]. And also, the Federal Ministry of Health formulates different policy and management protocols regarding managing labor and delivery.

Despite those efforts, the provision of pain relief in labor is neglected and has remained an ongoing challenge [12, 13]. A lack of awareness, unfavorable attitude, and availability of pharmacological methods are considered to be the main reasons why women in Ethiopia do not receive adequate pain relief [6, 12, 14]. Even though healthcare providers in Ethiopia understand that women suffer significant pain during labor, currently they do not provide effective pain relief [15].



Though it is known that effective labor pain management results in greater birth satisfaction for the mother. In many LMIC including Ethiopia, OCPs' awareness of utilization of pharmacological labor pain management methods is low and even zero in some parts of the country. Moreover, to the best knowledge of the investigators concerned, evidence is scarce regarding the utilization of pharmacological labor pain management methods in the study area. So, this study was carried out to determine the utilization of pharmacological labor pain management methods among obstetric caregivers in Harari regional state health facilities, Ethiopia, 2021.

## **METHOD AND MATERIAL**

### **Study Design, Study Setting, and Period**

A facility-based cross-sectional study was conducted from May 20 to June 10, 2021, in Harari regional state health facilities, Ethiopia. The region is found 526 km towards east of Addis Ababa, the capital city of Ethiopia. The region is circled by Oromia regional state. The region is divided into nine woredas with three of them are rural and six are urban. The urban districts are subdivided into 19 kebeles (the lowest administrative division in the country), and the rural districts are subdivided into 17 peasant associations (which is equivalent to kebeles in urban cases). The total population (2021 projection based on the 2007 Census, CSA) of the region is 270,000 from which 136,000 are males and 134,000 are females [16]. There are two public hospitals, two private hospitals and 2 governmental hospitals, eight public health centers, and 20 health posts in the region.

### **Study Population**

All obstetric care providers in Harari regional state health facilities during the study period were the study population. Since there were small numbers of source population in the study area, all obstetric caregivers available during the study period were included. Obstetrics care providers not available during data collection due to illness, annual leave, and left for training were excluded from the study.

### **Sample Size Determination and Sampling Procedure**

The study was conducted at those health facilities having maternal services. These include two public hospitals (HFSUH, and Jugol hospital), one private hospital (Harar General hospital), two private clinics, and nine health centers of the Harari regional state. Since the number of the source

population (OCPs) in all health facilities is too small, all obstetrics caregivers were included in the study. Totally 464 Obstetrics Care Providers (OCPs) were included in the study.

### **Data Collection Methods and Data Quality Assurance**

A structured pre-tested self-administered questionnaire was prepared by adapting from different studies considering the local situation of the study area [7, 14, 17]. The questionnaire had four essential components including providers' socio-demographic characteristics, knowledge, attitude, and utilization-related questions.

The data was collected by research-qualified five professional Nurses and supervised by two MSC holding Midwives. The quality of the data was guaranteed by pretesting using the 5% of the respondents who are not eligible for the study before actual data collection. The completeness of the data was checked daily. Double data entry was done by two data clerks and consistency of entered data is cross-checked by comparing two separately entered data on Epi-Data.

### **Measurement**

Good utilization of pharmacological labor pain management methods: Obstetrics Care Providers (OCPs) who were practicing greater than or equal to the mean value of listed labor pain management methods (0.65(SD= 0.769)) [7]. While Obstetrics Care Providers (OCPs) who answered less than the mean values of practice-related labor pain relief method questions labeled as poor utilization.

Obstetrics Care Providers (OCPs): Midwifery, Nurses, Health Officers, and residents.

Adequate knowledge: Obstetrics Care Providers (OCPs) who answered greater than or equal to the mean values of knowledge-related labor pain relief method questions. Those Obstetrics Care Providers (OCPs) who answered  $\geq 6.65$  from total knowledge-related questions were considered as Obstetrics Care Providers (OCPs) who had adequate knowledge about labor pain relief methods [7].

Inadequate knowledge: Obstetrics Care Providers (OCPs) who answered less than the mean values of knowledge-related labor pain relief method questions. Those Obstetrics Care Providers (OCPs) who answered  $< 6.649$  from total knowledge-related questions were considered as OCGs who had inadequate knowledge about labor pain relief methods [7]

Favorable attitude: Obstetrics Care Providers (OCPs) who answered greater than or equal to the mean value of attitude-related labor pain relief method questions. Those Obstetrics Care Providers (OCPs) who answered  $\geq 4.1$  from total attitude-related questions were considered as OCPs who had a favorable attitude about labor pain relief methods [6]

Unfavorable attitude: OCPs who answered less than the mean value of attitude-related labor pain relief method questions. Those OCPs who answered  $< 4.1$  from total attitude-related questions were considered as OCPs who had an unfavorable attitude about labor pain relief methods [6]

### **Statistical Analysis**

The data were entered into Epi data statistical software version 3.1 and then exported to SPSS window version 22 for analysis. Descriptive statistical analysis such as simple frequency was used to describe the characteristics of study participants such as socio-demographic, level of knowledge, and attitude of obstetrics care providers. Then the information was presented using frequencies, tables, and figures.

Bivariate analysis and multivariate analysis were done to see the association between each independent variable and outcome variables by using binary logistic regression. The assumptions of binary logistic regression were checked. The goodness of fit was checked by Hosmer-Lemeshow statistic and omnibus tests. All variables with  $p < 0.25$  in the Bivariate analysis were included in the final model of multivariate analysis to control all possible confounders. The multi co-linearity test was carried out to see the correlation between independent variables by using the standard error and collinearity statistics (variance inflation factors  $> 10$  and standard error  $> 2$  was considered as suggestive of the existence of multi co-linearity). The direction and strength statistical association was measured by odds ratio with 95% CI. Adjusted odds ratio along with 95% CI was estimated to identify the association between independent variables and utilization of pharmacologic labor pain managements by using multivariate analysis in binary logistic regression. In this study,  $P\text{-value} < 0.05$  was considered to declare a result statistically significant

### **Ethical Considerations**

Ethical clearance to conduct this study was obtained from Haramaya University, College of Health and Medical Sciences, Institutional Health Research Ethics Review Committee (HU-IHRERC). Official letters to make a study was also obtained from health facilities as needed. The health

facilities were informed about the purpose of the study and the risk and benefits of the study. Informed, voluntary, written, and signed consent was obtained from the health facilities administrators and study participants before the data collection.

## **RESULT**

### **Socio-demographic Characteristics**

A total of 464 obstetric health care providers were involved and yielding a response rate of 100%. Among them, 200 (43.1%) were males and 264 (56.9%) were females and male to female ratio is 0.76 to 1. More than half 233(53%) of the study participants were Midwives (Table 1).

### **Knowledge of Obstetric Care Providers**

Almost all 444 (95.7%) health care providers knew about labor pain. Among them, 328 (73.9%) of obstetric care providers knew both pharmacological and non-pharmacological pain relief methods while 60 (13.5%) of them knew only pharmacological pain relief methods.

Among pharmacological pain relief methods, the majority 240 (61.9%) of obstetric care providers knew about non-steroidal anti-inflammatory drugs (NSAIDs), while 156 (40.2%) of them knew about systemic opioids and 140 (36.1%) of obstetric care providers know about epidural analgesia pain relief methods (Table 2).

Among OCPs who knew about labor pain, 356 (80.2%) of them knew about the side effect of pharmacological pain relief methods while 88 (19.8%) of them don't know about the side effect of pharmacological pain relief methods. Two hundred thirty-six (66.3%) of obstetric care providers respond that pharmacological pain relief methods causing the labor progress to be delayed while 208 (58.4%) of OCPs respond that pharmacologic pain relief methods cause fetal distress (Figure 1).

More than two-third 332 (71.6%) of obstetric care providers were asked to give pain relief methods. Most 252 (54.3%) obstetric care providers knew about the WHO pain ladder.

The mean score of knowledge questions is 6.65(SD= 2.95) with minimum and maximum scores of 0 and 15 respectively. More than 230 (49.6%) of the respondent have adequate knowledge.

The mean score of attitude questions is 4.01(SD= 1.4) with minimum and maximum scores of 0 and 6 respectively. Three hundred four (65.5%) of the respondent have an unfavorable attitude.

### **Utilization of Pharmacological Pain Relief Methods**

From a total of 464 obstetric care providers, 364 (78.4%) of obstetric care providers were asked to provide pain relief for laboring mothers. More than three-fourth 344 (74.1%) of OCPs would provide labor pain relief methods if they had resources. The main barriers for patients receiving pain relief include high patient flow 172 (54.4%), limited numbers of staff 164 (51.9%), lack of knowledge and skill 104 (32.9%), no drug available to provide 88 (27.8%), and no equipment available 48 (15.2%).

Among those OCPs (364), 32 (8.8%) of obstetric care providers provide only pharmacological pain relief methods while 204 (44%) of OCPs provide both pharmacological and non-pharmacological pain relief methods. From pharmacologic pain relief methods, the most common pharmacologic pain relief methods were pethidine 132 (55.9%), diclofenac 76 (32.2%), and paracetamol 72 (30.5%) (Figure 2).

The mean score of utilization questions is 0.65(SD= 0.769) with minimum and maximum scores of 0 and 5 respectively. The overall utilization of pharmacological labor pain relief methods was 50.9% (95% CI 45.8,55.8).

### **Institutional Related Factors**

Of the total of 464 study participants, 106 (91.4%) of the respondent replied that pharmacological drugs were available in their health facility whereas 10 (8.6%) of the participants responded that there were no drugs available in their health facility. It was reported that the most commonly available drugs were pethidine 292 (68.9%), diclofenac 324 (76.4%), paracetamol 300 (70.8%), and hyoscine 40 (9.4%). Nearly two-thirds 296 (63.8%) of OCPs got training about labor pain management methods from their respected facilities.

### **Obstetric Care Providers Preference**

Most 276 (59.5%) of obstetric care providers prefer both pharmacological and non-pharmacological pain relief methods to manage labor pain while 136 (29.3%) of obstetric care

providers prefer non-pharmacological pain relief methods to manage labor pain and 52 (11.2%) of OCPs prefer pharmacological pain relief methods to manage labor pain.

Three hundred sixty-eight (79.3%) of OCPs expect the intensity of labor pain was severe pain while 60 (12.9%) of OCPs expect moderate pain and 36 (7.8%) of them expect mild pain.

### **Factors Associated with Utilization of Pharmacological labor pain management methods**

Variables like age, sex, profession, year of experience, caregiver preferences, level of education, Pain expectation of caregiver, availability of equipment & analgesics drugs, level of knowledge, and attitude of the respondents were associated with utilization of pharmacological labor pain management methods in bivariate logistic regression analysis. In multivariate logistic regression sex, training, and attitude were factors associated with utilization of obstetric analgesia.

Multivariate analysis indicated that female OCPs were 2.07 times [AOR=2.07, 95% CI (1.31,3.26)] less likely to utilize pharmacological labor pain relief methods than male respondents. The odds of poor utilization of pharmacological labor pain relief methods among OCPs who had not trained were 1.60 times that of OCPs having training [AOR=1.60, 95%CI (1.02,2.53)].

Obstetric care providers with unfavorable attitudes were 9.1 times less likely to utilize pharmacological labor pain relief methods as compared to OCPs with favorable attitudes [AOR=9.1, 95%CI: (5.34,15.35)] (Table 3).

## **DISCUSSION**

This study assessed the utilization of pharmacological labor pain management methods among OCPs in Harari regional state health facilities. Additionally, factors that affect the utilization of pharmacological labor pain management methods were isolated.

The overall utilization of pharmacological labor pain relief methods was 50.9% [(95% CI (45.8,55.8)]. Which was higher than studies done in Canada 16.7% [9], Nigeria 38.9% [5], Leku primary hospital, southern Ethiopia 25.7% [18], Amhara regional state zero [7], southern Ethiopia 37.9% [6]. This variation might be due to study period differences and sample size differences. Meanwhile, this study was lower than a study conducted in Ethiopia which was 66% [15]. This

discrepancy might be due to sampling size difference and the health facility setups because some of the setups could be well-equipped, the presence of skilled man powers (who took training may provide obstetric analgesia to laboring mothers).

In this study, the main barriers for laboring women receiving pain relief included high patient flow, limited numbers of staff, lack of knowledge and skill, no drug available to provide, and no equipment available. This is in line with studies done in Nigeria [19, 20], Egypt [13], and the Amhara region [7]. These findings indicated that a weak health care system, weak facility organizations, and limited distribution of resources obscure access to maternal health services. The health bureau of the region should work in addressing those problems by arranging in-service training to equip OCPs with knowledge and attitude and fulfilling necessary drugs and equipment.

In this study, training was found to be associated with the utilization of pharmacologic labor pain relief methods. This finding is in harmony with studies conducted in Amhara regional state [7], southern Ethiopia [6], and the Tigray region [21]. This may be due to failure to train obstetric care providers result in the inadequacy of the skill which in turn yields the inability to utilize pharmacologic labor pain relief methods

Our study pointed out that level of attitudes of care providers was associated with the utilization of pharmacologic labor pain relief methods This is consistent with studies done in Amhara regional state [7, 14], southern Ethiopia [6], and the Tigray region [21]. This may be due to if obstetric care providers had a favorable attitude, they can feel the need for pain relief methods, and then they would provide pain relief methods.

The limitation of this study was it might not indicate a cause-effect relationship because the study design was cross-sectional. Since the study is institution-based and the results might lack generalization to the entire population in the catchment area. The study didn't incorporate women in labor.

## **Conclusion**

The study revealed that the overall prevalence of utilization of pharmacological labor pain management methods among obstetric care providers was low. In general, this study identified that variables like sex, training, and attitude of caregivers were factors associated with the utilization of pharmacological labor pain management methods. All stakeholders need to work

together to exploit OCPs' skills and attitudes. The health facilities should give the training to enhance the skill of obstetric care providers and equip themselves with sufficient analgesic drugs. Obstetric caregivers need to update and make themselves familiar with obstetric analgesia.

### **Abbreviations**

AOR: Adjusted Odds Ratio; EmONC: Emergency Obstetric and Newborn Care LMIC; Low Mid Income Country; OCG: Obstetrics Care Givers; OCP: Obstetrics Care Providers; WHO: World Health Organization.

### ***Declarations***

#### **Ethics Approval and Consent to Participate**

Ethical clearance was taken from the Institutional Health Research Ethics Review Committee (IHRERC) of the College of Health and Medical Sciences of Haramaya University. In addition, permission to proceed was obtained from all public hospitals. And also, informed voluntary written and signed consent was obtained from each participant.

#### **Consent for Publication**

Not applicable.

#### **Availability of Data and Materials.**

All relevant data are included in this study. However, additional data will be available from the corresponding author on reasonable request.

#### **Competing Interests.**

The authors declare that there is no competing interest in this work.

#### **Funding**

The funder (Haramaya University) has no role in the design of the study, collection, analysis, and interpretation of data, and in writing the manuscript.

#### **Author Contributions**

Addis Eyeberu conceived the idea and had major roles in the data review, drafting, and writing the result. All co-authors contributed to data analysis, drafting, and manuscript preparation. Finally,



all authors read and approved the final version of the manuscript to be published and agreed on all aspects of this work.

## Acknowledgments

We would like to extend our deepest thanks to Haramaya University, College of Health and Medical Science for helping us in performing this paper and we also would like to extend our acknowledgment to our friends for providing their constructive support.

## REFERENCES

1. Labor, S. and S. Maguire, *The Pain of Labour*. Reviews in pain, 2008. **2**(2): p. 15-19.
2. Thomson, G., et al., *Women's experiences of pharmacological and non-pharmacological pain relief methods for labor and childbirth: a qualitative systematic review*. Reproductive Health, 2019. **16**(1): p. 71.
3. N, L., O. Yvonne, and S. Ghosh, *Physical and Psychological Aspects of Pain in Obstetrics*. 2012.
4. Czech, I., et al., *Pharmacological and Non-Pharmacological Methods of Labour Pain Relief-Establishment of Effectiveness and Comparison*. International journal of environmental research and public health, 2018. **15**(12): p. 2792.
5. Imarengiaye, C. and A. Ande, *Demand and utilisation of labour analgesia service by Nigerian women*. Journal of obstetrics and gynecology, 2006. **26**(2): p. 130-132.
6. Geltore, T.E., A. Taye, and A.G. Kelbore, *Utilization of obstetric analgesia in labor pain management and associated factors among obstetric care givers in public health facilities of Kembata Tembaro Zone, Southern Ethiopia*. Journal of pain research, 2018. **11**: p. 3089-3097.
7. Bitew, A., et al., *Utilization of obstetric analgesia in labor pain management and associated factors among obstetric caregivers in Amhara Regional State Referral Hospitals, Northwest Ethiopia. A hospital-based cross-sectional study*. Journal of Biomedical Sciences, 2016. **5**(2).
8. Adams, J., et al., *Use of pharmacological and non-pharmacological labor pain management techniques and their relationship to maternal and infant birth outcomes: an*

- examination of a nationally representative sample of 1835 pregnant women. Midwifery, 2015. 31(4): p. 458-463.*
9. Agaram, R., et al., *Inadequate pain relief with labor epidurals: a multivariate analysis of associated factors. International journal of obstetric anesthesia, 2009. 18(1): p. 10-14.*
  10. Barrett, S.J. and M.A. Stark, *Factors Associated With Labor Support Behaviors of Nurses. The Journal of perinatal Education, 2008. 19(1): p. 12-18.*
  11. FMOH, *ETHIOPIAN Emergency Obstetric and Newborn Care (EmONC) Assessment 2016. 2016.*
  12. Geltore, T.E., A.G. Kelbore, and A.T. Angelo, *Perceptions of obstetric analgesia: a qualitative study among midwives attending normal vaginal deliveries in Durame Hospital, Southern Ethiopia. J Pain Res, 2019. 12: p. 2187-2192.*
  13. Mousa, O., et al., *Perceptions and Practice of Labor Pain-Relief Methods among Health Professionals Conducting Delivery in Minia Maternity Units in Egypt. Obstet Gynecol Int, 2018. 2018: p. 3060953.*
  14. Bishaw, K.A., E.G. Sendo, and W.S. Abebe, *Knowledge, and use of labor pain relief methods and associated factors among obstetric care givers at public health centers of East Gojjam zone, Amhara region, Ethiopia: a facility-based cross-sectional study. BMC Pregnancy Childbirth, 2020. 20(1): p. 180.*
  15. McCauley, M., C. Stewart, and B. Kebede, *A survey of healthcare providers' knowledge and attitudes regarding pain relief in labor for women in Ethiopia. BMC Pregnancy Childbirth, 2017. 17(1): p. 56.*
  16. CSA, *CSA Website Analysis and Recommendation.pdf. 2009.*
  17. Getu, A.A., et al., *Non-pharmacological labor pain management and associated factor among skilled birth attendants in Amhara Regional State health institutions, Northwest Ethiopia. Reprod Health, 2020. 17(1): p. 183.*
  18. Siyoum, M. and S. Mekonnen, *Labor pain control and associated factors among women who gave birth at Leku primary hospital, southern Ethiopia. BMC Res Notes, 2019. 12(1): p. 619.*
  19. Ogboli-Nwasor, E., et al., *Pain relief in labor: a survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. Journal of Pain Research, 2011.*

20. Ohaeri, B., G. Owolabi, and J. Ingwu, *Skilled health attendants' knowledge and practice of pain management during labor in health care facilities in Ibadan, Nigeria*. European Journal of Midwifery, 2019. **3**(February).
21. Sahile, E., et al., *Practice of Labour Pain Management Methods and Associated Factors among Skilled Attendants Working at General Hospitals in Tigray Region, North Ethiopia: Hospital-Based Cross-Sectional Study Design*. Health Science Journal, 2017. **11**(4).

## List of Tables

Table 1 Socio-demographic characteristics of obstetric care providers at Harari regional state public health facilities, 2021 (n=464)

Variables	Categories	Frequency	Percentage
Sex	Male	200	43.1
	Female	264	56.9
Age	15-24	60	12.9
	25-54	404	87.1
Religion	Orthodox	220	47.4
	Muslim	176	37.9
	Protestant	52	11.2
	Others	16	3.4
Profession	Midwife	233	53
	Nurse	198	42.67
	Obstetrician	20	4.3
	Health officers	13	2.8
Level of education	Diploma	142	30.6
	BSC	284	61.2
	MSC	38	8.2
Clinical experience	<2years	88	19
	≥2years	376	81

\*others=Catholic and Adventist, BSC- Bachelor of Science, MSC- Masters of Sciences

Table 2 Knowledge about types of pharmacological pain relief methods among obstetric caregivers at Harari regional state health facilities (n=388) from May 20 to June 10, 2021

Variable	Categories	Frequency	Percentage
Systemic opioids	Yes	156	40.2
	No	232	59.8
Non-Steroidal Anti-inflammatory Drugs (NSAIDs)	Yes	240	61.9
	No	148	38.1
Epidural analgesia	Yes	140	36.1
	No	248	63.9
Inhalational analgesia	Yes	16	4.1
	No	372	95.9

Table 3 Factors associated with utilization of pharmacological pain relief methods among obstetric care providers at Harari regional state health facilities from May 20 to June 10, 2021.

Variables		Utilization of Pharmacological labor pain management		COR 95%CI	AOR 95%CI
		Poor	Good		
Sex	Male	112	88	1	1
	Female	116	148	1.62(1.12,2.35) *	2.07(1.31,3.26) *
Age	25-54	208	196	1	1
	15-24	20	40	2.12(1.19,3.76) *	0.51(0.26,1.01)
Level of education	BSc and above	168	148	1.67(1.12,2.47) *	1.22(0.76,1.95)
	Diploma	60	88	1	1
Training	No	108	60	2.64(1.78,3.91) *	1.60(1.02,2.53) *
	Yes	120	176	1	
Pain expectation	Moderate	29	31	0.412(0.17,0.984)	0.97(0.44,2.15)
	Sever	192	176	0.48(0.23,1.04)	1.28(0.69,2.38)
	Mild	25	11	1	1
Attitude	Unfavorable	200	104	9.07(5.66,14.53) *	9.1(5.34,15.35) *
	Favorable	28	132	1	1

Knowledge	Adequate	114	116	1	1
	Inadequate	114	120	1.03(0.72,1.49)	1.03(0.66,1.61)

\*P≤ 0.001, OR=Odd ratio, AOR=Adjusted odd ratio, CI=Confidence interval




Figure 1 knowledge of OCP about the side effect of pharmacological pain relief methods among obstetric caregivers at Harari regional state health facilities from May 20 to June 10, 2021.




Figure 2 Most common pharmacological pain relief methods provided at Harari regional state health facilities from May 20 to June 10, 2021.



# Utilization of pharmacological labour analgesia: a survey of obstetric care providers in eastern Ethiopia

Addis Eyeberu<sup>a,\*</sup>, Tamirat Getachew<sup>a</sup>, Adera Debella<sup>a</sup>, Bikila Balis<sup>a</sup>, Bajirond Eshetu<sup>a</sup>, Sinetibet Mesfin<sup>a</sup>, Habtamu Bekele<sup>a</sup>, Dawit Tamiru<sup>a</sup>, Getahun Tiruye<sup>a</sup>, Meron Degefa<sup>a</sup>, Addisu Alemu<sup>b,\*</sup>, Merga Dheressa<sup>a</sup> and Yadeta Dessie<sup>b</sup>

<sup>a</sup>School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia; <sup>b</sup>School of Public Health, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

\*Corresponding authors: Tel: +251910005436, +251911611147; E-mail: [addiseyeberu@gmail.com](mailto:addiseyeberu@gmail.com), [alemuadisu789@gmail.com](mailto:alemuadisu789@gmail.com)

Received 3 December 2021; revised 3 August 2022; editorial decision 17 August 2022; accepted 19 August 2022

**Background:** Despite a crucial role played by obstetric care providers (OCPs) in providing pain relief methods during labour, the evidence suggests that there is a large number of women in Ethiopia who did not have access to pain relief methods. Moreover, to the best of the investigators' knowledge, evidence is scarce regarding the utilization of pharmacological labour pain management methods in the study area. Thus this study was carried out to determine the practice of pharmacological labour pain management methods among OCPs in Harari regional state health facilities in Ethiopia.

**Methods:** A facility-based cross-sectional study was conducted from 20 May to 10 June 2021 in Harari regional state health facilities. All (n=464) OCPs in Harari regional state health facilities were included in the study. Data were collected using a structured questionnaire adapted from previous studies. The data were entered into Epi-data version 3.1 statistical software. Statistical analysis was carried out using SPSS for Windows version 22. Bivariate and multivariate logistic regression analyses were employed to determine the association between independent variables and the outcome variable. A p-value <0.05 was used to establish statistical significance.

**Results:** In this study, 464 OCPs participated. The overall utilization of pharmacological labour pain relief methods was 50.9% (95% confidence interval [CI] 45.8 to 55.8). The main barriers to labouring mothers receiving pain relief include high patient flow (54.4%), the limited number of staff (51.9%) and a lack of knowledge and skill (32.9%). Being female (adjusted odds ratio [AOR] 2.07 [95% CI 1.31 to 3.26]), not having training (AOR 1.60 [95% CI 1.02 to 2.53]) and having an unfavourable attitude (AOR 9.1 [95% CI 5.34 to 15.35]) were the predictors of utilization of pharmacological labour pain management methods.

**Conclusions:** The study revealed that half of the OCPs utilized the pharmacological labour pain management method. In general, this study identified that variables like sex, training and the attitude of OCPs were factors associated with the utilization of pharmacological labour pain management methods. The health facilities should provide training to enhance the skill of OCPs and equip themselves with sufficient analgesic drugs.

**Keywords:** analgesia, Harar, labour pain, pharmacological labour pain management, utilization.

## Introduction

Pain during labour and childbirth is a unique and the most severe pain that a woman ever has to face. The extent to which a woman feels in control of pain during labour is an indicator of maternal emotive well-being in childbirth.<sup>1</sup> Pharmacological labour pain management methods include oral tablets (paracetamol, codeine or tramadol), inhalations and injectable analgesics.<sup>2</sup>

Pain related to childbirth is present during pregnancy, labour and post-partum periods.<sup>3</sup> Evidence showed that the care provided to the parturient during labour and delivery is one of the indicators of the satisfaction of childbirth.<sup>4</sup> Pain during childbirth has an impact on maternal health, particularly its physiological and psychological influences.<sup>5,6</sup> Furthermore, pain has a profound impact on maternal health, including anxiety, depression,

and alteration of respiratory patterns.<sup>6,7</sup> Evidence shows that 51.9% of women had stress.<sup>8</sup> Psychosocial factors have been implicated in the pain experienced during childbirth, which can have both short- and long-term consequences on the mother's health and her relationship with her infant.<sup>3</sup>

Evidence shows that labour pain analgesia has an effect on the mother and newborn, and labour outcomes include increased duration of labour, rates of instrumental deliveries, hypotension, the likelihood of maternal fever, neonatal sepsis and increased duration of breastfeeding.<sup>9–12</sup> Even if those effects exist, labour pain is an individual phenomenon, so maternal request is considered a sufficient justification for pain relief.<sup>13</sup> Furthermore, evidence shows that pain relief should be provided to labouring women.<sup>14</sup>

Even if obstetric care providers (OCPs) have an imperative role in providing pain relief options during labour, the majority of women in low-income countries, including Ethiopia, have not accessed pain relief methods. Evidence shows that 85.1% of the patients want their labour pain to be relieved, but only 38.9% received analgesia during labour.<sup>15</sup> Studies done in Africa also report that the overall utilization of pharmacological labour pain methods varies and was 70.1% in Egypt<sup>16</sup> and 48.4% and 49% in Nigeria.<sup>14,17</sup> The utilization of pharmacological obstetric analgesia methods in Ethiopia also varies from place to place. A study done in Amhara and Tigray regions showed that the utilization of pharmacological labour analgesia was zero.<sup>13,18</sup> A similar study done in Addis Ababa and southern Ethiopia showed that the overall utilization of pharmacological labour analgesia was 36.6% and 45.2%, respectively.<sup>13,19</sup> The common factors associated with the non-utilization of pharmacological labour pain management methods were inadequate knowledge, negative attitude, unavailability of analgesic drugs in the health facilities, individual factors and institutional factors.<sup>12,13,18</sup>

Ethiopia has developed and implemented standards of care, including the provision of physical and psychological support and the use of pharmacological and non-pharmacological comfort measures during labour and birth.<sup>20</sup> The Ethiopian Food, Medicine and Health Care Administration and Control Authority (FMHACA) also recommended the use of pharmacological labour pain management methods.<sup>21</sup>

Despite these efforts, the provision of pain relief in labour is neglected and has remained an ongoing challenge.<sup>16,22</sup> A lack of awareness, unfavourable attitude and a lack of availability of pharmacological methods are considered to be the main reasons why women in Ethiopia do not receive adequate pain relief.<sup>13,22,23</sup> Even though healthcare providers in Ethiopia understand that women suffer significant pain during labour, currently they do not provide effective pain relief.<sup>24</sup>

Although it is known that effective labour pain management results in greater birth satisfaction for the mother, in many low- and middle-income countries (LMICs), including Ethiopia, OCPs' utilization of pharmacological labour pain management methods is low. Moreover, to the best of our knowledge, evidence is scarce regarding the utilization of pharmacological labour pain management methods in the study area. Thus this study was carried out to determine the utilization of pharmacological labour pain management methods among obstetric caregivers in Harari regional state health facilities.

## Methods

### Study design, study setting, and period

An institution-based cross-sectional study was conducted in Harari regional state health facilities from 20 May to 10 June 2021. The region is located 526 km east of Addis Ababa, the capital city of Ethiopia. The region is circled by Oromia regional state. The region is divided into nine *woredas*, with three of them being rural and six being urban. The urban districts are subdivided into 19 *kebeles* (the lowest administrative division in the country) and the rural districts are subdivided into 17 peasant associations (which is equivalent to *kebeles*). The total population (2021 projection based on the 2007 census) of the region is 270 000, of which 136 000 are males and 134 000 are females.<sup>25</sup> There are two public hospitals, two private hospitals and two governmental hospitals, eight public health centres and 20 health posts in the region. The number of healthcare providers in the Hiwot Fana Specialized University Hospital (HFSUH) and Jugol Hospital is 410 and 266, respectively.

### Study population

All OCPs in Harari regional state health facilities during the study period were the study population. These include two public hospitals (HFSUH and Jugol Hospital), one private hospital (Harar General Hospital), two private clinics and nine health centres in the Harari region. All of the 464 obstetric caregivers available during the study period were included. OCPs not available during data collection due to illness, annual leave or left for training were excluded from the study.

### Data collection methods and data quality assurance

A structured pretested self-administered questionnaire was prepared by adapting from different studies considering the local situation of the study area.<sup>18,23,26</sup> The questionnaire had four essential components: providers' sociodemographic characteristics and knowledge, attitude and utilization-related questions.

The data were collected by five research-qualified professional nurses and supervised by two MSc-holding midwives. The quality of the data was guaranteed by pretesting using 5% of the respondents who were not eligible for the study. The completeness of the data was checked daily. Double data entry was done by two data clerks and the consistency of entered data was cross-checked by comparing two separately entered data using Epi-Data.

### Variables

The outcome variable was the utilization of pharmacological labour pain management. Independent variables included sociodemographic factors such as age, sex, profession and year of experience; institutional factors included health institution policy, availability of equipment and analgesic drugs; and knowledge, attitude and individual factors included respondent's level of knowledge, qualifications, attitude and pain expectation.

## Measurement

OCPs who were practicing greater than or equal to the mean value of listed labour pain management methods (0.65 [standard deviation {SD} 0.769]) were considered to have good utilization of pharmacological labour pain management methods,<sup>18</sup> while OCPs who answered less than the mean values of practice-related labour pain relief method questions were considered to have poor utilization.

Knowledge was assessed by 10 items and the questions were answered with yes, no or I don't know responses. During analysis, the correct answer was coded as 1 and incorrect or I don't know answers were coded as 0. The sum score (range 0–10) was then computed and categorized. OCPs who scored  $\geq 6.65$  on the knowledge-related questions were considered to have adequate knowledge about labour pain relief methods, while OCPs who scored  $< 6.649$  were considered to have inadequate knowledge about labour pain relief methods.<sup>18</sup>

Attitude was assessed by a total of six items and the questions were answered with yes, no or I don't know responses. During analysis, the correct answer was coded as 1 and incorrect or I don't know answers were coded as 0. The sum score (range 0–10) was then computed and categorized. OCPs who scored  $\geq 4.1$  on the attitude-related questions were considered to have a favourable attitude towards labour pain relief methods, while OCPs who scored  $< 4.1$  were considered to have an unfavourable attitude towards labour pain relief methods.<sup>13</sup>

## Statistical analysis

The data were entered into EpiData statistical software version 3.1 (<http://epidata.dk/>) and then exported to SPSS for Windows version 22 (IBM, Armonk, NY, USA) for analysis. Descriptive statistical analysis such as simple frequency was used to describe the characteristics of study participants, including sociodemographics, level of knowledge and attitudes of OCPs. Then the information was presented using frequencies, tables and figures.

Bivariate analysis and multivariate binary logistic regression analysis were performed to determine the association between each independent variable and the outcome variable. The assumptions of binary logistic regression were checked. The goodness of fit was checked by Hosmer–Lemeshow statistic and omnibus tests. All variables with a p-value  $< 0.25$  in the bivariate analysis were included in the final model of multivariate analysis to control all possible confounders. A multicollinearity test was carried out to determine the correlation between independent variables by using the standard error and collinearity statistics (variance inflation factors  $> 10$  and standard error  $> 2$  were considered suggestive of the existence of multicollinearity). The direction and strength of the statistical association were measured by odds ratios (ORs) with 95% confidence intervals (CIs). Adjusted ORs along with 95% CIs were estimated to identify the association between independent variables and the utilization of pharmacological labour pain management by using multivariate analysis in binary logistic regression. In this study, p-values  $< 0.05$  were considered statistically significant.

**Table 1.** Sociodemographic characteristics of OCPs at Harari regional state public health facilities, 2021 (N=464)

Variables	Categories	n	%
Sex	Male	200	43.1
	Female	264	56.9
Age (years)	15–24	60	12.9
	25–54	404	87.1
Religion	Christian	220	47.4
	Muslim	176	37.9
	Protestant	52	11.2
	Others <sup>a</sup>	16	3.5
Profession	Midwife	233	50.2
	Nurse	198	42.7
	Obstetrician	20	4.3
	Health officer	13	2.8
Level of education	Diploma	142	30.6
	BSc	284	61.2
	MSc	38	8.2
Clinical experience (years)	$< 2$	88	19
	$\geq 2$	376	81

<sup>a</sup>Catholic and Adventist.

BSc: Bachelor of Science; MSc: Master's of Science.

## Ethical considerations

Ethical approval to conduct this study was obtained from the Haramaya University, College of Health and Medical Sciences, Institutional Health Research Ethics Review Committee (IHRERC/066/2021). Official letters to conduct a study were also obtained from health facilities as needed. The health facilities were informed about the purpose of the study and the risks and benefits of the study. Informed, voluntary, written and signed consent was obtained from the health facilities' administrators and study participants before data collection.

## Results

### Sociodemographic characteristics

A total of 464 OCPs were involved, yielding a response rate of 100%. Among them, 200 (43.1%) were males and 264 (56.9%) were females and the male:female ratio was 0.76:1. More than half (233 [53%]) of the study participants were midwives (Table 1).

### Knowledge of OCPs

Almost all (444 [95.7%]) OCPs knew about labour pain management methods. Of these, 328 (73.9%) knew both pharmacological and non-pharmacological pain relief methods and 60 (13.5%) of them knew only pharmacological pain relief methods.

Among pharmacological pain relief methods, the majority (240 [61.9%]) of OCPs knew about non-steroidal anti-inflammatory drugs (NSAIDs), while 156 (40.2%) of them knew

**Table 2.** Knowledge about types of pharmacological pain relief methods among OCPs at Harari regional state health facilities (n=388) from 20 May to 10 June 2021

Variables	Categories	n	%
Systemic opioids	Yes	156	40.2
	No	232	59.8
Non-steroidal anti-inflammatory drugs	Yes	240	61.9
	No	148	38.1
Epidural analgesia	Yes	140	36.1
	No	248	63.9
Inhalational analgesia	Yes	16	4.1
	No	372	95.9

about systemic opioids and 140 (36.1%) knew about epidural analgesia pain relief methods (Table 2).

Among OCPs who knew about labour pain management, 356 (80.2%) knew about the side effects of pharmacological pain relief methods, while 88 (19.8%) of them did not know about the side effects. A total of 236 (66.3%) OCPs responded that pharmacological pain relief methods cause the labour progress to be delayed, while 208 (58.4%) responded that pharmacological pain relief methods cause foetal distress (Figure 1).

More than two-thirds (332 [71.6%]) of OCPs had been asked to provide pain relief and most (252 [54.3%]) OCPs knew about the World Health Organization pain ladder.

The mean score of knowledge questions was 6.65 (SD 2.95), with minimum and maximum scores of 0 and 15, respectively. A total of 230 (49.6%) of the respondent had adequate knowledge.

Regarding attitudes of OCPs, more than half (280 [60.3%]) of OCPs believe that analgesics are necessary to manage labour pain and more than two-thirds (84.5%) believe that they have the

responsibility and obligation to manage labour pain. The mean score of attitude questions was 4.01 (SD 1.4), with minimum and maximum scores of 0 and 6, respectively. A total of 304 (65.5%) of the respondent had an unfavourable attitude (Table 3).

### Utilization of pharmacological pain relief methods


Of 464 OCPs, 364 (78.4%) were asked to provide pain relief for labouring mothers. Almost three-fourths (344 [74.1%]) of OCPs would provide labour pain relief methods if they had the resources. The main barriers for patients receiving pain relief include high patient flow (172 [54.4%]), limited staff (164 [51.9%]), lack of knowledge and skill (104 [32.9%]), no drugs available (88 [27.8%]) and no equipment available (48 [15.2%]).

Among the 364 OCPs, 32 (8.8%) provide only pharmacological pain relief methods while 204 (44%) provide both pharmacological and non-pharmacological pain relief methods. The most common pharmacological pain relief methods were pethidine (132 [55.9%]), diclofenac (76 [32.2%]) and paracetamol (72 [30.5%]) (Figure 2).

The mean score of utilization questions was 0.65 (SD 0.769), with minimum and maximum scores of 0 and 5, respectively. The overall utilization of pharmacological labour pain relief methods was 50.9% (95% CI 45.8 to 55.8).

### Institutional-related factors

Of the 464 study participants, 106 (91.4%) replied that pharmacological drugs were available in their health facility, whereas 10 (8.6%) of the participants responded that there were no drugs available in their health facility. It was reported that the most commonly available drugs were pethidine (292 [68.9%]), diclofenac (324 [76.4%]), paracetamol (300 [70.8%]) and hyoscine (40 [9.4%]). Nearly two-thirds (296 [63.8%]) of OCPs got training about labour pain management methods from their respective facilities.


**Figure 1.** Knowledge of the side effects of pharmacological pain relief methods among OCPs at Harari regional state health facilities from 20 May to 10 June 2021.



**Table 3.** Attitudes of OCPs about pharmacological pain relief methods in Harari regional state public health facilities, 2021

Variables	Categories	n	%
Believe that labour pain management methods alleviate labour pain (n=464)	Yes	392	84.5
	No	72	15.5
Believe every mother during labour should be managed (n=464)	Yes	320	69
	No	144	31
Believe that labour pain is natural and the mother has to face it (n=464)	Yes	344	74.1
	No	120	25.9
Believe pharmacology methods are necessary for managing labour pain (n=464)	Yes	280	60.3
	No	184	39.7
Believe that you have a responsibility and obligation to manage labour pain (n=464)	Yes	392	84.5
	No	72	15.5

pharmacological pain relief methods



**Figure 2.** Most common pharmacological pain relief methods provided at Harari regional state health facilities from 20 May to 10 June 2021.

**OCP preference**

Most OCPs (276 [59.5%]) prefer both pharmacological and non-pharmacological pain relief methods to manage labour pain while 136 (29.3%) OCPs prefer non-pharmacological pain relief methods and 52 (11.2%) OCPs prefer pharmacological pain relief methods. A total of 368 (79.3%) OCPs expected the intensity of labour pain to be severe, while 60 (12.9%) expected moderate pain and 36 (7.8%) expected mild pain.

**Factors associated with the utilization of pharmacological labour pain management methods**

Variables such as age, sex, profession, year of experience, caregiver preferences, level of education, pain expectation of the

caregiver, availability of equipment and analgesic drugs, level of knowledge and attitude of the caregiver were associated with the utilization of pharmacological labour pain management methods in bivariate logistic regression analysis. In multivariate logistic regression sex, training and attitude were factors associated with the utilization of obstetric analgesia.

Multivariate analysis indicated that female OCPs were 2.07 times (adjusted OR [AOR] 2.07 [95% CI 1.31 to 3.26]) more likely to utilize pharmacological labour pain relief methods than male respondents. The odds of poor utilization of pharmacological labour pain relief methods among OCPs who had not been trained were 1.60 times those of OCPs who had training (AOR 1.60 [95% CI 1.02 to 2.53]). OCPs with unfavourable attitudes were 9.1 times less likely to utilize pharmacological labour pain relief methods as compared with OCPs with favourable attitudes (AOR 9.1 [95% CI 5.34 to 15.35]) (Table 4).

**Discussion**

This study assessed the utilization of pharmacological labour pain management methods among OCPs in Harari regional state health facilities. Additionally, factors that affect the utilization of pharmacological labour pain management methods were isolated.

The overall utilization of pharmacological labour pain relief methods was 50.9% (95% CI 45.8 to 55.8), which was higher than in studies in Canada (16.7%),<sup>27</sup> Nigeria (38.9%),<sup>15</sup> Addis Ababa, Ethiopia (36.6%),<sup>19</sup> Amhara regional state (0%)<sup>18</sup> and southern Ethiopia (37.9%).<sup>13</sup> This variation might be due to study period differences and sample size differences. Meanwhile, the utilization in this study was lower than in studies conducted in Egypt (70.1%) and another study in Ethiopia (66%).<sup>24</sup> This discrepancy might be due to sampling size differences and health facilities because some of the health facilities may be well equipped, with skilled manpower.

In this study, the main barriers for labouring women receiving pain relief included high patient flow, limited staff, a lack of knowledge and skill, no drugs available and no equipment available. This is in line with studies done in Nigeria,<sup>28,29</sup> Egypt<sup>16</sup> and the Amhara region.<sup>18</sup> These findings indicated that a weak

**Table 4.** Factors associated with utilization of pharmacological pain relief methods among OCPs at Harari regional state health facilities from 20 May to 10 June 2021

Variables		Utilization of pharmacological labour pain management, n		OR (95% CI)	AOR (95% CI)
		Poor	Good		
Sex	Male	112	88	1	1
	Female	116	148	1.62 (1.12 to 2.35)*	2.07 (1.31 to 3.26)*
Age (years)	25–54	208	196	1	1
	15–24	20	40	2.12 (1.19 to 3.76)*	0.51 (0.26 to 1.01)
Level of education	BSc and above	168	148	1.67 (1.12 to 2.47)*	1.22 (0.76 to 1.95)
	Diploma	60	88	1	1
Training	No	108	60	2.6(1.78 to 3.91)*	1.6 (1.02 to 2.53)*
	Yes	120	176	1	1
Pain expectation	Moderate	29	31	0.412 (0.17 to 0.984)	0.97 (0.44 to 2.15)
	Severe	192	176	0.48 (0.23 to 1.04)	1.28 (0.69 to 2.38)
	Mild	25	11	1	1
Attitude	Unfavourable	200	104	9.07 (5.66 to 14.53)*	9.1 (5.34 to 15.35)*
	Favourable	28	132	1	1
Knowledge	Adequate	114	116	1	1
	Inadequate	114	120	1.03 (0.72 to 1.49)	1.03 (0.66 to 1.61)

\* $p \leq 0.001$ .

healthcare system, weak facility organizations and limited distribution of resources limit access to maternal health services. Arranging in-service training to equip OCPs with knowledge and providing necessary drugs and equipment could be the solution.

In this study, training was found to be associated with the utilization of pharmacological labour pain relief methods. This finding is in harmony with studies conducted in the Amhara region,<sup>18</sup> southern Ethiopia<sup>13</sup> and the Tigray region.<sup>30</sup> This may be due to failure to train OCPs, resulting in inadequate skills.

Our study pointed out that the attitudes of care providers was associated with the utilization of pharmacological labour pain relief methods, as those with unfavourable attitudes were less likely to utilize pharmacological pain relief methods. This is consistent with studies done in the Amhara region,<sup>18,23</sup> southern Ethiopia<sup>13</sup> and the Tigray region.<sup>30</sup> Moreover, female OCPs were more likely to utilize pharmacological labour pain relief methods than male respondents. This may be because female OCPs may have giving birth in the past and experienced the pain and they may not want their clients to suffer the pain.

The limitation of this study was that the assessment of the utilization of pharmacological labour pain management methods only encompassed OCPs and did not include the service receivers (mothers facing labour pain during childbirth).

## Conclusions

The study revealed that the overall prevalence of utilization of pharmacological labour pain management methods among

OCPs was low. In general, this study identified that variables like sex, training and the attitude of caregivers were factors associated with the utilization of pharmacological labour pain management methods. All stakeholders need to work together to enhance the skills and attitudes of OCPs. The health facilities should provide training to enhance the skills of OCPs and equip themselves with sufficient analgesic drugs and OCPs need to familiarize themselves with obstetric analgesia.

**Authors' contributions:** AE conceived the idea and was involved in the data review and drafting and writing the manuscript. TG, AD, BB, BE, SM and HB were responsible for proposal writing, project administration, methodology, data analysis and drafting the manuscript. AE, DT, GT, MD, AA, MDA and YD contributed to data analysis, drafting and manuscript preparation. Finally, all authors read and approved the final version of the manuscript to be published and agreed on all aspects of this work.

**Acknowledgments:** We would like to extend our deepest thanks to Haramaya University, College of Health and Medical Science, data collectors, and participants, for helping us with this research and we would also like to extend thanks to our friends for providing their constructive support.

**Funding:** The funder (Haramaya University) had no role in the design of the study; collection, analysis and interpretation of data; or writing the manuscript.

**Competing interests:** None declared.

**Data availability:** All relevant data are included in this study. However, additional data is available from the corresponding author upon reasonable request.

## References

- Labor S, Maguire S. The pain of labour. *Rev Pain*. 2008;2(2):15–19.
- Thomson G, Feeley C, Moran VH, et al. Women's experiences of pharmacological and non-pharmacological pain relief methods for labor and childbirth: a qualitative systematic review. *Reprod Health*. 2019;16(1):71.
- Ebirim LN, Yvonne O, Ghosh S. Physical and psychological aspects of pain in obstetrics. In: Ghosh S, editor. *Pain in perspective*. Available from: <https://www.intechopen.com/chapters/40395> [accessed 29 August 2022].
- Czech I, Fuchs P, Fuchs A, et al. Pharmacological and non-pharmacological methods of labour pain relief-establishment of effectiveness and comparison. *Int J Environ Res Public Health*. 2018;15(12):2792.
- Olde E, van der Hart O, Kleber R, et al. Posttraumatic stress following childbirth: a review. *Clin Psychol Rev*. 2006;26(1):1–16.
- Lowe NK. The nature of labor pain. *Am J Obstet Gynecol*. 2002;186(5 Suppl):S16–24.
- Junge C, von Soest T, Weidner K, et al. Labor pain in women with and without severe fear of childbirth: a population-based, longitudinal study. *Birth*. 2018;45(4):469–77.
- Tan HS, Agarhesh T, Tan CW, et al. Perceived stress during labor and its association with depressive symptomatology, anxiety, and pain catastrophizing. *Sci Rep*. 2021;11(1):17005.
- Lieberman E, O'Donoghue C. Unintended effects of epidural analgesia during labor: a systematic review. *Am J Obstet Gynecol*. 2002;186(5 Suppl):S31–68.
- Leighton BL, Halpern SH. The effects of epidural analgesia on labor, maternal, and neonatal outcomes: a systematic review. *Am J Obstet Gynecol*. 2002;186(5 Suppl):S69–77.
- Leeman L, Fontaine P, King V, et al. The nature and management of labor pain: part II. Pharmacologic pain relief. *Am Fam Physician*. 2003;68(6):1115–20.
- Adams J, Frawley J, Steel A, et al. Use of pharmacological and non-pharmacological labor pain management techniques and their relationship to maternal and infant birth outcomes: an examination of a nationally representative sample of 1835 pregnant women. *Midwifery*. 2015;31(4):458–63.
- Geltore TE, Taye A, Kelbore AG. Utilization of obstetric analgesia in labor pain management and associated factors among obstetric caregivers in public health facilities of Kembata Tembaro Zone, Southern Ethiopia. *J Pain Res*. 2018;11:3089–97.
- Ogboli-Nwasor E, Adaji SE, Bature SB, et al. Pain relief in labor: a survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. *J Pain Res*. 2011;4:227.
- Imarengiaye C, Ande A. Demand and utilisation of labour analgesia service by Nigerian women. *J Obstet Gynecol*. 2006;26(2):130–2.
- Mousa O, Abdelhafez AA, Abdelraheim AR, et al. Perceptions and practice of labor pain-relief methods among health professionals conducting delivery in Minia maternity units in Egypt. *Obstet Gynecol Int*. 2018;2018:3060953.
- Lawani LO, Eze JN, Anozie OB, et al. Obstetric analgesia for vaginal birth in contemporary obstetrics: a survey of the practice of obstetricians in Nigeria. *BMC Pregnancy Childbirth*. 2014;14:140.
- Bitew A, Workie A, Seyum T, et al. Utilization of obstetric analgesia in labor pain management and associated factors among obstetric caregivers in Amhara Regional State Referral Hospitals, Northwest Ethiopia. A hospital-based cross-sectional study. *J Biomed Sci*. 2016;5(2):15.
- Gido R, Yadeta TA, Tura AK. Utilization of obstetric analgesia for labor pain management and associated factors among obstetric care providers in public hospitals of Addis Ababa, Ethiopia: a cross-sectional study. *Obstet Gynecol Int*. 2021;2021:9973001.
- Ministry of Health. Ethiopian Emergency Obstetric and Newborn Care (EmONC) Assessment 2016. Addis Ababa: Ethiopian Public Health Institute; 2016.
- Food, Medicine and Healthcare Administration and Control Authority of Ethiopia. Standard treatment guidelines for general hospitals. Addis Ababa: Ethiopian Federal Ministry of health; 2014.
- Geltore TE, Kelbore AG, Angelo AT. Perceptions of obstetric analgesia: a qualitative study among midwives attending normal vaginal deliveries in Durame Hospital, Southern Ethiopia. *J Pain Res*. 2019;12:2187–92.
- Bishaw KA, Sendo EG, Abebe WS. Knowledge, and use of labor pain relief methods and associated factors among obstetric caregivers at public health centers of East Gojjam zone, Amhara region, Ethiopia: a facility based cross-sectional study. *BMC Pregnancy Childbirth*. 2020;20:180.
- McCauley M, Stewart C, Kebede B. A survey of healthcare providers' knowledge and attitudes regarding pain relief in labor for women in Ethiopia. *BMC Pregnancy Childbirth*. 2017;17:56.
- CSA. *CSA Population Projections for Ethiopia 2007–2037*. Available from: <https://www.statsethiopia.gov.et/wp-content/uploads/2019/05/ICPS-Population-Projection-2007-2037-produced-in-2012.pdf>.
- Getu AA, Getie SA, Gela GB, et al. Non-pharmacological labor pain management and associated factor among skilled birth attendants in Amhara Regional State health institutions, Northwest Ethiopia. *Reprod Health*. 2020;17:183.
- Agaram R, Douglas MJ, McTaggart RA, et al. Inadequate pain relief with labor epidurals: a multivariate analysis of associated factors. *Int J Obstet Anesth*. 2009;18(1):10–14.
- Ogboli-Nwasor E, Adaji SE, Bature SB, et al. Pain relief in labor: a survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. *J Pain Res*. 2011;4:227–32.
- Ohaeri B, Owolabi G, Ingwu J. Skilled health attendants' knowledge and practice of pain management during labor in health care facilities in Ibadan, Nigeria. *Eur J Midwifery*. 2019;3:3.
- Sahile E, Yemaneh Y, Alemu AA, et al. Practice of labour pain management methods and associated factors among skilled attendants working at general hospitals in Tigray Region, North Ethiopia: hospital based cross-sectional study design. *Health Sci J*. 2017;11(4).

## **2. Journal of health science and medical research**

**Q4 : SJR 0,15**

**Thailand**



99+

Tulis

Mail

Kotak Masuk 1.721

Chat

Berbintang

Ditunda

Terkirim

Meet

Draf 2

Selengkapnya

Label

journal of health science and medical research

X

☰

▼

←

📁

🕒

🗑️

✉️

🕒

↶

📧

📄

⋮

🌟 Summarize

risnah@uin-alauddin.ac.id: Review request- ANPC-22-RA-178 (risnah@uin-alauddin.ac.id: Permohonan review- ANPC-22-RA-178)

Eksternal Kotak Mas

👤

**review.ptz@manuscript-submission.com** <review.ptz@manuscript-submission.com>  
kepada saya

🌐

Inggris → Indonesia  
[Tampilkan versi asli](#)

⚙️

Risnah yang terhormat,

Semoga Anda terhindar dari COVID-19 dan tetap sehat.

Saya menghubungi Anda untuk meninjau naskah berikut. Jika memungkinkan dan Anda tertarik, mohon sampaikan pendapat Anda untuk meningkatkan kualitas naskah dan kami dapat memberikan informasi yang bebas kesalahan kepada komunitas ilmiah.

*Nama Jurnal: Arsip Praktik dan Perawatan Keperawatan*

*Nomor naskah: ANPC-22-RA-178*

*Judul Naskah: Pengaruh Tai Chi terhadap Kualitas Hidup pada Lansia*

Harap beri tahu kami kesediaan Anda untuk meninjau naskah ini tanpa penundaan.

Respon dan dukungan cepat Anda akan sangat kami harapkan.

99+

Tulis

Mail

Kotak Masuk 1.721

Chat

Berbintang

Ditunda

Terkirim

Meet

Draf 2

Selengkapnya

Label

journal of health science and medical research

X

☰

▼

←

📁

🕒

🗑️

✉️

🕒

↶

📧

📄

⋮

✦ Summarize

Journal of Health Science and Medical Research - Account Created in Manuscripts (Jurnal Ilmu Kesehatan dan Penelitian Medis - Akun Dibuat Manuscripts)

Eksternal

Kotak Masuk x



**Wit Wichaidit** <onbehalf@manuscriptcentral.com>  
kepada saya

🌐 Inggris → Indonesia  
[Tampilkan versi asli](#)

⚙️

14 November 2023

Kepada Yth. Dr. Risnah:

Selamat datang di situs Jurnal Ilmu Kesehatan dan Penelitian Medis - ScholarOne Manuscripts untuk pengiriman dan p telah ditambahkan ke basis data peninjau kami dengan harapan Anda dapat meninjau naskah untuk Jurnal yang terma

ID PENGGUNA untuk akun Anda adalah sebagai berikut:

URL SITUS: <https://mc03.manuscriptcentral.com/jhsmr>ID PENGGUNA: [risnah@uin-alauddin.ac.id](mailto:risnah@uin-alauddin.ac.id)

PASSWORD: Untuk mengatur kata sandi Anda, silakan klik tautan di bawah ini. Dengan mengklik tautan tersebut, Anda mengatur kata sandi permanen Anda.

[https://mc03.manuscriptcentral.com/jhsmr?URI\\_MASK=1448a98687474b97aa59cd41de1e01aa](https://mc03.manuscriptcentral.com/jhsmr?URI_MASK=1448a98687474b97aa59cd41de1e01aa)



✦ Summarize

Label



Saya menyadari bahwa peninjau ahli kami memberikan kontribusi besar terhadap standar tinggi Jurnal ini, dan saya be







✦ Summarize

Eksternal Kotak Masuk x

Label

**Web of Science researcher profiles** [noreply@webofscience.com](mailto:noreply@webofscience.com) [lewat](#) [amazonses.com](#)  
kepada saya

 Inggris → Indonesia  
[Tampilkan versi asli](#)



Ulasan terkini Anda tentang "Pencegahan Praktik Manipulasi Data Kecelakaan di Sektor Pertambangan Batubara Menggunakan Aplikasi Seluler di Kalimantan Timur, Indonesia" Jurnal Ilmu Kesehatan dan Penelitian Medis telah secara otomatis ditambahkan ke [pro/ Web of Science](#) Anda sebagai bagian dari kemitraan kami dengan Universitas Pangeran Songkla.



**Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia**

Journal:	<i>Journal of Health Science and Medical Research</i>
Manuscript ID	JHSMR-2023-0601
Manuscript Type:	Original Article
Keywords:	Accident prevention, Coal mining, Manipulation of data, Mobile application
Speciality:	Occupational health and safety, Accident reporting, Coal mining, Zero accident

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia**

**ABSTRACT**

**Objective:** The practice of manipulating data on work accidents and potential hazards still occurs, especially in the coal mining sector. A mobile application for reporting work accidents and potential hazards is needed so that data manipulation practices can be prevented.

**Materials and Methods:** The research design used was exploratory which aimed to introduce and explore the capabilities of the SIKAP application (Information System for Occupational Accidents, Occupational Diseases and Potential Hazards) to prevent data manipulation practices. Researchers involved thirty coal mine workers using the SIKAP application which had been tested on three information systems experts using the black box testing method.

**Results:** Every worker has the opportunity to report work accidents and potential dangers via the SIKAP application on the worker's account. Companies can receive and follow up on worker reports via the company account. Regional governments in the field of labor development and inspection can monitor reports through government accounts. The SIKAP application is integrated and synchronized between workers, companies and the government.

**Conclusion:** The SIKAP application is able to prevent the practice of manipulating work accident reporting data and potential hazards in the coal mining sector.

**Keywords:** Accident prevention, Coal mining, Manipulation of data, Mobile application



## Introduction

Coal companies in Indonesia are still one of the most lucrative business sectors and have even become one of the favorite business sectors in East Kalimantan to this day. However, the coal mining sector is also one of the sectors that contributes to the high number of work accidents and work-related diseases. It is known that all stages of mining activities contain dangers that have a high risk of causing workers to suffer minor, serious injuries and even death.

The Indonesian government continues to make improvements so that the mining sector is no longer a scary place to work, especially for mining workers. One of them is by issuing regulations such as the Regulation of the Minister of Energy and Mineral Resources of the Republic of Indonesia Number 26 of 2018 concerning the Implementation of Good Mining Principles and Supervision of Mineral and Coal Mining, and followed by the Decree of the Directorate General of Minerals and Coal Number 185.K/ 37.04/DJB/2019 concerning Technical Guidelines for the Implementation of Mining Safety and Implementation, Assessment and Reporting of Mineral and Coal Mining Safety Management Systems.

Apart from the government's role, companies also have the responsibility to implement occupational safety and health (OSH) efforts to ensure that workers continue to receive protection and safety while working. This has been regulated in Law Number 13 of 2003 concerning Employment<sup>1</sup>, and Government Regulation Number 50 of 2012 concerning the Implementation of an Occupational Safety and Health Management System which is integrated with the company management system.<sup>2</sup>

One of the efforts to prevent work accidents and work-related diseases is through reporting work accidents, work-related diseases and potential dangers, both routine reports and special reports when work accidents occur. This reporting contains a number of information related to work accidents,<sup>3</sup> and this information will help company management to carry out immediate control over potential dangers so as not to result in work accidents in the future.<sup>4</sup> However, in practice there are still problems with reporting that is not transparent and tends to be manipulative.

Space for the practice of manipulating data on work accidents and work-related illnesses reported from companies to the government as the agency supervising coal mining operations remains wide open. One of the reasons is that the reporting system used is still conventional, utilizing written work accident and occupational disease reporting forms.<sup>5</sup> The practice of manipulating report data can not only be carried out by workers, but also company K3 officers and/or government employees.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

A real-time, integrated and synchronized information system for reporting work accidents, work-related illnesses and potential hazards is needed to prevent the practice of data manipulation when reporting is carried out. This is also in accordance with previous research recommendations which stated that one of the efforts to control the practice of manipulating work accident and occupational disease data is to utilize a digitalization-based reporting information system.<sup>6</sup> Based on existing problems, research is needed to introduce and explore the capabilities of the SIKAP application to prevent data manipulation practices in coal mining in East Kalimantan, Indonesia.

**Methods**

This research uses an exploratory design by utilizing the SIKAP application in the reporting system for work accidents, work-related diseases and potential hazards in coal mining. This research involved thirty coal mining workers from several coal companies operating in Samarinda City, East Kalimantan Province, Indonesia. The workers involved were given the SIKAP reporting application which had previously been tested on three information systems experts using the black box testing method to test that the SIKAP application could be used properly. Every worker is given the opportunity to use the SIKAP application to report work accidents, work-related illnesses, and potential dangers that have been experienced or seen while working at a coal company. The research was carried out in October 2023 based on a permit to carry out research issued by the Faculty of Economics and Business, Mulawarman University Number: 202/UN17.1.36.1/DT/2023. The results of trials using the SIKAP application on coal mining workers were then analyzed to assess the ability of the SIKAP application to prevent the practice of manipulating work accident reporting data, work-related diseases and potential hazards in coal mining.

**Result**

The SIKAP application is an acronym for Information System for Reporting Work Accidents, Occupational Diseases and Potential Hazards. The initial display when opening the SIKAP application consists of information about SIKAP, information about OSH activities, OSH policies, and other information related to OSH. There is an access menu at the bottom of the information consisting of worker, company and government accounts.

The worker account is preceded by creating a username, name, work unit, position, cellphone number, uploading a photo, and account password. Next, the worker clicks Log in by entering the username and password for the account that has been created. After logging in,

several application menus will appear consisting of Home, Announcements, Notifications, and Profile. The Home menu consists of Today's Data, Work Accident Report, Occupational Disease Report, Potential Hazard Report, and Report History.

Today's data includes data on work accidents, occupational disease, and potential dangers that have been reported by the worker and other workers at the company where he works. If the worker wants to report a work accident, then click the Report Work Accident menu. In this menu, workers enter a number of information, namely the type of work accident, victims, date of incident, time of incident, location of incident, body parts injured as a result of the accident, and actions taken by workers when experiencing an accident by attaching documentation/photos of the time of the incident. After completing all the required information, the worker then clicks Report.

For workers who want to report work-related illnesses, click Report of occupational disease. In the occupational disease report menu, a number of information will be entered by the worker, namely the type of occupational disease, details of the occupational disease, the complaint felt, the date and time of the complaint, and the actions taken by the worker when experiencing the complaint accompanied by documentation/photos. After completing all the required information, the worker then clicks Report. If the worker wants to report a potential hazard, then click Report Potential Hazard and fill in some information, namely location, description of the type of hazard, risks, actions taken, and attach documentation/photos of witnessing the potential hazard. After completing all the required information, the worker then clicks Report.

Worker accounts also have a Report History menu. This menu functions to check/monitor the status or reporting history of work accidents, occupational disease, and potential hazards that have been carried out by workers to the company's OSH unit. Workers can see the progress of the reports that have been carried out. The history of Work Accident Reports, occupational disease, and Potential Hazards consists of the statuses Sent, Processed, and Completed/Rejected. In this section, workers can see a description of the accident, occupational disease and potential hazard report information that has been filled in when reporting.

The report that has been sent by the worker can then be seen that the report was processed/received by officers in the OSH unit accompanied by the identity of the report recipient. Information was also obtained regarding suggestions/directions from officers in the OSH unit to workers/reporters. Workers will also receive information that the report has been completed by the OSH officer, accompanied by a description of the final action report from the

OSH officer. In this section there is also information about rejection of reports made by OSH officers if the report is deemed to have been repeatedly reported or has been completed by OSH officers.

Available Announcements menu, workers will see schedules or other important agendas informed by the company's OSH officers. On the Notification menu, workers will receive notifications automatically from reports that have been made previously. The profile will display a number of workers' personal information, including name, work unit and position. Details of the worker's SIKAP account can be seen in Figure 1 below:

Creating and using a company account is almost the same as a worker account, namely creating a username, name, work unit, position, cellphone number, uploading photos and account password. After registering the account, the company OSH officer then clicks Log in by entering the username and password for the account that has been created. When you log in, several application menus will appear consisting of Home, Announcements, Notifications and Profile.

The Home menu consists of Today's Data, Incoming Report, Process Report, and Completed Report. Today's Data Menu which includes data on work accidents, occupational disease, and potential hazards that have been reported by workers at the company. In this section, company OSH officers can search for accident, occupational disease and potential hazard data, both daily data and data according to the desired time period by clicking Search Data. The desired data can be displayed in graphical form by clicking the Graph button.

In the Incoming Reports, Process Reports and Completed Reports menu, company OSH officers can see a number of incoming report data from workers. OSH officers can also accept or reject reports from workers by providing directions/information addressed to the worker. In the Announcements menu, OSH officers can delete or add other important events/agenda which will be informed to workers. In the Notification menu, the company's OSH officer will receive notification automatically of each stage of incoming reporting and the profile will display a number of company OSH officer information. Details of the company's SIKAP account can be seen in Figure 2 below:

Creating a government account represented by certain service employees who are given the task of regarding accident, occupational disease, and potential danger data by creating a username, name, work unit, position, cellphone number, uploading photos and account password. After registering an account, employees in certain departments then click Log in by entering the username and password for the account that has been created. After logging in, several application menus will appear consisting of Home, Notifications and Profile.

The Home menu consists of Today's Data and Reports. Today's data includes data on work accidents, occupational disease, and potential dangers that have been reported by workers at certain companies. In this section, certain service employees can search for accident, occupational disease and potential hazard data, both daily data and data according to the specific time period desired by clicking Search Data. The desired data can be displayed in graphical form by clicking the Graph button. In the Reports menu, accident, occupational disease and potential hazard reports will be displayed in the Incoming, Processing and Completed Reports menu.

Certain service employees can view a number of incoming report data from workers at certain companies along with the history of each report. Service employees can also provide suggestions/input aimed at the company regarding incidents reported by their workers. In the Notification menu, certain service employees will receive automatic notification of each reporting process/stage that is entered into the SIKAP account belonging to a certain service. The profile will display a number of information about certain service employees who handle work accident data, occupational disease and potential hazards. Details of the government's SIKAP account can be seen in Figure 3 below:

## Discussion

The Indonesian government continues to make improvements to improve the coal mining sector as one of the sectors at risk of work accidents and work-related diseases through regulations. In several countries such as China and Australia, regulations regarding coal mine safety have been developed.<sup>7</sup> Regulations that are consistent in their implementation can influence workers' behavior at work.<sup>8</sup>

Reporting of work accidents, work-related diseases and potential hazards in the workplace, including in the coal mining sector, has been regulated in Minister of Manpower Regulation Number 3 of 1998 concerning Procedures for Reporting and Inspecting Accidents and has been used since 1998 until now using a written report format and an online reporting system. conventional from the company to the local regional government and the Ministry of Manpower of the Republic of Indonesia. Such reporting management has a number of weaknesses, such as the reporting process being slow, the possibility of incomplete information being obtained due to the disappearance of witnesses and loss of evidence when a work accident occurs, the unavailability of forms at the scene of the incident, and various other weaknesses.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Incomplete information regarding work accidents, work-related diseases and potential workplace hazards will have a negative impact on the company's OSH performance. It is difficult for company management to take action to prevent and handle work accidents and work-related diseases if they are not supported by detailed data and information regarding the root causes or potential hazards that result in work accidents and work-related diseases. On the other hand, having complete information about potential hazards in the work environment makes it easier for companies to identify work areas that are at risk of work accidents and proceed with corrective action so that similar incidents do not happen again.<sup>9</sup>

Reporting work accidents, work-related diseases and potential hazards is not only the company's responsibility to the government, especially when work accidents and work-related diseases occur, but also requires worker participation in reporting.<sup>10</sup> This has also been regulated based on Decree of the Directorate General of Minerals and Coal Number 185.K/37.04/DJB/2019 that every mining worker is obliged to report any accidents or injuries caused by work or related to work. However, it is still found that mining workers think that reporting work accidents and potential dangers in the workplace is not important, so it is necessary to increase workers' perceptions about the importance of reporting.<sup>11</sup>

Work accident reporting aims to make it easier for company management to identify work areas that allow work accidents,<sup>9</sup> and at the same time identify the causes of accidents so that work accidents can be prevented early on.<sup>12</sup> In addition, Work accident reports can also estimate workers' unsafe actions.<sup>13</sup> However, the K3 efforts carried out are still less effective in preventing work accidents, so a better investigation system is needed.<sup>14</sup> Reporting coal mining accidents can also improve safety management in the sector mining.<sup>15</sup>

Reporting accidents and potential dangers by workers needs to receive support from the company.<sup>16</sup> Workers' participation in reporting work accidents is not enough to create an K3 culture in the workplace, but the role of the company is also needed.<sup>17</sup> Companies that have a high commitment to safety are positively correlated with workers' K3 behavior.<sup>18</sup> Solutions to preventing work accidents and occupational diseases require partnerships and synergy between workers, government, companies and other parties.<sup>19</sup>

Challenge faced in reporting work accidents, work-related diseases and potential hazards in coal mining is the large number of cases that occur but are hidden or not reported.<sup>20</sup> The practice of hiding, deleting data, changing some or all of the data is an act of manipulation that will harm various parties, especially workers. Likewise, companies will experience difficulty in following up on OSH improvement efforts. Therefore, a digitalization-based information system is needed in the form of an application for reporting work accidents, work-



related illnesses and potential hazards in coal mining. The information produced from the reporting system becomes the basis for companies to make appropriate decisions,<sup>21</sup> and has a positive impact on worker performance.<sup>22</sup> The use of mining information technology has a positive impact on increasing safety and productivity.<sup>23</sup>

The question is, is the SIKAP application able to prevent the practice of manipulating data on accidents, occupational disease and potential hazards in coal mining? The SIKAP application consists of three user groups, namely workers, companies and government. This is intended so that the reporting carried out does not only involve workers and companies for internal purposes, but also involves the government so that reports of work accidents, work-related illnesses and potential hazards in coal mining can be jointly accounted for by all parties related to accident data, work and potential hazards.

Every worker has the opportunity to report any work accidents, work-related illnesses and potential dangers experienced by themselves or their co-workers through the worker's account in the SIKAP application. Mining workers are able to find out in real-time the number of reports of work accidents, work-related illnesses and potential hazards that are entered into the application, whether reported by themselves or reports from other workers. Apart from that, workers have the opportunity to monitor the progress of each report made, so that it is not just reporting, but there is attention and follow-up from the company.

Reports of work accidents, work-related illnesses and potential hazards carried out by workers will be connected and synchronized with the SIKAP account owned by the company and government. Companies can follow up on reports of potential dangers reported by workers with various efforts or programs to prevent these potential dangers from causing work accidents and work-related diseases. Likewise, reports of work accidents and work-related illnesses reported in the SIKAP application are then followed up by the company in the form of immediate treatment so as not to cause more serious injuries.

The reporting data received by the government is also real-time so that the government can know the total data on work accidents, work-related diseases and potential hazards that have been carried out by workers. Apart from that, the government can also find out about the handling actions that have been taken by the company regarding work accidents and work-related illnesses experienced by workers. This means that there is no difference in the amount of data reported by workers and the data that will be received in company and government accounts. The compatibility of data received between workers, companies and the government will close the space for anyone to carry out data manipulation practices in the future, especially when companies make periodic reports and special reports to the government.

**Conclusion**

The SIKAP application is an innovative step in the reporting system for work accidents, work-related illnesses and potential hazards to prevent the practice of manipulating report data in coal mining in Indonesia.


**References**

1. Indonesia, R. Law of the Republic of Indonesia No.13 of 2003 concerning Employment. *Constitution. No.13 of 2003* (2003).
2. Indonesia, P. R. Republic of Indonesia Government Regulation Number 50 of 2012 concerning the Implementation of Occupational Health and Safety Management System. *President of Republik Indonesia* (2012).
3. Heraghty, D., Dekker, S. & Rae, A. Accident report interpretation. *Safety* **4**, 46 (2018).
4. Sneddon, A., Mearns, K. & Flin, R. Stress, fatigue, situation awareness and safety in offshore drilling crews. *Saf. Sci.* (2013) doi:10.1016/j.ssci.2012.05.027.
5. Molinero Ruiz, E., Pitarque, S., Fondevila-McDonald, Y. & Martin-Bustamante, M. How Reliable and Valid is the Coding of the Variables of the European Statistics on Accidents at Work (ESAW)? A Need to Improve Preventive Public Policies. *Saf. Sci.* (2015) doi:10.1016/j.ssci.2015.05.005.
6. Sultan, M. Collaboration Tripartite Elements Accident Reporting and Potential Hazards Management Coal Mining, Indonesia: Systematic Review. *J. Heal. Sci. Med. Res.* **41**, 1–5 (2023).
7. Quanlong Liu et al. Regulatory Regime on Coal Mine Safety in China and Australia: Comparative Analysis and Overall Findings. *Resour. Policy* **74**, (2021).
8. Wang, X., Zhang, C., Deng, J., Su, C. & Gao, Z. Analysis of Factors Influencing Miners Unsafe Behaviors in Intelligent Mines using a Novel Hybrid MCDM Model. *Int. J. Environ. Res. Public Health* **19**, (2022).




9. Weilli Fang et al. Automated text classification of near-misses from safety reports: An improved deep learning approach. *Adv. Eng. Informatics* **44**, (2020).
10. Winkler, M., Perlman, Y., & Westreich, S. Reporting Near-Miss Safety Events: Impacts and Decision-Making Analysis. *Saf. Sci.* **117**, 365–374 (2019).
11. Sultan, M., Setyadi, D. & Ramdan, I. M. Work Accident Reporting in Coal Mining, Indonesia: A Systematic Literature Review. *P.O.S.H Period. Occup. Saf. Heal.* **2**, 51–59 (2023).
12. Qiao, W., Li, X. & Liu, Q. Systemic approaches to incident analysis in coal mines: Comparison of the STAMP, FRAM and “2–4” models. *Resour. Policy* (2019) doi:10.1016/j.resourpol.2019.101453.
13. Kumara et al. Estimation of Human Error Rate in Underground Coal Mines Through Retrospective Analysis of Mining Accident Reports and Some Error Reduction Strategies. *Saf. Sci.* **123**, (2020).
14. Dash, A. K., Bhattacharjee, R. M., Ahmad, A., Sagesh, K. M. R. & Singh, C. S. Mine accident investigation in India: A system approach. *J. Mines, Met. Fuels* (2018).
15. Huang, Y. & Zhou, Q. Mine accident prediction and analysis based on multimedia big data. *Multimed. Tools Appl.* (2019) doi:10.1007/s11042-019-7175-6.
16. Westreich, M. W. Y. P. S. Reporting near-miss safety events: Impacts and decision-making analysis. *Saf. Sci.* **117**, 365–374 (2019).
17. Tetzlaff, Emily J., et al. Safety Culture: A Retrospective Analysis of Occupational Health and Safety Mining Reports. *Saf. Health Work* **12**, 201–208 (2021).
18. Xinfeng et al. The Mediating Role of Psychological Capital Between Perceived Management Commitment and Safety Behavior. *J. Safety Res.* **72**, 29–40 (2020).
19. Stewart, A. G. Mining is bad for health: a Voyage of Discovery. *Environ. Geochem. Health* **42**, 1153–1165 (2020).
20. Yang Xiuyun, Kees Krul, and D. S. Uncovering Coal Mining Accident Coverups: An Alternative Perspective on China’s New Safety Narrative. *Saf. Sci.* **148**, (2022).
21. Nugroho, A. S. *Management Information Systems: A Practitioner's Review of Information Technology*. (Teknosain, 2018).
22. Abualoush et al. The Role of Employees’ Empowerment as an Intermediary Variable Between Knowledge Management and Information Systems on Employees’ Performance. *VINE J. Inf. Knowl. Manag. Syst.* **48**, 217–237 (2018).
23. Yang, L., Birhane, G. E., Zhu, J. & Geng, J. Mining Employees Safety and the Application of Information Technology in Coal Mining: Review. *Front. Public Heal.* **9**, 1–12


1  
2  
3 330 (2021).  
4  
5 331  
6  
7 332



333  
334  
335 **Figure 1.** Display of worker accounts in the SIKAP application  
336  
337



338  
339  
340 **Figure 2.** Display of company accounts in the SIKAP application  
341



**Figure 3.** Display of government accounts in the SIKAP application

**3. Belitung Nursing  
Journal  
Q3 : SJR 0,3  
Indonesia**

99+

Tulis

Mail

Kotak Masuk 1.721

Berbintang

Chat

Ditunda

Terkirim

Meet

Draf 2

Selengkapnya

Label

joko gunawan

←

📧

🕒

🗑️

📧

🕒

🔄


📧

📧

⋮

✦ Summa

[BNJ] Article Review Request ([BNJ] Permintaan Ulasan Artikel) Eksterna



**Joko Gunawan, S.Kep. Ners., PhD** <editorbnj@gmail.com>  
kepada saya

🌐

Inggris → Indonesia  
[Tampilkan versi asli](#)

⚙️

Risnah:

Saya yakin Anda akan menjadi pengulas yang baik untuk naskah, "Praktik Kesehatan di antara Lansia Muslim yang Tir Selatan: Praktik Kesehatan di antara Lansia Muslim yang Tinggal di Rumah," yang telah diserahkan ke Jurnal Keperaw disisipkan di bawah ini, dan saya harap Anda akan mempertimbangkan untuk melakukan tugas penting ini bagi kami.

Silakan masuk ke situs web jurnal paling lambat 2022-11-29 untuk menunjukkan apakah Anda akan melakukan peninj kiriman dan mencatat tinjauan dan rekomendasi Anda.

Tinjauan itu sendiri akan jatuh tempo pada 2022-11-29.

URL pengiriman: <https://www.beliitungraya.org/BRP/index.php/bnj/reviewer/submission?submissionId=2406&reviewId=>

Jika Anda meninjau naskah ini, Anda akan memenuhi syarat untuk mendapatkan diskon 10% atas biaya pemrosesan s ke BNJ. Diskon ini harus diklaim pada saat penyerahan naskah dan berlaku selama satu tahun sejak selesainya peninj dapat diklaim atas naskah yang dikirimkan dan tidak dapat digabungkan dengan diskon lainnya.

99+

Tulis

Mail

Kotak Masuk1.721

Berbintang

Ditunda

Terkirim

Draf2

Selengkapnya

Chat

Meet

Label

joko gunawan

X

≡

▼

←

📁

🕒

🗑️

✉️

🕒

↶


📧

📄

⋮

🌟 Summa

[BNJ] Validate Your Account ([BNJ] Validasi Akun Anda) Eksternal Kotak Ma

 **Dr. Joko Gunawan** <editorbnj@gmail.com> kepada saya

🌐

Inggris → Indonesia

Tampilkan versi asli

⚙️

Risnah

Anda telah membuat akun di Belitung Nursing Journal, tetapi sebelum dapat menggunakannya, Anda perlu memvalida melakukannya, cukup ikuti tautan di bawah ini:  
<https://www.belitungraya.org/BRP/index.php/bnj/user/activateUser/risnah/mw67yF>

Terima kasih,

**Kantor Redaksi Belitung Raya**  
Departemen Publikasi, Yayasan Belitung Raya, Indonesia

Jurnal Keperawatan Belitung - Kantor Redaksi Belitung Raya

https://mail.google.com/mail/u/0/#search/joko+gunawan/FMfcgzGqRGWTWsClclSzNQDvsmhDBfdB

1/1

99+

Tulis

Mail

Kotak Masuk 1.721

Berbintang

Ditunda

Terkirim

Draf 2

Selengkapnya

Meet

Label

Q joko gunawan

X

≡

▼

←

📧

🕒

🗑️

📧

🕒

🔄

📧

📧

⋮

✦ Summa

## [BNJ] Article Review Acknowledgement ([BNJ] Ucapan Terima Kasih a Artikel) Eksternal Kotak Masuk x



**Joko Gunawan, S.Kep. Ners., PhD** <editorbnj@gmail.com>  
kepada saya



Inggris → Indonesia  
[Tampilkan versi asli](#)



Risnah yang terhormat,

Terima kasih telah menyelesaikan tinjauan atas kiriman, "Praktik Kesehatan di antara Lansia Muslim yang Tinggal di R Praktik Kesehatan di antara Lansia Muslim yang Tinggal di Rumah," untuk Jurnal Keperawatan Belitung. Kami mengha karya yang kami terbitkan.

Harap dicatat bahwa Anda kini berhak atas diskon 15% untuk biaya pemrosesan satu artikel untuk naskah yang dikirim ini harus diklaim pada saat penyerahan naskah dan berlaku selama satu tahun sejak selesainya peninjauan Anda. Har naskah yang dikirimkan dan tidak dapat digabungkan dengan diskon lainnya.

**Belitung Nursing Journal** adalah [anggota Reviewer Credits](#). Menyelesaikan peer-review yang Anda ikuti akan memt mendapatkan sertifikasi pihak ketiga yang disediakan oleh organisasi ini dan memberi Anda kredit yang dapat digunak. Daftarkan akun Reviewer Credits gratis sekarang dan bersiaplah untuk memulai. Cari tahu lebih lanjut di [SINI](#).

Salam,

...

## **Health practice among Muslim homebound older adults living in the Southern Thai community**

### **Abstract**

**Backgrounds:** Age friendly environment is helpful to promote older people's health practice and healthy aging. However, little is known about those elders living at home in a Thai Muslim community.

**Objective:** This is a part of the ethnographic study that aimed to explore health care practices of Thai Muslim Homebound older adult (HOA) based on beliefs and their experiences to maintain their holistic health in southern Thai Muslim context.

**Methods:** Purposive and snowball sampling were used to recruit 15 HOAs as key informants among whom 9 were living in an urban area and 6 living in a rural area. Data were collected using in-depth interview, participant observation, and field notes and analyzed using a four-step qualitative data analysis.

**Results:** The results showed that Muslim HOAs performed their health practice based on traditional health practice related to Muslim culture under the theme of life and health are designated by Allah for living with nature and comforts with their age. Every participant used religion to lead their practice, a natural way based on warmly environment, and integrated both folk and modern medicine. Various modalities for health care such as traditional massage, oil massage and herbal therapy following the Quran teaching were used. Health related religious practice was the main method and their practice regularly performed by all HOAs under the belief that religious practice would help them maintain both healthiness and happiness. Muslim prayer was believed to improve physical function of all body and personal hygiene. In addition,



their health practices were supported from their family, community health workers, and health volunteers including their Allah as a spiritual retreat.

**Conclusions:** Knowledge of cultural health practice would help nurses especially primary care professionals to promote self-care and provide culturally sensitive nursing care to maintain the holistic health of Muslim elders in long term.

**Keywords:** Homebound older adult, Health practice, Muslim cultural care

## **Introduction**

The increased number of older adults around the world affects the change in the structure of the population. In 2020, 13 percent of the world's population were aged over 60 years with an increase of 3% per year and the highest rank was in Asia since 2005 (United Nations, 2020). Thailand is one of the highest ranks where the older population accounts for 18 percent of the total population reported. Among this number, 21 percent are living with dependency conditions. Of these, 15% are the homebound older adults (Foundation of Thai Gerontology Research and Development Institute (TGRI), 2019)

The World Health Organization (WHO) has urged a strategy on healthy ageing and defined it as ‘the process of developing and maintaining the functional ability that enables well-being in older age’ (World Health Organization, 2017). This has necessitated the intervention to promote the health of older adults and prevent the health problems related to the aging process, chronic illness, and inability to perform daily activities. A previous study showed that many older people living at home face some physical health conditions which may deteriorate into being bed-ridden (Husebo & Storm, 2014). In addition, the deterioration of physical condition and chronic diseases is the main cause affecting quality of life of the older adults particularly

those who are homebound. Therefore, the homebound older are classified as a significant group regarding the great need of care to slow or prevent the deterioration of their physical condition and who require a special health service system in accordance with their health needs.

Research suggests that the development of an age friendly environment is necessary to promote health within and around living areas. Each older person is culturally different in lifestyle including social structure, for example the living characteristics and social conditions of urban and rural communities differ in terms of housing. The older people in a rural community often live together with their children in a community, helping each other in a Thai culture, and having traditions and beliefs that have been passed on from ancestors from generation to generation. The older adults are valuable social capital and affect the minds of their children, and have their own wisdom, while the urban community has a modular style of living. However, the living condition may be in different patterns such as many children leave home for work outside and leave the elders at home alone or living with their spouse without support. Therefore, it affects the quality of life and mental state of all the older adults (Foundation of Thai Gerontology Research and Development Institute, 2021). In addition, Thailand has prepared a 20-year National Strategic Plan (Public Health section) which focuses on people's life span, from youth up until old age. Its goal is to enable older adults to take care of themselves in 'active and healthy aging' (Ministry of Public Health, 2016). If health or social professionals have a better recognition of older people and their way of life in a community, a more 'health in the older adults targeted approach' could be applied to address health and social problems and the needs of senior citizens. These working methods should be both culturally sensitive and effective (Leininger & McFarland, 2002).

However, a previous study in one of the southern Thai provinces, showed that the Thai Buddhist homebound older adults were able to take care of themselves at home despite some health limitations (Detthippornpong, Songwathana, & Bourbonnais, 2022). Some perceived themselves as a healthy group and able to control their illness by taking self-care, sharing a positive view of life with others, and integrating their beliefs on folk with modern medicine to maintain health. Supportive family networks and Thai cultural beliefs on respecting older people in the community and participating in religious activities were identified as facilitating factors. However, the previous study was limited to Thai Buddhists, which suggests a need to explore another ethnic group, such as the older Thai Muslim population in southern Thailand. This study aimed to explore cultural care practices related to the health and wellness of Thai Muslim homebound older adults.

### **Health care culture in a Muslim context**

For the Muslim health care culture, it is often consistent with religious principles at each stage of life as a way of life, especially in the mental and spiritual aspects of Muslims who have faith in God. Muslims regard the body as a gift from God. Therefore, one has a duty to care of the body and use it within the limits to maintain good health. When individual is ill, the sickness needs to be healed. The recovery from sickness will depend on God's schedule because Muslims believe sickness that arises is ordained by God to test that person's strength (Mohmula, 2007). In addition, for a healthy person, there must be a good relationship between man and God, and between self and environment. Muslims then focus primarily on spiritual well-being under the belief that it leads to better overall well-being.

Many Thai Muslims believe that spirit is the power of life and it is taken as a goal in a person's daily life. Spiritual wellness is the power to hold the mind that creates motivation and hope, awareness of one's self-worth and self-confidence, and leads to interactions with others and the environment, and brings life satisfaction (Chansangrat, Phibal, Niltaya, Thiratchakul, & Hatthakit, 2014; Rojanadhamkul, 2020). In addition, Muslim religious health practice has established measures at the individual and community levels as a guideline for prevention and promotion of health, and consist of 3 aspects related to good health, namely cleanliness, exercise, and food consumption (Arihathairat, 2020). Religious practices are often aligned with Muslim lifestyles to become cultural health care. Islam believes that the actual health starts with a strong and stable spirit, that is, the power to make decisions for oneself and able to maintain one's own independently without falling to emotional desires or material values. To achieve holistic health, the value of life and health must be firmly contained in the soul, not the value seen by others except the value guided by Allah (Iman), followed by the conversion of value into action (Amal), which is the composition of "Ibadah" (Lohwithee, 2020)

Ibadah is something that must be done on a regular basis in order to cultivate spiritual strength. It is an integrated system of health and life that covers all aspects of body, spirit, individuality, and society in worshiping which must arise from a spirit of fidelity to Allah and a traditional spirit or the Five Principles (Alabdulwahab, Kachanathu, & Oluseye, 2013). It consists of 1) taking an oath to God (Shahaada), 2) praying, 3) fasting, 4) zakat, and 5) pilgrimage to Makkah (Hajj). It has been linked to Muslim health conditions such as prayer, a physical and mental reverence for God. It should be performed from the age of about 7 years onwards, 5 times a day. The activity during praying helps to increase the flexibility of joints and bones in the body, which helps to create a balance in the body such as balance due to the

movement of all body parts. When doing it on a regular basis, it could stimulate of various organs such as the neck, back, legs, muscles, joints, etc. (Alabdulwahab, Kachanathu, & Oluseye, 2013) and help the mind to have mental stability. In addition, to reduce the anxiety and suffering, praying zakat is a part of purifying the mind to have a shared mind (Arihathairat, 2020), helping the individual to follow human beings according to one's strength and one's determination to share things with others.

Previous studies focusing on experiences among older people living at home showed that they often enjoyed life and wished to continue life as long as possible, but those limited to active people. However, most HOAs who are virtually isolated and may have limited of mobility should be taken in consideration for appropriate strategies in providing services for health promotion. This study then focused on those Muslim homebound older adults living in a different sociocultural context from Thai Buddhist.

## **Objective**

To explore cultural care practice related to health and wellness of Thai Muslim homebound older adults.

## **Conceptual framework**

The concept of holistic health nursing (Dossey & Keegan, 2016) and Leininger's culture care diversity and universality theory (Leininger & McFarland, 2006) informed this study. According to this theory, the goal of nursing is to offer culturally congruent care for the benefit of people's health and well-being. Leininger defines a culture as a set of beliefs, values, norms, and lifeways shared and learned by a group of people. In her theory, a cultural group shares folk

and professional care systems which can be mediated by nursing care (Leininger & McFarland, 2006). It also described various cultural and social dimensions that can influence health in a cultural group, for example, technological, religious, or economic factors. The meanings of health and those dimensions present differences and similarities between cultures (Leininger & McFarland, 2002). Another concept of Leininger's theory is caring which considered the essence of nursing. It involves understanding and knowing about people being cared for by helping them improve their human condition and health. This understanding can only be developed by considering their cultural context. It allows nurses to take actions to preserve, accommodate or repattern cultural care (Leininger & McFarland, 2002) to improve their health.

Based on this theory, the health practice among Thai Muslim HOA in Southern Thailand is influenced by cultural dimensions. To provide culturally congruent care, nurses need to better understand those health practices, particularly from the views of folk or traditional care systems, and the influencing religious beliefs, to better support their holistic health and well-being. This theoretical perspective guided the various methodological aspects of this study such as sampling, data collection and analysis.

## **Method**

This ethnographic research aimed to explore cultural care among the Muslim older adults at home in order to maintain holistic health. The study was conducted in two subdistricts (one in urban and another from rural area) in Songkhla province. The areas were selected based on the 2020 annual report of Songkhla Public Health office as which reported them as having the highest number of Muslim older adults and where the Subdistrict Administrative Organization identified more number of cases who were partially dependent and were mainly living at home.

In this study, homebound older adults were approached as key participants. To ensure that those participants involved could provide rich information and were willing to participate in this study, we used participant observation by visiting the primary health center in the study site and joining with the community nurses and health volunteer, to gain access to information related to health records of the older population. The purposive sampling to recruit 15 key informants, selection criteria which included 1) a homebound older person, as defined based on the National Health Security Office: NHSO (2016), with a Barthel Index score between 5-11, 2) being a Thai Muslim older person (age over 60) who was able to take care or move with ease by his or herself, mainly living at home, 3) being healthy according to his or her own perception although there may be some underlying chronic diseases but no history of complications or hospital admission for at least one year, and 4) able to speak and understand the Thai language or southern Thai dialect. To obtain in depth information, the associate informants (family members) who were well recognized or referred by the key informants were approached to explain and clarify the studied phenomenon.

### **Research Instrument**

Semi-structured interview guideline was mainly used to capture in-depth data from both key and associate informants. The interview guideline was developed by the team based on the literature review and Leininger's transcultural care theory (Leininger & McFarland, 2002). The interview format was based on open-ended questions to encourage extended response. The general questions about the lives of participants were usually used at the start of each interview and then the meaning of health and care for themselves in everyday life situations were probed. For example, "What do you think about your health and what do you practice in daily life and

why do you do those activities, how was life and health perceived in your age?”. The associate participants were asked about the daily life and activities usually performed to support key informant’s health or maintain their healthy. Each of the interviews lasted 45-60 minutes. A group interview with Muslim health professionals took place at the local hospital to gain more knowledge about the health services offered to Muslim older people in the study site. All instruments modified from the previous study in a Thai Buddhist homebound group (Detthippornpong, Songwathana, Bourbonnais, 2022 ) were used among the Thai Muslim older adults (see Appendix).

### **Human subject’s protection**

The study was approved by the Institutional Research Board Committee of the Faculty of Nursing, Prince of Songkla University (PSU IRB code 2021-LL-Nur011). The objectives of the study, research methods, and potential risks and benefits were described to the informants before the data collection. Permission from the informants for audio recording was obtained, and data were kept confidential on a personal laptop secured by a password. The participants were free and had time to decide whether to participate in the research. Verbal consent was used instead of written form owing to eye problem and hand writing. During participation, if the participants wished to leave or withdraw from the research before the end of the action, it could be done without giving any reason.

### **Data collection**

According to the research process, researchers and research assistants collected data as follows:



### 1. Gain entry to the community

The researcher contacted and coordinated with the nurses working at the two selected Sub-District Health Promoting Hospitals which had recorded many healthy Thai Muslim HOAs to explain the objective of the research project and data collection process. The Muslim older adults living at home were listed based on the community database and health report and then the participants were introduced by community/home health care nurses. Due to the pandemic during the community survey, local research assistants in each area (health care workers who have public health backgrounds) were trained about the qualitative approach and interview skills. A skill building workshop on ethnographic research was held at the community hospital focusing on qualitative techniques such as focus group discussion, semi-structure interview, and participant observation as well as content analysis.

### 2. Data collection

The informants were interviewed in their home with the help from family members who were living with and referred by participants when some who had difficulty to explain. The observation of the informants' behaviors and activities in their daily lives during interview was also conducted using diary/record forms. In the field, a personal diary, field notes, as well as an audio tape recorder and camera were used. The field notes recorded the first author's reflection during the interviews and the general context during the researcher's stay in the field. Data were collected between August 2021 and February 2022 through participant observation, and semi-structured in-depth interviews until the data were saturated.

In addition, a meeting was organized to present the progress of data collection and data analysis with research assistants through the ZOOM system and video conference for researchers to verify that the data were consistent with the research objectives. This enabled the data to be

analyzed and issues that needed additional information to be explored. During the data collection period, there was an outbreak of COVID-19, each household was aware and protected the older adults by keeping them at home and avoiding taking them to the doctor. Mask wearing, hand washing and keeping the distance were strictly performed by researchers as well as associated informants in the community.

### **Data analysis**

Data analysis was performed in parallel with data collection until the data were saturated. Thematic analysis was used to identify, analyze, and report patterns of the health practices within data in this ethnographic research. Four steps in ethnonursing data analysis were used (Leininger, 2006). First, researchers initially transcribed from in-depth verbatim interviews. The recorded tape and field note was listened repeatedly including the text used several times to understand the context or words that appeared to ensure that all information was covered. Second, descriptors and components were identified code and categorized after becoming familiar with the data set. For example, the daily dietary practice of eating steamed rice, soup, and a lot of fish, avoiding spicy food, drinking honey with warm water, were grouped into a potential subtheme of eating soft diet to improve digestive function. Similarly, other subthemes were searched such as praying and reading the Qur'an to restore mind, body and spiritual health. Third, the researcher analyzed patterns and contexts related to cultural care and health practice of the Muslim HOAs. Lastly, researcher synthesized the major theme and summarized the findings.

## **Trustworthiness**

Trustworthiness was provided by prolonged engagement in the field, maintaining field notes and a reflexive journal, peer debriefing and a form of data triangulation based on the concept of Lincoln and Guba (1985). Data were collected through different methods and different groups of participants. The researchers visited HOAs and their family a few times, at least 2 hours each visit, in their natural settings. Rapport and trust with the informants were established before the visits, which were arranged at different times of the day to gain various perspectives for the data. Field notes and a reflexive journal were maintained to describe the research process in detail and the researcher's role. Each step of the research process was appraised and confirmed by the team to check the accuracy of coding in data analysis, and to confirm the consistency of the inferences.

## **Results**

The findings were presented into two parts: 1) characteristics of the informants; 2) cultural care of health practice of Thai Muslim HOAs.

### **Characteristics of the informants**

Demographic characteristic of the 15 key informants, 6 males and 9 females, was presented in Table 1. Nine older adults in urban areas and 6 older adults in rural areas participated. Their age was between 60-94 years, most of them had activities of daily living (ADL) although some remained working in the rubber plantations. All lived at home with at least one of their family members and five of them lived with their wife. Most of them had at least one illness, the most common illnesses were hypertension and diabetes as shown in Table 1. In this

study, 6 family members who were also interviewed to gain more information based on the HOAs' referred persons were daughters (Ma1, Pa2, Ma6, Ma7, MaT2, MaT6).

### **Cultural care of health practices among Thai Muslim HOAs**

In describing their health, the HOAs lived at home defined health mostly related to their physical functioning, emotional and spiritual well-being. This definition included how these components affected their daily lives and their health. Although physical health and medical issues were part of living, participants viewed their health in a holistic way, more in terms of basic needs, close to nature within a simple and self-sufficient way. "Life and health are designated by Allah for living with a nature and comforts with their age" was perceived and guided their health practice as important for Muslim HOAs. Their daily health practices consisted of four patterns: 1) maintaining day to day functioning to stay independence, 2) having a simple and comfortable life with support, 3) performing religious activities as a priority of life for well-being, and 4) managing symptoms to gain a balance and restore health.

#### ***1. Maintaining day to day functioning to stay independence***

In daily life, participants described the importance of maintaining independence, which they considered as primary source to stay strong in health. They valued their ability to support themselves without burdening others. Activity of daily living (ADL) such as eating, bathing and dressing, were considered the most important of individual ability that should be maintained. Some talked about their self-care ability by keeping mobile as much they could in and out of their home. Some participants performed regular exercise as routine for their recovery from

illness. Some had ability to go out where they wanted (even with limitations), and enjoy their life when they felt they could do by themselves. As one participant who had just recovered from paralysis, she would walk around the house for an hour a day, keep kicking her legs, turning her arms, and flicking her arms every morning-evening or when she had a free time.

*... My former occupation was gardening, growing vegetables and trading, selling vegetables, bringing vegetables to sell at the market in the morning. After selling, I returned to gardening, planting fruits and vegetables, and having people come to home for body massage. Currently, I have stopped working because I'm old, my children do not allow me to leave the house since falling last year...Now, I try to help myself as much as I can, picking up things, moving to the bathroom, taking a shower when staying at home alone because I can't go anywhere. I try not to stand still, but try to exercise, move arms and legs, flicking leg, swinging arms, and kicking legs daily. I sometimes use massage oil to loosen the stretched muscles. (Ma1)*

Another participant also demonstrated her body movement by daily exercise on legs and arms after she stopped working.

*... I do exercise regularly, sit still and kick legs, turn around arms, flick arms every morning-evening or when I have free time...In the past, I sold fish at the market, working hard, I worked hard every day. After I had diabetes, I stopped selling stuff and stayed at home for the past 5 years... Always exercise to keep my body strength. (Ma5)*

In the rural area, some participants enjoyed working at farm by lifting muscles exercise, taking the cows for a walk in the early morning and evening, as a field note and interview:

*...I exercise every day after waking up. I do many of the same postures. I can't remember them all. There are body lifts, legs lifts, arms up. There are about 6 positions that the nurse has taught me. Do each position and move for 20 times. (PaT3)*

*...before...I was still strong, after wake up, I went out by bicycle in the morning every day, going back and forth 3-4 kilometers a day. I sometimes stopped some places for having a lunch. I think, if I did not exercise like that, I may not stay strong until now. (MaT6)*

## ***2. Performing religious activities as a priority of life for well-being***

Participants addressed the importance of religious activities, by accepting the laws of God's circumstances, believing their life and health as a destiny originated by God. The natural health care linked to divine blessings was performed daily with some modern medicine as necessary to restore health. All participants accepted their body's limitations and viewed the Muslim spiritual practices as important to life, health and death. They believe that through praying and reciting the Quran, enable individual to balance his/her own mind and body, gain spiritual well-being by achieving inner peace and wisdom and bring good health.

*... I pray 5 times and pray to God for good health. Reading Quran is necessary for all Muslims. I believe in the laws of God that giving birth, being old age, having sickness, and death are normal. I always relax the body, keeping no stress or worry about health problems... Every day I pray, sometimes recite the remembrance of God 1,000 times and read Yaseen (Chapter one of the Qur'an). Every morning, it helps me remember things to do with good mood. Many people say that I am a kind person, have never felt angry with anyone...and it helps to sleep better. (Ma1)*

*...During the illness, there is a period of stress, too much thinking, feel angry, irritability, troubling, self-inflicted. I have no strength at all after having a stroke. I need a great power from God to walk. must be a great power, not a little power. Let's look at others to heal. We feel empowered by God. Helping to keep getting better.... In the past, I couldn't sit and stand. I couldn't help myself... I can solve it by praying for blessings from God for good health, I pray and Zikr Ullah (Remembrance of God) every morning and evening 300 times. I then feel better...my mood has improved. (Pa4)*

In addition, adhering to religious doctrines is the culture where older people are regarded as the most important in later life when compared to other ages. Under the belief that practicing religious principles brings merit, having a happy mind to gain happiness in this world and the next and also have a long life. Practicing a good deed and faith in religion throughout life is essential. Then, strict practice on religious teachings as a way of life is helpful to enhance peacefulness of mind and happiness, as some participants reflected:

*... Allah said that by doing Prue (whatever way) we can combine our minds to Allah. You can lie or sit down, can think about it. You can go in your heart. Ask Allah and you will feel comfortable... Allah. To hurt, to be sick, to have a fever, to die is the duty of Allah. Allah is in control of everything.... Allah give rest when we have been working a lot to rest with grandchildren, I feel comfortable when thinking like this. It's time to go. Don't be afraid. It's time Allah takes you back. We are not capable of holding on to life. Allah takes everyone back and leaves no one behind. It depends on whether it's slow or fast. (PaT3)*

Moreover, the participants were concerned with having good hygiene habits consistent with religious teachings, such as eating quality and wholesome food, cleaning the body cleaning clothes. Muslim older adults staying at home tend to place personal hygiene as a top priority by

believing that cleanliness is the most basic form of health, which is a part of Muslim's obligation and responsibility to himself. Islam has defined the daily roles and duties of Muslims from waking up to going to bed to cover and commit to keep clean body, clothes, food and others, both at private and public settings. As some participants described:

*...After taking a bath and praying 5 times a day, use clean water to wash hands, mouth, face, nose, arms, head, ears, legs. I have to keep my body clean even if I can't walk. If it is clean, it will make the body strong. During nighttime, the child will help me a bath. Now, I can do it myself...I feel happy that I can take good self-care. (PaT5)*

### ***3. Having a simple and comfortable life with support***

In everyday life, participants described their living within a simple and self-sufficient way. Having a simple and comfortable life by consuming safe food in the adequate amount, getting enough sleep and being careful of falling, keeping the mind relaxed as the following details:

#### ***3.1 Consuming safe food in the adequate amount***

Most of the Muslim older adults living in their homes focused on eating natural food but in moderation, not overeating. They often used a product from the local market and cooked for themselves rather than buying from outside. They focused on eating fresh food, avoiding sweet, oily, salty foods. The indispensable food is real honey, brown rice and bananas. Their diet had been modified, avoiding eating sweets, salty foods, and focusing on eating Tom Yum fish, fish curry, and vegetables; reducing eating meat. In addition, non-toxic food was concerned by



growing vegetables in the kitchen themselves. Some used herbs and ate dietary supplements such as *Ganoderma lucidum*, *Hemowit*, royal jelly, and focused on drinking enough water.

*...food, we must eat things that are grown by ourselves. I eat vegetables and fruits that are grown in the garden. There are grapefruit, papaya, and chickens to treat themselves. My favorite food is kaeng som (spicy curry), yellow curry. Sometimes I have my kids make curry with kua yuk chicken at home, by bringing their own chickens to make curry. After eating, I must eat one banana every time (can eat all types of bananas) (Ma1)*

*...food that makes me strong...the food that I like to eat is Tom Yum Snapper. I can eat every day, never get bored. Wife makes sea bass tom yum. The fish is marine fish, not the farmed fish to cook, it must be fresh, 2-3 pieces per plate. Sometimes relatives bring the fresh fish from the lake, Songkhla, to make tom yum, add tomatoes, spring onions and coriander, add fresh lime, a little chili paste so it's not spicy....don't like spicy food. Cooking still have some good flavor, do not eat salty, but eating plain flavor following a doctor's advice...When I woke up and grabbed water and drank 2 glasses of water every day because the body is dehydrated during the night, the blood becomes viscous, it is inconvenient to feed the brain. Therefore, drinking water helps for better blood circulation. This idea came from a conversation with a friend a long time ago, who is a Chinese doctor, said that we had to drink 6 glasses of water after waking up, saying that one glass goes to the heart, another glass to the liver and the rest to the body and other organs, so use it. (Pa4)*

*Not only eating non-toxic food but also avoiding eating too much dinner under the beliefs that it would harm the digestive system as a participant who had a gastritis, she reflected "during dinner time try not to eat a lot, because if excreting heavy I do not want to call my children to come and help me at night. I don't want them to drive during the night. (Ma1)*

In addition, more organic foods and vegetables which they can find around the house and grow for their own farms were consumed, as a participant said:

*...I eat vegetables that I grow myself in the backyard. There are eggplants, long beans, morning glory, Chinese cabbage, and chili (PaT6)*

### *3.2 Getting enough sleep and be careful of falling*

Moreover, participants described the important practice in daily life by getting enough sleep and be careful of falling when staying home alone. Although Muslims participants had a lot of time to sleep at home, none of the participants reported difficult sleeping. They always find a way to manage to get enough sleep by looking for some activities to do in helping their sleep, as a participant said.

*"I will not sleep too much. If I can't sleep, I will recite a prayer for blessings. (Ma3)*

In addition, some participants had emphasized on carelessness and the need to be careful of falling due to muscle weakness as well as having an enough sleep. As a participant who had a recovered from paralysis reflected on sleeping is important as being safe when mobilized at home.

*...when I get better, I can start walking on my own. I do not underestimate my risk wherever I walk, keep in mind, don't forget to walk with assistance or use four-legged cane, don't think to walk on your own. I also sleep well and get the nap in the afternoon because feeling sleepy can increase my own risk of fall. I always reminded myself that don't underestimate the risk of falling due to muscle weakness.... be careful not to fall again It's going to be difficult... I used to go to the bathroom, couldn't lift a cane on four legs, almost fell down, couldn't move forward. Since that day, I have been very careful. (Pa4)*

### *3.3 Keeping relax the mind with a simple life that does not make a burden to children*

Some informants tried to live a simple life so as not to cause trouble for children especially those whose children have low income and must go out to work. They would not be offended, not complain, and stay for the children. One said that living a simple life would not increase the burden on the children.

*...I tried to live easily, eat easily. I have breakfast to eat, my children bought only 10 baht of Khanom Krok (local dessert) and it's okay. One banana is enough to eat, a piece of fruit, will not think much, never insulted children. I live for the children with love and cherish, think of Allah only to relax the mind. (Ma5)*

*...not stressed, not thinking too much, not thinking what it has to be just go to the bathroom by yourself, you're comfortable and satisfied. When you have a stroke, you don't have to stress anymore. I know that stress affects our mind and body. After that, it's not stressful anymore. If there is stress, don't think, try to leave it as much. (MaT1)*

All participants try to find simple stress-relieving activities for themselves in order to find happiness in doing what they enjoy, relieving emotions and stress, such as watching the favorite TV program, talking to your children and neighbors, making product from crops near the house for sale as some participants reflected.

*...In the morning, I will take a seat in front of the house. When someone I know walk or ride passes by my house, I would greet him or her. If he or she is not in a hurry, he'll stop by and talk for a while (for a moment). I like talking to other people, it is enough to relieve loneliness (MaT1)*

A field not from observation also reflect her daily life. Ma (she call herself) often comes out and sit in front of the house. Later, she watches the favorite TV shows, likes to watch the news, watch advertisements for alternative medicines, dietary supplements. Watching TV has some benefits as she described “it can help relieve boredom, it's like having a TV as a friend”.

*...Men would go out and sit in the sun in front of the house every morning. Get around the sun outside...help me Bye Jai (Thai word means happy). It is more comfort in the sun, and feel warm rather than cold inside. When sitting in the sun in front of the house, neighbors drove past and greeted each other. If some riding a bicycle, they would stop and talk for a while when they pass by my house and then go. This just makes me feel comfortable. (PaT 3)*

*...When feeling bored or in the evening, I will talk to my children and grandchildren, sometimes I go to the end of the three-way intersection. to greet the neighbors. Getting out of the house to talk to other people sometimes helps a lot to relieve loneliness. (PaT 5)*

Moreover, preparing for his/her death was also thought in advance. Some people have planned for the possessions by managing the assets available to the children. They felt no worry because it brings about unity among relatives as clear information. Some participants who had a well plan for their inheritance, they felt relax as it would not cause a burden to their children in the future. As one participant said:

*...The properties are divided equally and prepared for each child. That stuff has already been set up... Let's allocate the land to every child, 3-4 rai (acre) each, so that each child won't have a debate, I have no stress, no worries. (MaT2)*

#### **4) Managing symptoms to gain a balance and restore health**

The participants in this study usually enjoyed staying at home. Some who had symptoms related to chronic illness such as pain at the knee, backache, and pain with numbness from his or her illness, they often used self-management to relieve symptoms with natural methods and some alternative health care methods, such as taking wind pills, using herbal oils to relax the muscles, herbal bath and using holy water, etc., as detailed below.

*...I usually drink honey every day, about 1 bottle/month to help my appetite. By dissolving warm water with 2 tablespoons of honey, eat before breakfast - evening. and take some pills to reduce wind when having symptoms of nausea, it helps the circulation of blood and wind... eats real honey obtained from the island. (Ma3)*

*...I sometimes used massage, herbal bath and use holy water to rub the area that is painful or weak It is also a method that is widely used in Muslim elders and has been recognized that it works well... In the past, my grandmother used to be a masseuse to treat people (this is an additional occupation learnt from a great-grandfather). I used when I had aches and pains, tends to massage by myself by touching, squeezing, and ironing. Aches and pains will use massage oil when having sprains.*

Moreover, most participants could take care of their health when their illness was not severe or can control by using alternative medicine to treat the illness and restore health, such as using herbal medicines, relaxation massage, or seeking a local healer especially for neuromuscular symptoms, such as some older adults with limb weakness from ischemic stroke. Most of the rehabilitation was done by chiropractors in the community so that the muscles would

return to strong normal use. Participants also shared time and information about the methods to heal and restore the body by folk healers in self-help group in doing daily activities.

*...I had a stroke and weak limbs. In the past, there was a masseuse in the village who came to give massages with assistance from a daughter. I then able to eat and go to the bathroom by myself. Now my daughter is urged to rejuvenate herself every day, by squeezing my arms and legs by herself. My legs had more strength (MaT1)*

Older adults also had the ability to treat their own minor ailments by using first aid and using herbs that can be found in the community. They reflected their perspectives partly because it is easy-to-get access, as an informant reflected:

*...We use Plai oil that Yo (Khao) made to treat toothache, gum pain, swollen gums. Why is it not difficult to make it? Use about 1 kg of Plai head, cloves, borneol, camphor, menthol, and keratin oil. When we have pain, we apply it on the gums. it will not be inflamed Cloves will help healing teeth. Put some oil on your hands and rub it inside when applied, it becomes hot, feels numb, has no sensation. If we put the medicine in about half an hour, it will be better. When we have some symptoms such as abdominal pain, flatulence, body aches, we put oil on it and will get better soon. (PaT4)*

In addition, most participants sought ways to keep the mood clear when they spent time at home particularly during the Covid-19 pandemic. The activities such as relaxing the mind by praying, finding a stress-relieving activity, watching a favorite TV program, meeting neighbors on occasion

*...During this period, you can only watch television to relieve stress. Love to watch boxing shows. In the past, when there was no covid (Covid-19), I went to sit and talk with friends*

*who used to work in Malay and able to speak Multilanguage, both Malay, Chinese and Thai. It helps to encourage, relax, and reduce boredom. (PaT2)*

*...This year there is an epidemic. I don't go to the doctor. My children said it was difficult to go to the doctor. I have diabetes and heart disease. If I don't take medicine. I cannot walk a lot due to tired. I must sit and rest...God has given us what we are going to be, we must accept it and pray and pray. God bless you so you can live and help yourself. I pray 5 times and prays to God for good health. When I still see it, I read the Qur'an and donate some stuff on an occasional basis because we believe giving to others will make merit and create good health for ourself. ... During this time, my daughter said that it's inconvenient and unsafe to take me to the hospital. She concerned of old age group who could take more risk when go out. (Ma5)*

## **Discussion**

From the study findings, it showed that participants in the urban and rural districts, described their health practice in similar way based on their cultural beliefs for the maintenance of holistic health although they were living in different areas. Religious beliefs related to health are regarded as part of life which guide their health practice. The living conditions, way of life, and culture were integrated according to the traditional and traditional Islamic principles (Lillahkul and Supanakul, 2020). Some participants preferred to treat their ailments using alternative medicine such as massage, using oil to massage the painful area, eating herbs to prevent disease including a selection of dietary supplements under the belief that it helps to improve health more than using the modern medicine treatment alone.

Their daily health practices under the meaning of “life and health is designated by Allah for living with a nature and comforts with their age” consisted of four patterns: 1) maintaining day to day functioning to stay independence, 2) having a simple and comfortable life with support, 3) performing religious activities as a priority of life for well-being, and 4) managing symptoms to gain a balance and restore health.

Muslim older adults in this study spent most of the time at home to take care of their body, mind, and spirit integrated between the principles of Islam and the principles of medicine and public health. The participants had a belief and lifestyle that strictly adheres to the religious principles and teachings of the Quran. Holistic health care and health-promoting behaviors of Muslim elders are therefore continually practiced together until they become part of their daily lives. Maintaining day to day functioning to stay independence is important and link with having a simple and comfort life with support.

In addition, the participants described personal relationships with friends or neighbors, to maintaining their social networks as important. Although they were at home, the social context and interactions with others influenced their lives positively and were described as essential. It means being able to see and interact with family members and friends daily. Their social context of being central to this view is shared with Garbaccio et al. (2018) who found that active older adults generally enjoy their life of one able to stay in the private home, enjoying relationships with family and friends as part of life.

Older adults need to take care of their physical, mental, social, and spiritual health in order to stay healthy and happy, by eating natural food, emphasis on organic food or local vegetables are available locally and seasonally. Most of them eat freshly cooked food, and soft food, and eat in moderation. They addressed the important issues which can promote healthy and



strength by not overeating, avoid sugary, oily, salty foods and eat foods that are appropriate for the disease or foods on farms. This practice accords with the commandments set forth in the Quran regarding the care of food consumed, such as eating enough food (not too much) to sustain life and strengthen the body, eating nutritious, clean, chemical-free food, appropriate for each person's age and condition. The older adults should choose foods that are easily digestible because the stomach, intestines, liver, and kidneys begin to deteriorate, and should avoid very salty foods. (Sutheravut and Nima, 2009). In addition, older people take herbs and immune-boosting supplements to prevent disease by nourishing health according to beliefs such as bitter gourd, honey, Lingzhi, and dietary supplements according to health problems such as royal jelly, poppy oil, and rice germ oil. The participants also preferred to eat herbs and supplements because they believed that it helps to maintain a physical health. This is consistent with the study by Panyathorn and Tajaeng (2020) who addressed the attitudes towards products and consumption of dietary supplements among the older adults in a district, Udon Thani Province that supplements can give strength in life. The Quran also mentions honey as a cure for human ailments because honey contains nectar from all fruits (Lillahkul and Suphankul, 2020).

The overall theme derived from the four categories suggested that Muslim HOAs in our study addressed health practice to pursue an independent life, maintain their health with the purpose of God and a given tradition and value relationships and social networks for as long as possible. Although they did realize that life would change, it was thought as nature and return to the God under the religious concept. The ability to lead a full and balanced life close to family, friends, nature and activities were important for their feelings of good health and well-being. Therefore, Muslim older adults living at home performed self-care through natural methods which similar to a previous study (Sombat, Phiriyapong, & Atnarong, 2003). Consuming

foods, supplements and alternative medicine to gain a balance and restore health are considered as important for healthy older adults. Most participants in this study use herbal massage oil to rub and massage the sprained area by themselves. Some older adults tend to relax with massage oils at home. Similar pattern of drinking holy water has also found from the study of local wisdom on self-care of Muslims in the southern border provinces. Healing with holy water is a very popular treatment among the local people particularly from Muslim folk healers. It is also a drug listed in the Quran and Hadith. However, it is a treatment in conjunction with modern medicine (Hemman, 2016). In addition, Muslims HOAs described their exercises for health according to religious principles, such as swinging arms, swinging legs, and walking regularly to help keep the body healthy. This may be similar to the study by Lillahkul and Supanakul (2018) who explored about activities among Thai Muslim older adults performed for health promotion. In this study, most Muslim older adults stay at home tend to do some exercise with low intensity that does not focus on strengthening the heart and lungs. This may be due to the presence of physical limitations.

However, the participants living in their homes also have good hygiene habits, by using clean water to wash their four organs, namely the face, both hands and arms, head and feet before praying and linked it with the belief that those who are physically unclean will not be able to practice certain religions. Cleanliness is also a part of the faith and link with their mental health. They were taught as an inner purification by clinging to one Allah and having faith will make the soul pure, calm, and energetic, ready to worship God. As a result, those who maintain cleanliness will have good health (Sutheravut & Nima, 2009). In addition, some older adults drink warm water and drink enough water to meet the physical needs. It is believed that blood circulation is good and excretion is convenient.

In mental health care, Muslims HOAs try to keep their minds refreshed and relaxed by doing stress-relieving activities such as watching television, and greeting their children and grandchildren. Meeting the neighbors on occasion would help them to enjoy life based on the principle of satisfaction in the present life, which is the practice of religious doctrine in enhancing mental health that satisfaction with Allah's destiny is extremely important to good mental health. A person's dissatisfaction with his or her condition would make him or her unhappy partly because they do not appreciate what Allah said. In contrast, all participants in this study believe that Allah had given to most of the Muslim older adults living in their homes to have good human relations and have smiling faces with their children and neighbors. They spent a simple life by not acting as a burden to their children. In addition, they were taught to consider the charity given to others and keep their own mental health care according to the teachings of religion. So, the Muslim older adults stayed at home and relied on their religion as a mental relief. The older adults prayed regularly, read and listened to the Quran on a regular basis. Prayer is one of the religious practices that helps to relax the mind, and have peace of mind from purifying the mind without any sin or any bad thing. As Allah says, "The reward of performing Five prayers and Friday prayers is that Allah will atone (cleanse)" and "even prayer suppresses badness and evil" (Sutheravut & Nima, 2009)

Greeting their neighbors on occasions and participating in religious activities at the mosque were usually performed in everyday life. Although some older adults could not walk and suffered from chronic diseases, they would greet their neighbors from a distance. Having children take them to religious activities and attend activities at the mosque when needed. Most of the Muslim older adults families who live in their homes are extended families surrounding with their relatives and children. This allows the older adults to receive close relationships from

family members in meeting neighbors and participating in religious activities. Similar to a study by Thongket et al (2019) who focused on the health behaviors of the Muslims older adults in Satun province, they received support and assistance from their family, relatives, and neighbors whenever they need it.

Spiritual health care was found for all HOAs in this study who always performed strictly according to their religious practice. They have brought the religious doctrines set forth in the Quran as a way of life and took care of their spiritual health for a variety of vitality and spiritual health as peaceful life. Older adults prayed 5 times a day by sitting or lying down for those who cannot stand. Say Zikrullah (Remembrance of Allah) After the praying, asked du'ar (blessings) to Allah for good health, as well as read the Quran daily to accept the laws of God's circumstances. Some elders read the Azkar Annabawi, morning and evening prayers of the Prophet) Surah Surah Qulhu Allah (Doctrine), listened to religious teachings from Baba (Tho Chru) on YouTube and read Muslim religious practices regularly, and donated money for merit-making in some occasions. Have faith in Allah is regarded as the heart of being a Muslim (Sutheravut & Nima, 2009). They all believe that believing in Allah would bring them good health, happiness, and peace of mind. Having strong willpower in the face of illness gained a lot of merit in this world and the next world. This is in line with the study by Lillahkul and Suphankul (2020). Therefore, HOAs Muslims have spiritual health care at all stages of their lives under the faith of religion until the end of life.

Some Muslim older adults who lived in their homes had a plan to manage their wealth for their children to feel relief of anxiety when leaving with the allocation of the existing assets to the children equally. It was a preparation in advance for a peaceful departure without commitment in which the division of inheritance is organized according to religious principles

confirmed by the Quranic text (Mhadman, 2014). Therefore, the older adults have an inheritance plan to prevent conflicts among their children. It's faith in the Quran which bring them to be happy both physically and mentally and goals in the next world and to be freed from punishment and to enter paradise happily ever after (Onkhom, 2009).

Family support has also facilitated the health of the HOAs living at home. Most of the older adults have children as their primary caregivers. The children keep rotating to take care of various aspects such as doing daily activities. Having disease-appropriate diet, movement, excretion, religious and ritual activities at the mosque, doctor visits, appointments, etc. Islam has urged all Muslims to do good, showing gratitude to their parents is a noble and important duty. This is a duty that is obligatory (compulsory) for children to take care of their parents to the best of their ability, otherwise, it will be considered ungrateful (Tohma et al., 2018). It is consistent with a study by Aree (2013), who found that the children of older Muslims in urban communities have religious beliefs and beliefs about having the older adults live with them is a very good thing. The longer the older adults have, the more it is considered good and auspicious. Because of that longevity is to have the opportunity to do good deeds more than others. In addition, Muslim older adults who received regular family care feel worthwhile have a warm heart and have the power to continue living. Getting visits and encouragement from neighbors is associated with the health of the Muslim older adults living at home (Phithakkumpol, 2004). Older adults feel comfortable talking to health workers when they visit their homes. This is consistent with a previous study (Detthippornpong, Songwathana, Bourbonnais, 2022) which found that village health volunteers in the community are often related to the people in the community in the form of kinship. This makes it easy to reach people in the community. In the community, VHV supports the work of nurses in following home visits, taking care of the comfort, teaching

relatives about basic care and reporting any abnormal symptoms to the nurse, providing psychological support, and encouragement, and coordinating with government organizations to provide financial and social support to patients. It is also consistent with the study by Kuhirunyaratn et al. (2018), which examined health promotion behaviors of older adults living in suburban communities in Khon Kaen Province. All received a pension from the government, some people received a disability allowance. Most of them received money that their children provided for personal expenses and the money would be used to spend on health care and making merit for peace of mind.

In this study, all the Muslim older adults living in their homes had God as their spiritual dependency. This is the most important factor that promotes good health for the older adults. Older adults have a strong commitment and faith in God, and are strict in performing religious duties. Doing Du'a' (blessings) to keep themselves healthy always, make them feel warm have peace of mind with good mood control and accept everything that happens to them because it is the will of God. This is consistent with the previous study (Thiprat, 2008), who found that the older adults had a high level of religious beliefs related to health. When they were stressed or worried about illness, they prayed to help and calm their minds. Therefore, it can be concluded that the Muslim HOAs have faith in Allah, and viewed their health in a holistic way, as "Life and health are designated by Allah for living with nature and comfort with their age" Allah was always in their heart of health practice.

## **Limitations**

Some limitations could be 1) the participants were recruited from small areas which may influence transferability to apply the findings in other areas, 2) Covid-19 pandemic may have influenced the relationship between researchers and the informants and could have influenced their response during data collection.

## **Conclusions and implications**

The study provides a holistic picture of traditional health practice in a natural setting in Thai Muslim community which describes the daily life activities of how homebound older maintain their health at home. The health practice of HOAs in this study are strongly related to religion and nature. The knowledge of health practice which they viewed in four domains could be included in the education of nurses and primary health care workers to cultivate the cultural awareness and provide better older adults care at home or in long term care. Interventions need to consider cultural sensitivity and family value of older people to strengthen and promote holistic health among Thai Muslim HOAs.

## **Acknowledgement**

The authors wish to thank the informants for their time and contributions throughout this research. Acknowledgements also go to all health center and community leaders who kindly assisted us to approach participants. We also thank the Thailand National Research Institute for funding support.

## References

- Aree, W. (2013). Self-care of Muslim older adults: *Evidence from one community in Bangkok metropolitan (Master's thesis)*. Mahidol University, Bangkok.  
[https://doi.nrct.go.th/ListDoi/listDetail?Resolve\\_DOI=10.14457/MU.the.2013.329](https://doi.nrct.go.th/ListDoi/listDetail?Resolve_DOI=10.14457/MU.the.2013.329)
- Alabdulwahab, S. S., Kachanathu, S. J., & Oluseye, K. (2013). Physical activity associated with prayer regimes improves standing dynamic balance of healthy people. *Journal of Physical Therapy Science*, 25(12), 1565–1568. <https://doi.org/10.1589/jpts.25.1565>
- Arihathairat, S. (2020). *Promoting Nursing Practice based on Islamic Ways Experienced by Head Nurses of Community Hospitals in Three Southernmost Province, Thailand* [Master Minor Thesis, Prince of Songkla University]. <https://kb.psu.ac.th/psukb/bitstream/2016/13041/1/Subaidah%20Arihathairat.pdf>
- Chansangrat, N., Phibal, A., Niltaya, T., Thiratchakul, J., & Hatthakit, U (2014). *The need of spiritual health promotion of the Muslim older adults in Narathiwat Province*. Prince of Naradhiwas University.
- Detthippornpong, S., Songwathana, P., & Bourbonnais, A. (2022). Holistic Health Practices of Rural Thai Homebound Older Adults: A Focused Ethnographic Study. *Journal of Transcultural Nursing*, 33(4), 521–528. <https://doi.org/10.1177/10436596221090270>
- Dossey, B. M. & Keegan. (2016). *Holistic Nursing A Handbook for Practice* (7th Ed.). United Stage of America.
- Foundation of Thai Gerontology Research and Development Institute: TGRI. (2019). *Situation of the Thai older adults 2017*. Institute for Population and Social Research Mahidol University.
- Garbaccio, J. L., Tonaco, L., Estêvão, W. G., & Barcelos, B. J. (2018). Aging and quality of life of older adults in rural areas. *Revista brasileira de enfermagem*, 71 Suppl 2, 724–732.  
<https://doi.org/10.1590/0034-7167-2017-0149>
- Hemman, M. (2016). *Muslim caregivers' experiences in caring for patients with end stage renal disease receiving palliative care (Master's thesis)*. Prince of Songkla University, Songkla.



- Husebo, A. M., & Storm, M. (2014). Virtual visits in home health care for older adults. *The Scientific World Journal*, 2014, 689873. <https://doi.org/10.1155/2014/689873>
- Kuhirunyaratn, P., Ratanasiri, A., Jindawong, B., Natiboot, P., Watchanapan, L., Junthakhun, C., & Supapinij, C. (2018). Health promotion behaviors of older adults living in an urban community of Khon Kaen province. *Srinagarind Medical Journal*, 33(2), 153-160. <https://li01.tci-thaijo.org/index.php/SRIMEDJ/article/view/115767>
- Leininger, M.M. & McFarland, M.R. (2002) *Transcultural nursing: Concepts, theories, research and practice*. 3rd Edition, New York: Mc Graw-Hill Medical Publishing Division.
- Leininger, M.M. & McFarland, M.R. (2006). *Culture care diversity and universality: a worldwide nursing theory*. 2<sup>nd</sup> Edition, Sudbury, MA: Jones and Bartlett.
- Lincoln, Y.S., & Guba, E.G. (1985). *Naturalistic Inquiry*. Sage.
- Lillahkul, N., & Supanakul, P. (2020). Homebound older adults health promoting behaviors and related factors in a Muslim cultural context: A case study in Yala province. *The Southern College Network Journal of Nursing and Public Health*, 7(2), 194-205. <https://he01.tci-thaijo.org/index.php/scnet/article/view/177774>
- Lohwithee, W. (2020). The strengthening of well-being in accordance with guidelines suggested by the messenger of Islam. *Journal of Liberal Arts, Rangsit University*, 15(2). 65-76.
- Mhadman, M. (2014). *The heritage management in accordance in the Islamic law of Muslims in Satun province* (Master's thesis). Prince of Songkla University, Songkla.
- Mohmula, M. (2007). Chronic care in the crisis of unrest situation in three southern Thai borders. In *health system management during the crisis in multicultural society*. (pp. 41-54). TIMA: Thailand Islamic Medical Association, Southern Health Systems Research Institute Prince of Songkla University, Songkhla and National Health Security Office, Songkhla. [https://k4ds.psu.ac.th/k4ds\\_search/pdf/projects/TC0007.pdf](https://k4ds.psu.ac.th/k4ds_search/pdf/projects/TC0007.pdf)
- Ministry of Public Health. (2016). *20-year National Strategic Plan* (Public Health section). <http://www.oic.go.th/FILEWEB/CABINFOCENTER2/DRAWER023/GENERAL/DATA0000/00000077.PDF>.

- National Health Security Office: NHSO. (2016). *Manual of supporting administration in long term care services for public health in caring for dependent elders*.  
[https://www.nhso.go.th/storage/files/shares/PDF/fund\\_ltc13.pdf](https://www.nhso.go.th/storage/files/shares/PDF/fund_ltc13.pdf)
- Onkhom, S. (2009). *Comparative religion*. (8th ed.). Bangkok: Duangkaew.
- Panyathorn, K., & Beengmum, N (2019). Medication use behaviors in the older adults patient with chronic disease at Chomsri village, Udonthani province. *Udonthani Hospital Medical Journal*. 27(2), 119-127.  
<https://he02-old.tci-thaijo.org/index.php/udhhosmj/article/view/213438/148454>
- Panyathorn, K., & Thajang, S. (2020). Opinions and Consumption Behaviors of Supplementary food products Among the older adults in Nongbu village, Sam Phrao Sub-district, UdonThani Province. *Udonthani Hospital Medical Journal*, 28(2), 121-129.
- Phithakkumpol, A. (2004). *Life, sickness and death. Lecture documents Caring for terminally ill patients according to Islamic guidelines*, 7-8 May 2004. Songkhla: Faculty of Medicine. Prince of Songkla University.
- Rojanadhamkul, N. (2020). *Nursing care for older adults: physiology and mental health aspect*. Mission Faculty of Nursing, Asia-Pacific International University.
- Sombat, K., Phiriyapong, N., & Atnarong, S. (2003). *Local wisdom in self-care of Muslims in the southern border provinces: research report*. Office of the national research council of Thailand, Bangkok.
- Suthravut, P., & Nima, Y. (2009). *Integration of Islamic law knowledge and health promotion: Chronic diseases (Cardiovascular Diseases, Diabetes, Hypertension)* (1 st ed.) Southern Health Systems Research Institute Prince of Songkla University, Songkhla.
- Thiprat, S. (2008). *Selected factors related to health promotion behaviors of Thai-Muslim older adult members of older adults' clubs in upper Southern region (Master's thesis)*. Chulalongkorn University, Bangkok.  
[https://doi.nrct.go.th/ListDoi/listDetail?Resolve\\_Doi=10.14457/CU.the.2008.1010](https://doi.nrct.go.th/ListDoi/listDetail?Resolve_Doi=10.14457/CU.the.2008.1010)
- Thongket, J., Saradee, M., Cheuchom, J., Saetae, S. (2019). *Older adult's health behavior in era subdistrict, Tapae district Satool province*. In Research and Development Office, Prince

of Songkla University (Editors) Proceedings in the 10<sup>th</sup> National and International Conference (pp. 1662-1663), at Hatyai University.

Tohma, H., Al-fariti, A., Masarakama, N., Masoh, A., Shawo, L., Radeang, R., Awaeni, P., & Saeyiu, L. (2019). *The mental health care, according to the Islamic way of terminally ill patients: Case study Muslims Thailand patients in the three southern border provinces*. Research report. Health Systems Research Institute, Nonthaburi. <https://kb.hsri.or.th/dspace/handle/11228/5053>.

United Nations: Department of Economic and Social Affairs, Population Division (2020). *World Population Ageing 2020 Highlights: Living arrangements of older persons* (ST/ESA/SER.A/451). [https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/files/undesapd-2020\\_world\\_population\\_ageing\\_highlights.pdf](https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/files/undesapd-2020_world_population_ageing_highlights.pdf)

Vongmonta, A. (2017). Doctrines in the Quran consistent with folk wisdom in health care. Thai Muslim mothers and children in Pattani Province. *Rusamilae Journal*, 38(2), 95–99. <https://so05.tci-thaijo.org/index.php/rusamelae/article/view/95424>

World Health Organization. (2017). *Global Aging 2017*. [www.seniornet.org/index.php?option=com\\_content &view=article](http://www.seniornet.org/index.php?option=com_content&view=article).

Table 1 Characteristics of the informants

ID	Sex	Age	Marital status	Education level	Occupation	Past illness (year)	ADL score	No of child	Caregiver
Ma 1	F	94	widow	Grade 2	Trade market	-Gastric ulcer with HT (1)	11	11	Son/ Daughter/ Nephew
Pa 2	M	84	married	Grade 4	Rubber cutting/ gardening	-DM & HT (10) -Stroke (3)	11	4	Daughter/Son
Ma 3	F	86	widow	Grade 2	Trade market	-Kyphosis and chronic anemia	5	5	Daughter/Son/ Nephew
Pa 4	M	73	married	Grade 4	tailor	-Stroke (4) and HT	11	2	Wife/ Daughter/Son
Ma 5	F	84	widow	none	selling fish in the market	- DM & HT (5), MI (1)	11	13	Daughter/ Nephew
Ma 6	F	89	widow	Grade 4	Farming/	- DM & HT (4)	8	8	Son/ Daughter
Ma 7	F	93	widow	none	trade market	- no history of illness	10	4	Daughter/ Nephew
Ma 8	F	67	widow	Grade 4	construction manager	- DM & ischemic stroke (1)	11	4	Daughter
Pa 9	M	76	married	Grade 4	General contracting, cattle raising	- Hemiparalysis & HT (10) - BPH (5) and CA colon (1)	11	5	Wife
Ma-T1	F	68	married	Grade 4	Rubber farmer	- HT (13) - Ischemic stroke (8)	11	5	Daughter
Ma-T2	F	82	widow	Grade 4	Rubber farmer	- HT (29) - Herniated disc disease (12)	11	8	Daughter /Son
Pa-T3	M	60	married	Grade 7	Rubber agent	- DM (10), chronic kidney disease & HT (4 months)	11	6	Wife/ Daughter
Pa-T4	M	67	married	Grade 4	Rubber farmer	- Gastritis 15 years & Ischemic stroke & Lipidema 1 year - Paraplegia 19 years	5	6	Wife
Pa-T5	M	76	married	None	Rubber farmer	- DM & Lipidemia 16 years Post amputation right leg 3 years	11	7	Wife/ Daughter
Ma-T6	F	85	widow	Grade 4	Rubber farmer	- HT & Lipedema (4), -Ischemic stroke (1)	11	3	Daughter / Nephew

