

1.Jurnal Plus One

Q1 : SJR 0,84

United States

Review Thank You



Thank you for reviewing Manuscript Number PONE-D-21-33982.

[Return to Main Menu](#)

[Log out of Editorial Manager](#)

← Completed Reviewer Assignments

Page: 1 of 1 (1 total assignments)

Action 	My Reviewer Number 	Manuscript Number 	Article Type 	Article Title 	Current Status 	Final Disposition 	Date Reviewer Invited 	Date Reviewer Agreed 	Date Review Due 	Date Review Submitted 
Action Links	1	PONE-D-21-33982	Research Article	Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.	Under Review		Apr 25 2022 7:11AM	Apr 25 2022 8:31PM	May 9 2022 11:59PM	May 2022 8:19

Page: 1 of 1 (1 total assignments)

ments

Results per page 10

Manuscript Number	Article Type	Article Title	Current Status	Final Disposition	Date Reviewer Invited	Date Reviewer Agreed	Date Review Due	Date Review Submitted	Days Taken	Editor's Name	Corr. Author
PONE-D-21-33982	Research Article	Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.	Under Review		Apr 25 2022 7:11AM	Apr 25 2022 8:31PM	May 9 2022 11:59PM	May 6 2022 8:19AM	11	R&R PLOS	Addis Eyeberu

Results per page 10

Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.

--Manuscript Draft--

Manuscript Number:	PONE-D-21-33982
Article Type:	Research Article
Full Title:	Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.
Short Title:	Utilization of Pharmacological Labour Analgesia
Corresponding Author:	Addis Eyeberu Haramaya University College of Health Sciences: Haramaya University College of Health and Medical Sciences harar, ETHIOPIA
Keywords:	Utilization; Analgesia; labor pain; pharmacological labor pain management; Harar
Abstract:	<p>Background</p> <p>Despite a crucial role played by Obstetric care providers (OCPs) in providing pain relief methods during labor, the evidence suggests that there is an ample number of women in Ethiopia failed to have access to pain relief methods. Moreover, to the best knowledge of the investigators concerned, evidence is scarce regarding the utilization of pharmacological labor pain management methods in the study area. Thus, this study was carried out to determine the practice of pharmacological labor pain management methods among obstetric caregivers in Harari regional state health facilities, Ethiopia.</p> <p>Method</p> <p>A facility-based cross-sectional study was conducted from May 20 to June 10, 2021, in Harari regional state health facilities, Ethiopia. All obstetric caregivers in Harari regional state health facilities were included in the study. Data were collected using a structured questionnaire adapted from the previous studies. The data were entered into Epi-data version 3.1 statistical software. Statistical analysis was carried out by using SPSS for windows version 22. Bivariate and multivariate analyses were employed to determine the association between independent variables and the outcome variable.</p> <p>Result</p> <p>The overall utilization of pharmacological labor pain relief methods was 50.9% [(95% CI (45.8,55.8)]. The main barriers for laboring mothers receiving pain relief include high patient flow (54.4%), the limited number of staff (51.9%), and lack of knowledge and skill (32.9%). Being female [AOR=2.07, 95% CI (1.31,3.26)], not having training [AOR=1.60, 95%CI (1.02,2.53)], having unfavorable attitude [AOR=9.1, 95%CI: (5.34,15.35)] were the predictors of utilization of pharmacological labor pain management methods.</p> <p>Conclusion</p> <p>The study revealed that the overall prevalence of utilization of pharmacological labor pain management methods was low. In general, this study identified that variables like sex, training, and attitude of care providers were factors associated with utilization of pharmacological labor pain management methods. The health facilities should give the training to enhance the skill of obstetric care providers and equip themselves with sufficient analgesics drugs.</p>
Order of Authors:	<p>Addis Eyeberu</p> <p>Tamirat Getachew</p> <p>Adera Debela</p>

	Bikila Balis
	Bajirond Eshetu
	Sinetibet Mesfin
	Habtam Bekele
	Dawit Tamiru
	Getahun Tiruye
	Meron Degefa
	Addisu Alemu
	Merga Dheressa
	Yadeta Dessie
Additional Information:	
Question	Response
<p>Financial Disclosure</p> <p>Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS ONE for specific examples.</p> <p>This statement is required for submission and will appear in the published article if the submission is accepted. Please make sure it is accurate.</p> <p>Unfunded studies Enter: <i>The author(s) received no specific funding for this work.</i></p> <p>Funded studies Enter a statement with the following details:</p> <ul style="list-style-type: none"> • Initials of the authors who received each award • Grant numbers awarded to each author • The full name of each funder • URL of each funder website • Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript? • NO - Include this sentence at the end of your statement: <i>The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</i> • YES - Specify the role(s) played. 	<p>The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</p>

<p><i>* typeset</i></p> <p>Competing Interests</p> <p>Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any competing interests that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests.</p> <p>This statement is required for submission and will appear in the published article if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.</p> <p>View published research articles from PLOS ONE for specific examples.</p>	<p>The authors have declared that no competing interests exist.</p>
<p>NO authors have competing interests</p> <p>Enter: <i>The authors have declared that no competing interests exist.</i></p> <p>Authors with competing interests</p> <p>Enter competing interest details beginning with this statement:</p> <p><i>I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]</i></p>	
<p><i>* typeset</i></p> <p>Ethics Statement</p> <p>Enter an ethics statement for this submission. This statement is required if the study involved:</p> <ul style="list-style-type: none"> • Human participants • Human specimens or tissue 	<p>Ethical clearance was taken from the Institutional Health Research Ethics Review Committee (IHRERC) of the College of Health and Medical Sciences of Haramaya University. In addition, permission to proceed was obtained from all public hospitals. And also, informed voluntary written and signed consent was obtained from each participant.</p>

- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below.

Consult the [submission guidelines](#) for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

Format for specific study types

Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare exceptions to address legal and ethical concerns. See the [PLOS Data Policy](#) and [FAQ](#) for detailed information.

Yes - all data are fully available without restriction

A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and **will be published in the article**, if accepted.

Important: Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.

- If the data are **held or will be held in a public repository**, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: *All XXX files are available from the XXX database (accession number(s) XXX, XXX).*
- If the data are all contained **within the manuscript and/or Supporting Information files**, enter the following:
All relevant data are within the manuscript and its Supporting Information files.
- If neither of these applies but you are able to provide **details of access elsewhere**, with or without limitations, please do so. For example:

Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for access to confidential data.

The data underlying the results presented in the study are available from (include the name of the third party

All relevant data are within the manuscript and its Supporting Information files.

<p><i>and contact information or URL).</i></p> <ul style="list-style-type: none">• This text is appropriate if the data are owned by a third party and authors do not have permission to share the data. <p>* typeset</p>	
Additional data availability information:	

Utilization of Pharmacological Labour Analgesia: A Survey of Obstetric Care Providers in Eastern Ethiopia.

Addis Eyeberu^{1*}, Tamirat Getachew¹, Adera Debela¹, Bikila Balis¹, Bajirond Eshetu¹, Sinetibet Mesfin¹, Habtamu Bekele¹, Dawit Tamiru¹, Getahun Tiruye¹, Meron Degefa¹, Addisu Alemu², Merga Dheressa¹, Yadeta Dessie²

Author's affiliation

1. School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar.
2. School of Public Health, College of Health and Medical Sciences, Haramaya University, Harar.

Authors email address

AE- addiseyeberu@gmail.com , TG- tamirget@gmail.com , AD- aksanadera62@gmail.com , BB- bik.balis2008@gmail.com , BE- bajeshetu@gmail.com , SM- sine3686@gmail.com , HB- habeti.bekele@gmail.com , DT- dawittamru5@gmail.com , GT- gtiruye@gmail.com , MD- merrydegefa1@gmail.com , AA: alemuadisu789@gmail.com , MD- mderesa@yahoo.com , YD: yad_de2005@yahoo.com

Corresponding author information.

- Addis Eyeberu
- addiseyeberu@gmail.com : Institutional email: Addis.Eyeberu@haramaya.edu.et
- Haramaya University College of Health and Medical Science, Harar, Ethiopia
- ORCID: <https://orcid.org/0000-0002-3147-3770>; Institutional link: www.haramaya.edu.et
- P.O. BOX 138 Dire Dawa, Ethiopia

Abstract

Background: Despite a crucial role played by Obstetric care providers (OCPs) in providing pain relief methods during labor, the evidence suggests that there is an ample number of women in Ethiopia failed to have access to pain relief methods. Moreover, to the best knowledge of the investigators concerned, evidence is scarce regarding the utilization of pharmacological labor pain management methods in the study area. Thus, this study was carried out to determine the practice of pharmacological labor pain management methods among obstetric caregivers in Harari regional state health facilities, Ethiopia.

Method: A facility-based cross-sectional study was conducted from May 20 to June 10, 2021, in Harari regional state health facilities, Ethiopia. All obstetric caregivers in Harari regional state health facilities were included in the study. Data were collected using a structured questionnaire adapted from the previous studies. The data were entered into Epi-data version 3.1 statistical software. Statistical analysis was carried out by using SPSS for windows version 22. Bivariate and multivariate analyses were employed to determine the association between independent variables and the outcome variable.

Result: The overall utilization of pharmacological labor pain relief methods was 50.9% [(95% CI (45.8,55.8)]. The main barriers for laboring mothers receiving pain relief include high patient flow (54.4%), the limited number of staff (51.9%), and lack of knowledge and skill (32.9%). Being female [AOR=2.07, 95% CI (1.31,3.26)], not having training [AOR=1.60, 95%CI (1.02,2.53)], having unfavorable attitude [AOR=9.1, 95%CI: (5.34,15.35)] were the predictors of utilization of pharmacological labor pain management methods.

Conclusion: The study revealed that the overall prevalence of utilization of pharmacological labor pain management methods was low. In general, this study identified that variables like sex, training, and attitude of care providers were factors associated with utilization of pharmacological labor pain management methods. The health facilities should give the training to enhance the skill of obstetric care providers and equip themselves with sufficient analgesics drugs.

Keywords: Utilization, Analgesia, labor pain, pharmacological labor pain management, Harar

INTRODUCTION

Pain during labor and childbirth is a unique and the most severe pain that a woman ever has to face. The extent to which a woman feels in control of pain during labor is an indicator of maternal emotive wellbeing in childbirth [1]. Pharmacological labor pain management methods include oral tablets (paracetamol, codeine, or tramadol), inhalations, and injectable analgesics [2].

Pain related to childbirth is present during pregnancy, labor, and post-partum periods [3]. The satisfaction of childbirth depends on the care provided to the parturient during labor and delivery [4]. Evidence showed that 85.1% of the patients want their labor pain to be relieved, but 38.9% did receive analgesia during labor [5].

Even if Obstetric care providers (OCPs) have an imperative role in providing pain relief options during labor, the majority of women in Ethiopia have not accessed pain relief methods. The utilization of pharmacologic obstetric analgesia methods was found to be zero [6, 7]. The common factors that affect the utilization of pharmacological labor pain management methods were inadequate knowledge, negative attitude, unavailability of analgesic drugs in the health facilities, individual factor, and institutional factor [6-8].

Mother with intense labor pain affected by emotional distress, depression, and anxiety. Fear and anxiety are significant influences on pain experiences which is one reason why mothers are accompanied by another person during childbirth [9, 10]. Psychosocial factors have been implicated in the pain experienced during childbirth, which can have both short and long-term consequences on the mother's health and her relationship with her infant [3].

Ethiopia develops and implements Ethiopian Emergency Obstetric and Newborn Care (EmONC) assessment standards to enhance appropriate management of labor pain and management of complications during labor and delivery [11]. And also, the Federal Ministry of Health formulates different policy and management protocols regarding managing labor and delivery.

Despite those efforts, the provision of pain relief in labor is neglected and has remained an ongoing challenge [12, 13]. A lack of awareness, unfavorable attitude, and availability of pharmacological methods are considered to be the main reasons why women in Ethiopia do not receive adequate pain relief [6, 12, 14]. Even though healthcare providers in Ethiopia understand that women suffer significant pain during labor, currently they do not provide effective pain relief [15].

Though it is known that effective labor pain management results in greater birth satisfaction for the mother. In many LMIC including Ethiopia, OCPs' awareness of utilization of pharmacological labor pain management methods is low and even zero in some parts of the country. Moreover, to the best knowledge of the investigators concerned, evidence is scarce regarding the utilization of pharmacological labor pain management methods in the study area. So, this study was carried out to determine the utilization of pharmacological labor pain management methods among obstetric caregivers in Harari regional state health facilities, Ethiopia, 2021.

METHOD AND MATERIAL

Study Design, Study Setting, and Period

A facility-based cross-sectional study was conducted from May 20 to June 10, 2021, in Harari regional state health facilities, Ethiopia. The region is found 526 km towards east of Addis Ababa, the capital city of Ethiopia. The region is circled by Oromia regional state. The region is divided into nine woredas with three of them are rural and six are urban. The urban districts are subdivided into 19 kebeles (the lowest administrative division in the country), and the rural districts are subdivided into 17 peasant associations (which is equivalent to kebeles in urban cases). The total population (2021 projection based on the 2007 Census, CSA) of the region is 270,000 from which 136,000 are males and 134,000 are females [16]. There are two public hospitals, two private hospitals and 2 governmental hospitals, eight public health centers, and 20 health posts in the region.

Study Population

All obstetric care providers in Harari regional state health facilities during the study period were the study population. Since there were small numbers of source population in the study area, all obstetric caregivers available during the study period were included. Obstetrics care providers not available during data collection due to illness, annual leave, and left for training were excluded from the study.

Sample Size Determination and Sampling Procedure

The study was conducted at those health facilities having maternal services. These include two public hospitals (HFSUH, and Jugol hospital), one private hospital (Harar General hospital), two private clinics, and nine health centers of the Harari regional state. Since the number of the source

population (OCPs) in all health facilities is too small, all obstetrics caregivers were included in the study. Totally 464 Obstetrics Care Providers (OCPs) were included in the study.

Data Collection Methods and Data Quality Assurance

A structured pre-tested self-administered questionnaire was prepared by adapting from different studies considering the local situation of the study area [7, 14, 17]. The questionnaire had four essential components including providers' socio-demographic characteristics, knowledge, attitude, and utilization-related questions.

The data was collected by research-qualified five professional Nurses and supervised by two MSC holding Midwives. The quality of the data was guaranteed by pretesting using the 5% of the respondents who are not eligible for the study before actual data collection. The completeness of the data was checked daily. Double data entry was done by two data clerks and consistency of entered data is cross-checked by comparing two separately entered data on Epi-Data.

Measurement

Good utilization of pharmacological labor pain management methods: Obstetrics Care Providers (OCPs) who were practicing greater than or equal to the mean value of listed labor pain management methods ($0.65(SD= 0.769)$) [7]. While Obstetrics Care Providers (OCPs) who answered less than the mean values of practice-related labor pain relief method questions labeled as poor utilization.

Obstetrics Care Providers (OCPs): Midwifery, Nurses, Health Officers, and residents.

Adequate knowledge: Obstetrics Care Providers (OCPs) who answered greater than or equal to the mean values of knowledge-related labor pain relief method questions. Those Obstetrics Care Providers (OCPs) who answered ≥ 6.65 from total knowledge-related questions were considered as Obstetrics Care Providers (OCPs) who had adequate knowledge about labor pain relief methods [7].

Inadequate knowledge: Obstetrics Care Providers (OCPs) who answered less than the mean values of knowledge-related labor pain relief method questions. Those Obstetrics Care Providers (OCPs) who answered < 6.649 from total knowledge-related questions were considered as OCGs who had inadequate knowledge about labor pain relief methods [7]

Favorable attitude: Obstetrics Care Providers (OCPs) who answered greater than or equal to the mean value of attitude-related labor pain relief method questions. Those Obstetrics Care Providers (OCPs) who answered ≥ 4.1 from total attitude-related questions were considered as OCPs who had a favorable attitude about labor pain relief methods [6]

Unfavorable attitude: OCPs who answered less than the mean value of attitude-related labor pain relief method questions. Those OCPs who answered < 4.1 from total attitude-related questions were considered as OCPs who had an unfavorable attitude about labor pain relief methods [6]

Statistical Analysis

The data were entered into Epi data statistical software version 3.1 and then exported to SPSS window version 22 for analysis. Descriptive statistical analysis such as simple frequency was used to describe the characteristics of study participants such as socio-demographic, level of knowledge, and attitude of obstetrics care providers. Then the information was presented using frequencies, tables, and figures.

Bivariate analysis and multivariate analysis were done to see the association between each independent variable and outcome variables by using binary logistic regression. The assumptions of binary logistic regression were checked. The goodness of fit was checked by Hosmer-Lemeshow statistic and omnibus tests. All variables with $p < 0.25$ in the Bivariate analysis were included in the final model of multivariate analysis to control all possible confounders. The multi co-linearity test was carried out to see the correlation between independent variables by using the standard error and collinearity statistics (variance inflation factors > 10 and standard error > 2 was considered as suggestive of the existence of multi co-linearity). The direction and strength statistical association was measured by odds ratio with 95% CI. Adjusted odds ratio along with 95% CI was estimated to identify the association between independent variables and utilization of pharmacologic labor pain managements by using multivariate analysis in binary logistic regression. In this study, $P\text{-value} < 0.05$ was considered to declare a result statistically significant

Ethical Considerations

Ethical clearance to conduct this study was obtained from Haramaya University, College of Health and Medical Sciences, Institutional Health Research Ethics Review Committee (HU-IHRERC). Official letters to make a study was also obtained from health facilities as needed. The health

facilities were informed about the purpose of the study and the risk and benefits of the study. Informed, voluntary, written, and signed consent was obtained from the health facilities administrators and study participants before the data collection.

RESULT

Socio-demographic Characteristics

A total of 464 obstetric health care providers were involved and yielding a response rate of 100%. Among them, 200 (43.1%) were males and 264 (56.9%) were females and male to female ratio is 0.76 to 1. More than half 233(53%) of the study participants were Midwives (Table 1).

Knowledge of Obstetric Care Providers

Almost all 444 (95.7%) health care providers knew about labor pain. Among them, 328 (73.9%) of obstetric care providers knew both pharmacological and non-pharmacological pain relief methods while 60 (13.5%) of them knew only pharmacological pain relief methods.

Among pharmacological pain relief methods, the majority 240 (61.9%) of obstetric care providers knew about non-steroidal anti-inflammatory drugs (NSAIDs), while 156 (40.2%) of them knew about systemic opioids and 140 (36.1%) of obstetric care providers know about epidural analgesia pain relief methods (Table 2).

Among OCPs who knew about labor pain, 356 (80.2%) of them knew about the side effect of pharmacological pain relief methods while 88 (19.8%) of them don't know about the side effect of pharmacological pain relief methods. Two hundred thirty-six (66.3%) of obstetric care providers respond that pharmacological pain relief methods causing the labor progress to be delayed while 208 (58.4%) of OCPs respond that pharmacologic pain relief methods cause fetal distress (Figure 1).

More than two-third 332 (71.6%) of obstetric care providers were asked to give pain relief methods. Most 252 (54.3%) obstetric care providers knew about the WHO pain ladder.

The mean score of knowledge questions is 6.65(SD= 2.95) with minimum and maximum scores of 0 and 15 respectively. More than 230 (49.6%) of the respondent have adequate knowledge.

The mean score of attitude questions is 4.01(SD= 1.4) with minimum and maximum scores of 0 and 6 respectively. Three hundred four (65.5%) of the respondent have an unfavorable attitude.

Utilization of Pharmacological Pain Relief Methods

From a total of 464 obstetric care providers, 364 (78.4%) of obstetric care providers were asked to provide pain relief for laboring mothers. More than three-fourth 344 (74.1%) of OCPs would provide labor pain relief methods if they had resources. The main barriers for patients receiving pain relief include high patient flow 172 (54.4%), limited numbers of staff 164 (51.9%), lack of knowledge and skill 104 (32.9%), no drug available to provide 88 (27.8%), and no equipment available 48 (15.2%).

Among those OCPs (364), 32 (8.8%) of obstetric care providers provide only pharmacological pain relief methods while 204 (44%) of OCPs provide both pharmacological and non-pharmacological pain relief methods. From pharmacologic pain relief methods, the most common pharmacologic pain relief methods were pethidine 132 (55.9%), diclofenac 76 (32.2%), and paracetamol 72 (30.5%) (Figure 2).

The mean score of utilization questions is 0.65(SD= 0.769) with minimum and maximum scores of 0 and 5 respectively. The overall utilization of pharmacological labor pain relief methods was 50.9% (95% CI 45.8,55.8).

Institutional Related Factors

Of the total of 464 study participants, 106 (91.4%) of the respondent replied that pharmacological drugs were available in their health facility whereas 10 (8.6%) of the participants responded that there were no drugs available in their health facility. It was reported that the most commonly available drugs were pethidine 292 (68.9%), diclofenac 324 (76.4%), paracetamol 300 (70.8%), and hyoscine 40 (9.4%). Nearly two-thirds 296 (63.8%) of OCPs got training about labor pain management methods from their respected facilities.

Obstetric Care Providers Preference

Most 276 (59.5%) of obstetric care providers prefer both pharmacological and non-pharmacological pain relief methods to manage labor pain while 136 (29.3%) of obstetric care

providers prefer non-pharmacological pain relief methods to manage labor pain and 52 (11.2%) of OCPs prefer pharmacological pain relief methods to manage labor pain.

Three hundred sixty-eight (79.3%) of OCPs expect the intensity of labor pain was severe pain while 60 (12.9%) of OCPs expect moderate pain and 36 (7.8%) of them expect mild pain.

Factors Associated with Utilization of Pharmacological labor pain management methods

Variables like age, sex, profession, year of experience, caregiver preferences, level of education, Pain expectation of caregiver, availability of equipment & analgesics drugs, level of knowledge, and attitude of the respondents were associated with utilization of pharmacological labor pain management methods in bivariate logistic regression analysis. In multivariate logistic regression sex, training, and attitude were factors associated with utilization of obstetric analgesia.

Multivariate analysis indicated that female OCPs were 2.07 times [AOR=2.07, 95% CI (1.31,3.26)] less likely to utilize pharmacological labor pain relief methods than male respondents. The odds of poor utilization of pharmacological labor pain relief methods among OCPs who had not trained were 1.60 times that of OCPs having training [AOR=1.60, 95%CI (1.02,2.53)].

Obstetric care providers with unfavorable attitudes were 9.1 times less likely to utilize pharmacological labor pain relief methods as compared to OCPs with favorable attitudes [AOR=9.1, 95%CI: (5.34,15.35)] (Table 3).

DISCUSSION

This study assessed the utilization of pharmacological labor pain management methods among OCPs in Harari regional state health facilities. Additionally, factors that affect the utilization of pharmacological labor pain management methods were isolated.

The overall utilization of pharmacological labor pain relief methods was 50.9% [(95% CI (45.8,55.8)]. Which was higher than studies done in Canada 16.7% [9], Nigeria 38.9% [5], Leku primary hospital, southern Ethiopia 25.7% [18], Amhara regional state zero [7], southern Ethiopia 37.9% [6]. This variation might be due to study period differences and sample size differences. Meanwhile, this study was lower than a study conducted in Ethiopia which was 66% [15]. This

discrepancy might be due to sampling size difference and the health facility setups because some of the setups could be well-equipped, the presence of skilled man powers (who took training may provide obstetric analgesia to laboring mothers).

In this study, the main barriers for laboring women receiving pain relief included high patient flow, limited numbers of staff, lack of knowledge and skill, no drug available to provide, and no equipment available. This is in line with studies done in Nigeria [19, 20], Egypt [13], and the Amhara region [7]. These findings indicated that a weak health care system, weak facility organizations, and limited distribution of resources obscure access to maternal health services. The health bureau of the region should work in addressing those problems by arranging in-service training to equip OCPs with knowledge and attitude and fulfilling necessary drugs and equipment.

In this study, training was found to be associated with the utilization of pharmacologic labor pain relief methods. This finding is in harmony with studies conducted in Amhara regional state [7], southern Ethiopia [6], and the Tigray region [21]. This may be due to failure to train obstetric care providers result in the inadequacy of the skill which in turn yields the inability to utilize pharmacologic labor pain relief methods

Our study pointed out that level of attitudes of care providers was associated with the utilization of pharmacologic labor pain relief methods This is consistent with studies done in Amhara regional state [7, 14], southern Ethiopia [6], and the Tigray region [21]. This may be due to if obstetric care providers had a favorable attitude, they can feel the need for pain relief methods, and then they would provide pain relief methods.

The limitation of this study was it might not indicate a cause-effect relationship because the study design was cross-sectional. Since the study is institution-based and the results might lack generalization to the entire population in the catchment area. The study didn't incorporate women in labor.

Conclusion

The study revealed that the overall prevalence of utilization of pharmacological labor pain management methods among obstetric care providers was low. In general, this study identified that variables like sex, training, and attitude of caregivers were factors associated with the utilization of pharmacological labor pain management methods. All stakeholders need to work

together to exploit OCPs' skills and attitudes. The health facilities should give the training to enhance the skill of obstetric care providers and equip themselves with sufficient analgesic drugs. Obstetric caregivers need to update and make themselves familiar with obstetric analgesia.

Abbreviations

AOR: Adjusted Odds Ratio; EmONC: Emergency Obstetric and Newborn Care LMIC; Low Mid Income Country; OCG: Obstetrics Care Givers; OCP: Obstetrics Care Providers; WHO: World Health Organization.

Declarations

Ethics Approval and Consent to Participate

Ethical clearance was taken from the Institutional Health Research Ethics Review Committee (IHRERC) of the College of Health and Medical Sciences of Haramaya University. In addition, permission to proceed was obtained from all public hospitals. And also, informed voluntary written and signed consent was obtained from each participant.

Consent for Publication

Not applicable.

Availability of Data and Materials.

All relevant data are included in this study. However, additional data will be available from the corresponding author on reasonable request.

Competing Interests.

The authors declare that there is no competing interest in this work.

Funding

The funder (Haramaya University) has no role in the design of the study, collection, analysis, and interpretation of data, and in writing the manuscript.

Author Contributions

Addis Eyeberu conceived the idea and had major roles in the data review, drafting, and writing the result. All co-authors contributed to data analysis, drafting, and manuscript preparation. Finally,

all authors read and approved the final version of the manuscript to be published and agreed on all aspects of this work.

Acknowledgments

We would like to extend our deepest thanks to Haramaya University, College of Health and Medical Science for helping us in performing this paper and we also would like to extend our acknowledgment to our friends for providing their constructive support.

REFERENCES

1. Labor, S. and S. Maguire, *The Pain of Labour*. Reviews in pain, 2008. **2**(2): p. 15-19.
2. Thomson, G., et al., *Women's experiences of pharmacological and non-pharmacological pain relief methods for labor and childbirth: a qualitative systematic review*. Reproductive Health, 2019. **16**(1): p. 71.
3. N, L., O. Yvonne, and S. Ghosh, *Physical and Psychological Aspects of Pain in Obstetrics*. 2012.
4. Czech, I., et al., *Pharmacological and Non-Pharmacological Methods of Labour Pain Relief-Establishment of Effectiveness and Comparison*. International journal of environmental research and public health, 2018. **15**(12): p. 2792.
5. Imarengiaye, C. and A. Ande, *Demand and utilisation of labour analgesia service by Nigerian women*. Journal of obstetrics and gynecology, 2006. **26**(2): p. 130-132.
6. Geltore, T.E., A. Taye, and A.G. Kelbore, *Utilization of obstetric analgesia in labor pain management and associated factors among obstetric care givers in public health facilities of Kembata Tembaro Zone, Southern Ethiopia*. Journal of pain research, 2018. **11**: p. 3089-3097.
7. Bitew, A., et al., *Utilization of obstetric analgesia in labor pain management and associated factors among obstetric caregivers in Amhara Regional State Referral Hospitals, Northwest Ethiopia. A hospital-based cross-sectional study*. Journal of Biomedical Sciences, 2016. **5**(2).
8. Adams, J., et al., *Use of pharmacological and non-pharmacological labor pain management techniques and their relationship to maternal and infant birth outcomes: an*

- examination of a nationally representative sample of 1835 pregnant women. Midwifery, 2015. 31(4): p. 458-463.*
9. Agaram, R., et al., *Inadequate pain relief with labor epidurals: a multivariate analysis of associated factors. International journal of obstetric anesthesia, 2009. 18(1): p. 10-14.*
 10. Barrett, S.J. and M.A. Stark, *Factors Associated With Labor Support Behaviors of Nurses. The Journal of perinatal Education, 2008. 19(1): p. 12-18.*
 11. FMOH, *ETHIOPIAN Emergency Obstetric and Newborn Care (EmONC) Assessment 2016. 2016.*
 12. Geltore, T.E., A.G. Kelbore, and A.T. Angelo, *Perceptions of obstetric analgesia: a qualitative study among midwives attending normal vaginal deliveries in Durame Hospital, Southern Ethiopia. J Pain Res, 2019. 12: p. 2187-2192.*
 13. Mousa, O., et al., *Perceptions and Practice of Labor Pain-Relief Methods among Health Professionals Conducting Delivery in Minia Maternity Units in Egypt. Obstet Gynecol Int, 2018. 2018: p. 3060953.*
 14. Bishaw, K.A., E.G. Sendo, and W.S. Abebe, *Knowledge, and use of labor pain relief methods and associated factors among obstetric care givers at public health centers of East Gojjam zone, Amhara region, Ethiopia: a facility-based cross-sectional study. BMC Pregnancy Childbirth, 2020. 20(1): p. 180.*
 15. McCauley, M., C. Stewart, and B. Kebede, *A survey of healthcare providers' knowledge and attitudes regarding pain relief in labor for women in Ethiopia. BMC Pregnancy Childbirth, 2017. 17(1): p. 56.*
 16. CSA, *CSA Website Analysis and Recommendation.pdf. 2009.*
 17. Getu, A.A., et al., *Non-pharmacological labor pain management and associated factor among skilled birth attendants in Amhara Regional State health institutions, Northwest Ethiopia. Reprod Health, 2020. 17(1): p. 183.*
 18. Siyoum, M. and S. Mekonnen, *Labor pain control and associated factors among women who gave birth at Leku primary hospital, southern Ethiopia. BMC Res Notes, 2019. 12(1): p. 619.*
 19. Ogboli-Nwasor, E., et al., *Pain relief in labor: a survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. Journal of Pain Research, 2011.*

20. Ohaeri, B., G. Owolabi, and J. Ingwu, *Skilled health attendants' knowledge and practice of pain management during labor in health care facilities in Ibadan, Nigeria*. European Journal of Midwifery, 2019. **3**(February).
21. Sahile, E., et al., *Practice of Labour Pain Management Methods and Associated Factors among Skilled Attendants Working at General Hospitals in Tigray Region, North Ethiopia: Hospital-Based Cross-Sectional Study Design*. Health Science Journal, 2017. **11**(4).

List of Tables

Table 1 Socio-demographic characteristics of obstetric care providers at Harari regional state public health facilities, 2021 (n=464)

Variables	Categories	Frequency	Percentage
Sex	Male	200	43.1
	Female	264	56.9
Age	15-24	60	12.9
	25-54	404	87.1
Religion	Orthodox	220	47.4
	Muslim	176	37.9
	Protestant	52	11.2
	Others	16	3.4
Profession	Midwife	233	53
	Nurse	198	42.67
	Obstetrician	20	4.3
	Health officers	13	2.8
Level of education	Diploma	142	30.6
	BSC	284	61.2
	MSC	38	8.2
Clinical experience	<2years	88	19
	≥2years	376	81

*others=Catholic and Adventist, BSC- Bachelor of Science, MSC- Masters of Sciences

Table 2 Knowledge about types of pharmacological pain relief methods among obstetric caregivers at Harari regional state health facilities (n=388) from May 20 to June 10, 2021

Variable	Categories	Frequency	Percentage
Systemic opioids	Yes	156	40.2
	No	232	59.8
Non-Steroidal Anti-inflammatory Drugs (NSAIDs)	Yes	240	61.9
	No	148	38.1
Epidural analgesia	Yes	140	36.1
	No	248	63.9
Inhalational analgesia	Yes	16	4.1
	No	372	95.9

Table 3 Factors associated with utilization of pharmacological pain relief methods among obstetric care providers at Harari regional state health facilities from May 20 to June 10, 2021.

Variables		Utilization of Pharmacological labor pain management		COR 95%CI	AOR 95%CI
		Poor	Good		
Sex	Male	112	88	1	1
	Female	116	148	1.62(1.12,2.35) *	2.07(1.31,3.26) *
Age	25-54	208	196	1	1
	15-24	20	40	2.12(1.19,3.76) *	0.51(0.26,1.01)
Level of education	BSc and above	168	148	1.67(1.12,2.47) *	1.22(0.76,1.95)
	Diploma	60	88	1	1
Training	No	108	60	2.64(1.78,3.91) *	1.60(1.02,2.53) *
	Yes	120	176	1	
Pain expectation	Moderate	29	31	0.412(0.17,0.984)	0.97(0.44,2.15)
	Sever	192	176	0.48(0.23,1.04)	1.28(0.69,2.38)
	Mild	25	11	1	1
Attitude	Unfavorable	200	104	9.07(5.66,14.53) *	9.1(5.34,15.35) *
	Favorable	28	132	1	1

Knowledge	Adequate	114	116	1	1
	Inadequate	114	120	1.03(0.72,1.49)	1.03(0.66,1.61)

*P≤ 0.001, OR=Odd ratio, AOR=Adjusted odd ratio, CI=Confidence interval




Figure 1 knowledge of OCP about the side effect of pharmacological pain relief methods among obstetric caregivers at Harari regional state health facilities from May 20 to June 10, 2021.




Figure 2 Most common pharmacological pain relief methods provided at Harari regional state health facilities from May 20 to June 10, 2021.

Utilization of pharmacological labour analgesia: a survey of obstetric care providers in eastern Ethiopia

Addis Eyeberu^{a,*}, Tamirat Getachew^a, Adera Debella^a, Bikila Balis^a, Bajirond Eshetu^a, Sinetibet Mesfin^a, Habtamu Bekele^a, Dawit Tamiru^a, Getahun Tiruye^a, Meron Degefa^a, Addisu Alemu^{b,*}, Merga Dheressa^a and Yadeta Dessie^b

^aSchool of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia; ^bSchool of Public Health, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

*Corresponding authors: Tel: +251910005436, +251911611147; E-mail: addiseyeberu@gmail.com, alemuadisu789@gmail.com

Received 3 December 2021; revised 3 August 2022; editorial decision 17 August 2022; accepted 19 August 2022

Background: Despite a crucial role played by obstetric care providers (OCPs) in providing pain relief methods during labour, the evidence suggests that there is a large number of women in Ethiopia who did not have access to pain relief methods. Moreover, to the best of the investigators' knowledge, evidence is scarce regarding the utilization of pharmacological labour pain management methods in the study area. Thus this study was carried out to determine the practice of pharmacological labour pain management methods among OCPs in Harari regional state health facilities in Ethiopia.

Methods: A facility-based cross-sectional study was conducted from 20 May to 10 June 2021 in Harari regional state health facilities. All (n=464) OCPs in Harari regional state health facilities were included in the study. Data were collected using a structured questionnaire adapted from previous studies. The data were entered into Epi-data version 3.1 statistical software. Statistical analysis was carried out using SPSS for Windows version 22. Bivariate and multivariate logistic regression analyses were employed to determine the association between independent variables and the outcome variable. A p-value <0.05 was used to establish statistical significance.

Results: In this study, 464 OCPs participated. The overall utilization of pharmacological labour pain relief methods was 50.9% (95% confidence interval [CI] 45.8 to 55.8). The main barriers to labouring mothers receiving pain relief include high patient flow (54.4%), the limited number of staff (51.9%) and a lack of knowledge and skill (32.9%). Being female (adjusted odds ratio [AOR] 2.07 [95% CI 1.31 to 3.26]), not having training (AOR 1.60 [95% CI 1.02 to 2.53]) and having an unfavourable attitude (AOR 9.1 [95% CI 5.34 to 15.35]) were the predictors of utilization of pharmacological labour pain management methods.

Conclusions: The study revealed that half of the OCPs utilized the pharmacological labour pain management method. In general, this study identified that variables like sex, training and the attitude of OCPs were factors associated with the utilization of pharmacological labour pain management methods. The health facilities should provide training to enhance the skill of OCPs and equip themselves with sufficient analgesic drugs.

Keywords: analgesia, Harar, labour pain, pharmacological labour pain management, utilization.

Introduction

Pain during labour and childbirth is a unique and the most severe pain that a woman ever has to face. The extent to which a woman feels in control of pain during labour is an indicator of maternal emotive well-being in childbirth.¹ Pharmacological labour pain management methods include oral tablets (paracetamol, codeine or tramadol), inhalations and injectable analgesics.²

Pain related to childbirth is present during pregnancy, labour and post-partum periods.³ Evidence showed that the care provided to the parturient during labour and delivery is one of the indicators of the satisfaction of childbirth.⁴ Pain during childbirth has an impact on maternal health, particularly its physiological and psychological influences.^{5,6} Furthermore, pain has a profound impact on maternal health, including anxiety, depression,

and alteration of respiratory patterns.^{6,7} Evidence shows that 51.9% of women had stress.⁸ Psychosocial factors have been implicated in the pain experienced during childbirth, which can have both short- and long-term consequences on the mother's health and her relationship with her infant.³

Evidence shows that labour pain analgesia has an effect on the mother and newborn, and labour outcomes include increased duration of labour, rates of instrumental deliveries, hypotension, the likelihood of maternal fever, neonatal sepsis and increased duration of breastfeeding.^{9–12} Even if those effects exist, labour pain is an individual phenomenon, so maternal request is considered a sufficient justification for pain relief.¹³ Furthermore, evidence shows that pain relief should be provided to labouring women.¹⁴

Even if obstetric care providers (OCPs) have an imperative role in providing pain relief options during labour, the majority of women in low-income countries, including Ethiopia, have not accessed pain relief methods. Evidence shows that 85.1% of the patients want their labour pain to be relieved, but only 38.9% received analgesia during labour.¹⁵ Studies done in Africa also report that the overall utilization of pharmacological labour pain methods varies and was 70.1% in Egypt¹⁶ and 48.4% and 49% in Nigeria.^{14,17} The utilization of pharmacological obstetric analgesia methods in Ethiopia also varies from place to place. A study done in Amhara and Tigray regions showed that the utilization of pharmacological labour analgesia was zero.^{13,18} A similar study done in Addis Ababa and southern Ethiopia showed that the overall utilization of pharmacological labour analgesia was 36.6% and 45.2%, respectively.^{13,19} The common factors associated with the non-utilization of pharmacological labour pain management methods were inadequate knowledge, negative attitude, unavailability of analgesic drugs in the health facilities, individual factors and institutional factors.^{12,13,18}

Ethiopia has developed and implemented standards of care, including the provision of physical and psychological support and the use of pharmacological and non-pharmacological comfort measures during labour and birth.²⁰ The Ethiopian Food, Medicine and Health Care Administration and Control Authority (FMHACA) also recommended the use of pharmacological labour pain management methods.²¹

Despite these efforts, the provision of pain relief in labour is neglected and has remained an ongoing challenge.^{16,22} A lack of awareness, unfavourable attitude and a lack of availability of pharmacological methods are considered to be the main reasons why women in Ethiopia do not receive adequate pain relief.^{13,22,23} Even though healthcare providers in Ethiopia understand that women suffer significant pain during labour, currently they do not provide effective pain relief.²⁴

Although it is known that effective labour pain management results in greater birth satisfaction for the mother, in many low- and middle-income countries (LMICs), including Ethiopia, OCPs' utilization of pharmacological labour pain management methods is low. Moreover, to the best of our knowledge, evidence is scarce regarding the utilization of pharmacological labour pain management methods in the study area. Thus this study was carried out to determine the utilization of pharmacological labour pain management methods among obstetric caregivers in Harari regional state health facilities.

Methods

Study design, study setting, and period

An institution-based cross-sectional study was conducted in Harari regional state health facilities from 20 May to 10 June 2021. The region is located 526 km east of Addis Ababa, the capital city of Ethiopia. The region is circled by Oromia regional state. The region is divided into nine *woredas*, with three of them being rural and six being urban. The urban districts are subdivided into 19 *kebeles* (the lowest administrative division in the country) and the rural districts are subdivided into 17 peasant associations (which is equivalent to *kebeles*). The total population (2021 projection based on the 2007 census) of the region is 270 000, of which 136 000 are males and 134 000 are females.²⁵ There are two public hospitals, two private hospitals and two governmental hospitals, eight public health centres and 20 health posts in the region. The number of healthcare providers in the Hiwot Fana Specialized University Hospital (HFSUH) and Jugol Hospital is 410 and 266, respectively.

Study population

All OCPs in Harari regional state health facilities during the study period were the study population. These include two public hospitals (HFSUH and Jugol Hospital), one private hospital (Harar General Hospital), two private clinics and nine health centres in the Harari region. All of the 464 obstetric caregivers available during the study period were included. OCPs not available during data collection due to illness, annual leave or left for training were excluded from the study.

Data collection methods and data quality assurance

A structured pretested self-administered questionnaire was prepared by adapting from different studies considering the local situation of the study area.^{18,23,26} The questionnaire had four essential components: providers' sociodemographic characteristics and knowledge, attitude and utilization-related questions.

The data were collected by five research-qualified professional nurses and supervised by two MSc-holding midwives. The quality of the data was guaranteed by pretesting using 5% of the respondents who were not eligible for the study. The completeness of the data was checked daily. Double data entry was done by two data clerks and the consistency of entered data was cross-checked by comparing two separately entered data using Epi-Data.

Variables

The outcome variable was the utilization of pharmacological labour pain management. Independent variables included sociodemographic factors such as age, sex, profession and year of experience; institutional factors included health institution policy, availability of equipment and analgesic drugs; and knowledge, attitude and individual factors included respondent's level of knowledge, qualifications, attitude and pain expectation.

Measurement

OCPs who were practicing greater than or equal to the mean value of listed labour pain management methods (0.65 [standard deviation {SD} 0.769]) were considered to have good utilization of pharmacological labour pain management methods,¹⁸ while OCPs who answered less than the mean values of practice-related labour pain relief method questions were considered to have poor utilization.

Knowledge was assessed by 10 items and the questions were answered with yes, no or I don't know responses. During analysis, the correct answer was coded as 1 and incorrect or I don't know answers were coded as 0. The sum score (range 0–10) was then computed and categorized. OCPs who scored ≥ 6.65 on the knowledge-related questions were considered to have adequate knowledge about labour pain relief methods, while OCPs who scored < 6.649 were considered to have inadequate knowledge about labour pain relief methods.¹⁸

Attitude was assessed by a total of six items and the questions were answered with yes, no or I don't know responses. During analysis, the correct answer was coded as 1 and incorrect or I don't know answers were coded as 0. The sum score (range 0–10) was then computed and categorized. OCPs who scored ≥ 4.1 on the attitude-related questions were considered to have a favourable attitude towards labour pain relief methods, while OCPs who scored < 4.1 were considered to have an unfavourable attitude towards labour pain relief methods.¹³

Statistical analysis

The data were entered into EpiData statistical software version 3.1 (<http://epidata.dk/>) and then exported to SPSS for Windows version 22 (IBM, Armonk, NY, USA) for analysis. Descriptive statistical analysis such as simple frequency was used to describe the characteristics of study participants, including sociodemographics, level of knowledge and attitudes of OCPs. Then the information was presented using frequencies, tables and figures.

Bivariate analysis and multivariate binary logistic regression analysis were performed to determine the association between each independent variable and the outcome variable. The assumptions of binary logistic regression were checked. The goodness of fit was checked by Hosmer–Lemeshow statistic and omnibus tests. All variables with a p-value < 0.25 in the bivariate analysis were included in the final model of multivariate analysis to control all possible confounders. A multicollinearity test was carried out to determine the correlation between independent variables by using the standard error and collinearity statistics (variance inflation factors > 10 and standard error > 2 were considered suggestive of the existence of multicollinearity). The direction and strength of the statistical association were measured by odds ratios (ORs) with 95% confidence intervals (CIs). Adjusted ORs along with 95% CIs were estimated to identify the association between independent variables and the utilization of pharmacological labour pain management by using multivariate analysis in binary logistic regression. In this study, p-values < 0.05 were considered statistically significant.

Table 1. Sociodemographic characteristics of OCPs at Harari regional state public health facilities, 2021 (N=464)

Variables	Categories	n	%
Sex	Male	200	43.1
	Female	264	56.9
Age (years)	15–24	60	12.9
	25–54	404	87.1
Religion	Christian	220	47.4
	Muslim	176	37.9
	Protestant	52	11.2
	Others ^a	16	3.5
Profession	Midwife	233	50.2
	Nurse	198	42.7
	Obstetrician	20	4.3
	Health officer	13	2.8
Level of education	Diploma	142	30.6
	BSc	284	61.2
	MSc	38	8.2
Clinical experience (years)	< 2	88	19
	≥ 2	376	81

^aCatholic and Adventist.
BSc: Bachelor of Science; MSc: Master's of Science.

Ethical considerations

Ethical approval to conduct this study was obtained from the Haramaya University, College of Health and Medical Sciences, Institutional Health Research Ethics Review Committee (IHRERC/066/2021). Official letters to conduct a study were also obtained from health facilities as needed. The health facilities were informed about the purpose of the study and the risks and benefits of the study. Informed, voluntary, written and signed consent was obtained from the health facilities' administrators and study participants before data collection.

Results

Sociodemographic characteristics

A total of 464 OCPs were involved, yielding a response rate of 100%. Among them, 200 (43.1%) were males and 264 (56.9%) were females and the male:female ratio was 0.76:1. More than half (233 [53%]) of the study participants were midwives (Table 1).

Knowledge of OCPs

Almost all (444 [95.7%]) OCPs knew about labour pain management methods. Of these, 328 (73.9%) knew both pharmacological and non-pharmacological pain relief methods and 60 (13.5%) of them knew only pharmacological pain relief methods. Among pharmacological pain relief methods, the majority (240 [61.9%]) of OCPs knew about non-steroidal anti-inflammatory drugs (NSAIDs), while 156 (40.2%) of them knew

Table 2. Knowledge about types of pharmacological pain relief methods among OCPs at Harari regional state health facilities (n=388) from 20 May to 10 June 2021

Variables	Categories	n	%
Systemic opioids	Yes	156	40.2
	No	232	59.8
Non-steroidal anti-inflammatory drugs	Yes	240	61.9
	No	148	38.1
Epidural analgesia	Yes	140	36.1
	No	248	63.9
Inhalational analgesia	Yes	16	4.1
	No	372	95.9

about systemic opioids and 140 (36.1%) knew about epidural analgesia pain relief methods (Table 2).

Among OCPs who knew about labour pain management, 356 (80.2%) knew about the side effects of pharmacological pain relief methods, while 88 (19.8%) of them did not know about the side effects. A total of 236 (66.3%) OCPs responded that pharmacological pain relief methods cause the labour progress to be delayed, while 208 (58.4%) responded that pharmacological pain relief methods cause foetal distress (Figure 1).

More than two-thirds (332 [71.6%]) of OCPs had been asked to provide pain relief and most (252 [54.3%]) OCPs knew about the World Health Organization pain ladder.

The mean score of knowledge questions was 6.65 (SD 2.95), with minimum and maximum scores of 0 and 15, respectively. A total of 230 (49.6%) of the respondent had adequate knowledge.

Regarding attitudes of OCPs, more than half (280 [60.3%]) of OCPs believe that analgesics are necessary to manage labour pain and more than two-thirds (84.5%) believe that they have the

responsibility and obligation to manage labour pain. The mean score of attitude questions was 4.01 (SD 1.4), with minimum and maximum scores of 0 and 6, respectively. A total of 304 (65.5%) of the respondent had an unfavourable attitude (Table 3).

Utilization of pharmacological pain relief methods

Of 464 OCPs, 364 (78.4%) were asked to provide pain relief for labouring mothers. Almost three-fourths (344 [74.1%]) of OCPs would provide labour pain relief methods if they had the resources. The main barriers for patients receiving pain relief include high patient flow (172 [54.4%]), limited staff (164 [51.9%]), lack of knowledge and skill (104 [32.9%]), no drugs available (88 [27.8%]) and no equipment available (48 [15.2%]).

Among the 364 OCPs, 32 (8.8%) provide only pharmacological pain relief methods while 204 (44%) provide both pharmacological and non-pharmacological pain relief methods. The most common pharmacological pain relief methods were pethidine (132 [55.9%]), diclofenac (76 [32.2%]) and paracetamol (72 [30.5%]) (Figure 2).

The mean score of utilization questions was 0.65 (SD 0.769), with minimum and maximum scores of 0 and 5, respectively. The overall utilization of pharmacological labour pain relief methods was 50.9% (95% CI 45.8 to 55.8).

Institutional-related factors

Of the 464 study participants, 106 (91.4%) replied that pharmacological drugs were available in their health facility, whereas 10 (8.6%) of the participants responded that there were no drugs available in their health facility. It was reported that the most commonly available drugs were pethidine (292 [68.9%]), diclofenac (324 [76.4%]), paracetamol (300 [70.8%]) and hyoscine (40 [9.4%]). Nearly two-thirds (296 [63.8%]) of OCPs got training about labour pain management methods from their respective facilities.




Figure 1. Knowledge of the side effects of pharmacological pain relief methods among OCPs at Harari regional state health facilities from 20 May to 10 June 2021.

Table 3. Attitudes of OCPs about pharmacological pain relief methods in Harari regional state public health facilities, 2021

Variables	Categories	n	%
Believe that labour pain management methods alleviate labour pain (n=464)	Yes	392	84.5
	No	72	15.5
Believe every mother during labour should be managed (n=464)	Yes	320	69
	No	144	31
Believe that labour pain is natural and the mother has to face it (n=464)	Yes	344	74.1
	No	120	25.9
Believe pharmacology methods are necessary for managing labour pain (n=464)	Yes	280	60.3
	No	184	39.7
Believe that you have a responsibility and obligation to manage labour pain (n=464)	Yes	392	84.5
	No	72	15.5

pharmacological pain relief methods




Figure 2. Most common pharmacological pain relief methods provided at Harari regional state health facilities from 20 May to 10 June 2021.

OCP preference

Most OCPs (276 [59.5%]) prefer both pharmacological and non-pharmacological pain relief methods to manage labour pain while 136 (29.3%) OCPs prefer non-pharmacological pain relief methods and 52 (11.2%) OCPs prefer pharmacological pain relief methods. A total of 368 (79.3%) OCPs expected the intensity of labour pain to be severe, while 60 (12.9%) expected moderate pain and 36 (7.8%) expected mild pain.

Factors associated with the utilization of pharmacological labour pain management methods

Variables such as age, sex, profession, year of experience, caregiver preferences, level of education, pain expectation of the

caregiver, availability of equipment and analgesic drugs, level of knowledge and attitude of the caregiver were associated with the utilization of pharmacological labour pain management methods in bivariate logistic regression analysis. In multivariate logistic regression sex, training and attitude were factors associated with the utilization of obstetric analgesia.

Multivariate analysis indicated that female OCPs were 2.07 times (adjusted OR [AOR] 2.07 [95% CI 1.31 to 3.26]) more likely to utilize pharmacological labour pain relief methods than male respondents. The odds of poor utilization of pharmacological labour pain relief methods among OCPs who had not been trained were 1.60 times those of OCPs who had training (AOR 1.60 [95% CI 1.02 to 2.53]). OCPs with unfavourable attitudes were 9.1 times less likely to utilize pharmacological labour pain relief methods as compared with OCPs with favourable attitudes (AOR 9.1 [95% CI 5.34 to 15.35]) (Table 4).

Discussion

This study assessed the utilization of pharmacological labour pain management methods among OCPs in Harari regional state health facilities. Additionally, factors that affect the utilization of pharmacological labour pain management methods were isolated.

The overall utilization of pharmacological labour pain relief methods was 50.9% (95% CI 45.8 to 55.8), which was higher than in studies in Canada (16.7%),²⁷ Nigeria (38.9%),¹⁵ Addis Ababa, Ethiopia (36.6%),¹⁹ Amhara regional state (0%)¹⁸ and southern Ethiopia (37.9%).¹³ This variation might be due to study period differences and sample size differences. Meanwhile, the utilization in this study was lower than in studies conducted in Egypt (70.1%) and another study in Ethiopia (66%).²⁴ This discrepancy might be due to sampling size differences and health facilities because some of the health facilities may be well equipped, with skilled manpower.

In this study, the main barriers for labouring women receiving pain relief included high patient flow, limited staff, a lack of knowledge and skill, no drugs available and no equipment available. This is in line with studies done in Nigeria,^{28,29} Egypt¹⁶ and the Amhara region.¹⁸ These findings indicated that a weak

Table 4. Factors associated with utilization of pharmacological pain relief methods among OCPs at Harari regional state health facilities from 20 May to 10 June 2021

Variables		Utilization of pharmacological labour pain management, n		OR (95% CI)	AOR (95% CI)
		Poor	Good		
Sex	Male	112	88	1	1
	Female	116	148	1.62 (1.12 to 2.35)*	2.07 (1.31 to 3.26)*
Age (years)	25–54	208	196	1	1
	15–24	20	40	2.12 (1.19 to 3.76)*	0.51 (0.26 to 1.01)
Level of education	BSc and above	168	148	1.67 (1.12 to 2.47)*	1.22 (0.76 to 1.95)
	Diploma	60	88	1	1
Training	No	108	60	2.6(1.78 to 3.91)*	1.6 (1.02 to 2.53)*
	Yes	120	176	1	1
Pain expectation	Moderate	29	31	0.412 (0.17 to 0.984)	0.97 (0.44 to 2.15)
	Severe	192	176	0.48 (0.23 to 1.04)	1.28 (0.69 to 2.38)
	Mild	25	11	1	1
Attitude	Unfavourable	200	104	9.07 (5.66 to 14.53)*	9.1 (5.34 to 15.35)*
	Favourable	28	132	1	1
Knowledge	Adequate	114	116	1	1
	Inadequate	114	120	1.03 (0.72 to 1.49)	1.03 (0.66 to 1.61)

* $p \leq 0.001$.

healthcare system, weak facility organizations and limited distribution of resources limit access to maternal health services. Arranging in-service training to equip OCPs with knowledge and providing necessary drugs and equipment could be the solution.

In this study, training was found to be associated with the utilization of pharmacological labour pain relief methods. This finding is in harmony with studies conducted in the Amhara region,¹⁸ southern Ethiopia¹³ and the Tigray region.³⁰ This may be due to failure to train OCPs, resulting in inadequate skills.

Our study pointed out that the attitudes of care providers was associated with the utilization of pharmacological labour pain relief methods, as those with unfavourable attitudes were less likely to utilize pharmacological pain relief methods. This is consistent with studies done in the Amhara region,^{18,23} southern Ethiopia¹³ and the Tigray region.³⁰ Moreover, female OCPs were more likely to utilize pharmacological labour pain relief methods than male respondents. This may be because female OCPs may have giving birth in the past and experienced the pain and they may not want their clients to suffer the pain.

The limitation of this study was that the assessment of the utilization of pharmacological labour pain management methods only encompassed OCPs and did not include the service receivers (mothers facing labour pain during childbirth).

Conclusions

The study revealed that the overall prevalence of utilization of pharmacological labour pain management methods among

OCPs was low. In general, this study identified that variables like sex, training and the attitude of caregivers were factors associated with the utilization of pharmacological labour pain management methods. All stakeholders need to work together to enhance the skills and attitudes of OCPs. The health facilities should provide training to enhance the skills of OCPs and equip themselves with sufficient analgesic drugs and OCPs need to familiarize themselves with obstetric analgesia.

Authors' contributions: AE conceived the idea and was involved in the data review and drafting and writing the manuscript. TG, AD, BB, BE, SM and HB were responsible for proposal writing, project administration, methodology, data analysis and drafting the manuscript. AE, DT, GT, MD, AA, MDA and YD contributed to data analysis, drafting and manuscript preparation. Finally, all authors read and approved the final version of the manuscript to be published and agreed on all aspects of this work.

Acknowledgments: We would like to extend our deepest thanks to Haramaya University, College of Health and Medical Science, data collectors, and participants, for helping us with this research and we would also like to extend thanks to our friends for providing their constructive support.

Funding: The funder (Haramaya University) had no role in the design of the study; collection, analysis and interpretation of data; or writing the manuscript.

Competing interests: None declared.

Data availability: All relevant data are included in this study. However, additional data is available from the corresponding author upon reasonable request.

References

- Labor S, Maguire S. The pain of labour. *Rev Pain*. 2008;2(2):15–19.
- Thomson G, Feeley C, Moran VH, et al. Women's experiences of pharmacological and non-pharmacological pain relief methods for labor and childbirth: a qualitative systematic review. *Reprod Health*. 2019;16(1):71.
- Ebirim LN, Yvonne O, Ghosh S. Physical and psychological aspects of pain in obstetrics. In: Ghosh S, editor. *Pain in perspective*. Available from: <https://www.intechopen.com/chapters/40395> [accessed 29 August 2022].
- Czech I, Fuchs P, Fuchs A, et al. Pharmacological and non-pharmacological methods of labour pain relief-establishment of effectiveness and comparison. *Int J Environ Res Public Health*. 2018;15(12):2792.
- Olde E, van der Hart O, Kleber R, et al. Posttraumatic stress following childbirth: a review. *Clin Psychol Rev*. 2006;26(1):1–16.
- Lowe NK. The nature of labor pain. *Am J Obstet Gynecol*. 2002;186(5 Suppl):S16–24.
- Junge C, von Soest T, Weidner K, et al. Labor pain in women with and without severe fear of childbirth: a population-based, longitudinal study. *Birth*. 2018;45(4):469–77.
- Tan HS, Agarhesh T, Tan CW, et al. Perceived stress during labor and its association with depressive symptomatology, anxiety, and pain catastrophizing. *Sci Rep*. 2021;11(1):17005.
- Lieberman E, O'Donoghue C. Unintended effects of epidural analgesia during labor: a systematic review. *Am J Obstet Gynecol*. 2002;186(5 Suppl):S31–68.
- Leighton BL, Halpern SH. The effects of epidural analgesia on labor, maternal, and neonatal outcomes: a systematic review. *Am J Obstet Gynecol*. 2002;186(5 Suppl):S69–77.
- Leeman L, Fontaine P, King V, et al. The nature and management of labor pain: part II. Pharmacologic pain relief. *Am Fam Physician*. 2003;68(6):1115–20.
- Adams J, Frawley J, Steel A, et al. Use of pharmacological and non-pharmacological labor pain management techniques and their relationship to maternal and infant birth outcomes: an examination of a nationally representative sample of 1835 pregnant women. *Midwifery*. 2015;31(4):458–63.
- Geltore TE, Taye A, Kelbore AG. Utilization of obstetric analgesia in labor pain management and associated factors among obstetric caregivers in public health facilities of Kembata Tembaro Zone, Southern Ethiopia. *J Pain Res*. 2018;11:3089–97.
- Ogboli-Nwasor E, Adaji SE, Bature SB, et al. Pain relief in labor: a survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. *J Pain Res*. 2011;4:227.
- Imarengiaye C, Ande A. Demand and utilisation of labour analgesia service by Nigerian women. *J Obstet Gynecol*. 2006;26(2):130–2.
- Mousa O, Abdelhafez AA, Abdelraheim AR, et al. Perceptions and practice of labor pain-relief methods among health professionals conducting delivery in Minia maternity units in Egypt. *Obstet Gynecol Int*. 2018;2018:3060953.
- Lawani LO, Eze JN, Anozie OB, et al. Obstetric analgesia for vaginal birth in contemporary obstetrics: a survey of the practice of obstetricians in Nigeria. *BMC Pregnancy Childbirth*. 2014;14:140.
- Bitew A, Workie A, Seyum T, et al. Utilization of obstetric analgesia in labor pain management and associated factors among obstetric caregivers in Amhara Regional State Referral Hospitals, Northwest Ethiopia. A hospital-based cross-sectional study. *J Biomed Sci*. 2016;5(2):15.
- Gido R, Yadeta TA, Tura AK. Utilization of obstetric analgesia for labor pain management and associated factors among obstetric care providers in public hospitals of Addis Ababa, Ethiopia: a cross-sectional study. *Obstet Gynecol Int*. 2021;2021:9973001.
- Ministry of Health. Ethiopian Emergency Obstetric and Newborn Care (EmONC) Assessment 2016. Addis Ababa: Ethiopian Public Health Institute; 2016.
- Food, Medicine and Healthcare Administration and Control Authority of Ethiopia. Standard treatment guidelines for general hospitals. Addis Ababa: Ethiopian Federal Ministry of health; 2014.
- Geltore TE, Kelbore AG, Angelo AT. Perceptions of obstetric analgesia: a qualitative study among midwives attending normal vaginal deliveries in Durame Hospital, Southern Ethiopia. *J Pain Res*. 2019;12:2187–92.
- Bishaw KA, Sendo EG, Abebe WS. Knowledge, and use of labor pain relief methods and associated factors among obstetric caregivers at public health centers of East Gojjam zone, Amhara region, Ethiopia: a facility based cross-sectional study. *BMC Pregnancy Childbirth*. 2020;20:180.
- McCauley M, Stewart C, Kebede B. A survey of healthcare providers' knowledge and attitudes regarding pain relief in labor for women in Ethiopia. *BMC Pregnancy Childbirth*. 2017;17:56.
- CSA. *CSA Population Projections for Ethiopia 2007–2037*. Available from: <https://www.statsethiopia.gov.et/wp-content/uploads/2019/05/ICPS-Population-Projection-2007-2037-produced-in-2012.pdf>.
- Getu AA, Getie SA, Gela GB, et al. Non-pharmacological labor pain management and associated factor among skilled birth attendants in Amhara Regional State health institutions, Northwest Ethiopia. *Reprod Health*. 2020;17:183.
- Agaram R, Douglas MJ, McTaggart RA, et al. Inadequate pain relief with labor epidurals: a multivariate analysis of associated factors. *Int J Obstet Anesth*. 2009;18(1):10–14.
- Ogboli-Nwasor E, Adaji SE, Bature SB, et al. Pain relief in labor: a survey of awareness, attitude, and practice of health care providers in Zaria, Nigeria. *J Pain Res*. 2011;4:227–32.
- Ohaeri B, Owolabi G, Ingwu J. Skilled health attendants' knowledge and practice of pain management during labor in health care facilities in Ibadan, Nigeria. *Eur J Midwifery*. 2019;3:3.
- Sahile E, Yemaneh Y, Alemu AA, et al. Practice of labour pain management methods and associated factors among skilled attendants working at general hospitals in Tigray Region, North Ethiopia: hospital based cross-sectional study design. *Health Sci J*. 2017;11(4).

2. Journal of health science and medical research

Q4 : SJR 0,15

Thailand

Invitation to Review for the J x

Pesan Asli x

(53) WhatsApp x

google translate - Penelusur x

Unduh file | iLovePDF x

+ -

mail.google.com/mail/u/0/#search/Journal+of+health+science+and+medical+research/FMfcgzGwHfqWHgDnLkmPKFTRrNRHQklh

Gmail

99+

Tulis

Mail

Chat

Meet

Kotak Masuk 1.729

Berbintang

Ditunda

Terkirim

Draf

Selengkapnya

Label +

Journal of health science and medical research

Aktif

Summarize 21 dari banyak

Invitation to Review for the Journal of Health Science and Medical Research

Eksternal Kotak Masuk x

Wit Wichaidit <onbehalf@manuscriptcentral.com>

Sel, 14 Nov 2023, 06.46

kepada saya

Terjemahkan ke Indonesia

14-Nov-2023

Dear Dr. Risnah:

Manuscript ID JHSMR-2023-0601 entitled "Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia" has been submitted to the Journal of Health Science and Medical Research.

I invite you to review this manuscript. The abstract appears at the end of this letter. Please let me know as soon as possible if you will be able to accept my invitation to review. If you are unable to review at this time, I would appreciate you recommending another expert reviewer. You may e-mail me with your reply or click the appropriate link at the bottom of the page to automatically register your reply with our online manuscript submission and review system.

Once you accept my invitation to review this manuscript, you will be notified via e-mail about how to access ScholarOne Manuscripts, our online manuscript submission and review system. You will then have access to the manuscript and reviewer instructions in your Reviewer Center.

I realize that our expert reviewers greatly contribute to the high standards of the Journal, and I thank you for your present and/or future participation.

Type here to search

USD/JPY -0.67%

21:24 30/05/2024

Manuscript ID JHSMR-2023-
Pesan Asli
(54) WhatsApp
google translate - Penelusur
Unduh file | iLovePDF

mail.google.com/mail/u/0/#search/Journal+of+health+science+and+medical+research/FMfcgzGwHpPGSbKqcTbQFgLCsdFmngKp

Journal of health science and medical research

Aktif

AT ALUDDIN

99+

Mail

Chat

Meet

Tulis

Kotak Masuk 1.724

Berbintang

Ditunda

Terkirim

Draf

Selengkapnya

Label +

Summarize

19 dari banyak

Manuscript ID JHSMR-2023-0601 now in your Reviewer Center - Journal of Health Science and Medical Research

Eksternal

Kotak Masuk x

Wit Wichaidit <onbehalf@manuscriptcentral.com>

kepada saya

Sel, 21 Nov 2023, 20.24

Terjemahkan ke Indonesia

21-Nov-2023

Dear Dr. Risnah:

Thank you for agreeing to review Manuscript ID JHSMR-2023-0601 entitled "Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia" for Journal of Health Science and Medical Research. Please try your best to complete your review by 05-Dec-2023.

In your review, please answer all questions. On the review page, there is a space for "Comments to Editor" and a space for "Comments to the Author." Please be sure to put your comments to the author in the appropriate space.

To access just the manuscript for review directly with no need to enter log in details, click the link below:

https://mc03.manuscriptcentral.com/jhsmr?URL_MASK=f4e4339800094a369bfcc517bce60e0.

To login to your account on the Journal of Health Science and Medical Research - ScholarOne Manuscripts site at <https://mc03.manuscriptcentral.com/jhsmr>

Type here to search

27°C Cerah

21:26
30/05/2024

Reminder: Review Due Today x | Pesan Asli x | (54) WhatsApp x | google translate - Penelusur x | Unduh file | iLovePDF x | + -

mail.google.com/mail/u/0/#search/Journal+of+health+science+and+medical+research/FMfcgzGwJRvgSWTrKRMkrfhdNWGRzRjG

Gmail

Journal of health science and medical research

Aktif

99+ Tulis

Mail

Chat

Meet

Kotak Masuk 1.724

Berbintang

Ditunda

Terkirim

Draf

Selengkapnya

Label +

Reminder: Review Due Today - Journal of Health Science and Medical Research

Eksternal Kotak Masuk x

Wit Wichaidit <onbehalf@manuscriptcentral.com> kepada saya

Sab, 23 Des 2023, 13.02

Terjemahkan ke Indonesia

23-Dec-2023

Dear Dr. Risnah:

Recently, you agreed to review Manuscript ID JHSMR-2023-0601.R1, entitled "Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia." This e-mail is a reminder that your review is now due. Please complete and submit your review.

To access just the manuscript for review directly with no need to enter log in details, click the link below:

https://mc03.manuscriptcentral.com/jhsmr?URL_MASK=90f73eb4553e4484833767e707df1c5b

To login to your account on the Journal of Health Science and Medical Research - ScholarOne Manuscripts site at <https://mc03.manuscriptcentral.com/jhsmr>, your case-sensitive USER ID is risnah@uin-alauddin.ac.id. For security purposes your password is not listed in this email. If you are unsure of your password you may click the link below to set a new password.



Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia

Journal:	<i>Journal of Health Science and Medical Research</i>
Manuscript ID	JHSMR-2023-0601
Manuscript Type:	Original Article
Keywords:	Accident prevention, Coal mining, Manipulation of data, Mobile application
Speciality:	Occupational health and safety, Accident reporting, Coal mining, Zero accident

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Preventing Practice of Manipulating Accident Data in the Coal Mining Sector Using Mobile Applications in East Kalimantan, Indonesia

ABSTRACT

Objective: The practice of manipulating data on work accidents and potential hazards still occurs, especially in the coal mining sector. A mobile application for reporting work accidents and potential hazards is needed so that data manipulation practices can be prevented.

Materials and Methods: The research design used was exploratory which aimed to introduce and explore the capabilities of the SIKAP application (Information System for Occupational Accidents, Occupational Diseases and Potential Hazards) to prevent data manipulation practices. Researchers involved thirty coal mine workers using the SIKAP application which had been tested on three information systems experts using the black box testing method.

Results: Every worker has the opportunity to report work accidents and potential dangers via the SIKAP application on the worker's account. Companies can receive and follow up on worker reports via the company account. Regional governments in the field of labor development and inspection can monitor reports through government accounts. The SIKAP application is integrated and synchronized between workers, companies and the government.

Conclusion: The SIKAP application is able to prevent the practice of manipulating work accident reporting data and potential hazards in the coal mining sector.

Keywords: Accident prevention, Coal mining, Manipulation of data, Mobile application

Introduction

Coal companies in Indonesia are still one of the most lucrative business sectors and have even become one of the favorite business sectors in East Kalimantan to this day. However, the coal mining sector is also one of the sectors that contributes to the high number of work accidents and work-related diseases. It is known that all stages of mining activities contain dangers that have a high risk of causing workers to suffer minor, serious injuries and even death.

The Indonesian government continues to make improvements so that the mining sector is no longer a scary place to work, especially for mining workers. One of them is by issuing regulations such as the Regulation of the Minister of Energy and Mineral Resources of the Republic of Indonesia Number 26 of 2018 concerning the Implementation of Good Mining Principles and Supervision of Mineral and Coal Mining, and followed by the Decree of the Directorate General of Minerals and Coal Number 185.K/ 37.04/DJB/2019 concerning Technical Guidelines for the Implementation of Mining Safety and Implementation, Assessment and Reporting of Mineral and Coal Mining Safety Management Systems.

Apart from the government's role, companies also have the responsibility to implement occupational safety and health (OSH) efforts to ensure that workers continue to receive protection and safety while working. This has been regulated in Law Number 13 of 2003 concerning Employment¹, and Government Regulation Number 50 of 2012 concerning the Implementation of an Occupational Safety and Health Management System which is integrated with the company management system.²

One of the efforts to prevent work accidents and work-related diseases is through reporting work accidents, work-related diseases and potential dangers, both routine reports and special reports when work accidents occur. This reporting contains a number of information related to work accidents,³ and this information will help company management to carry out immediate control over potential dangers so as not to result in work accidents in the future.⁴ However, in practice there are still problems with reporting that is not transparent and tends to be manipulative.

Space for the practice of manipulating data on work accidents and work-related illnesses reported from companies to the government as the agency supervising coal mining operations remains wide open. One of the reasons is that the reporting system used is still conventional, utilizing written work accident and occupational disease reporting forms.⁵ The practice of manipulating report data can not only be carried out by workers, but also company K3 officers and/or government employees.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A real-time, integrated and synchronized information system for reporting work accidents, work-related illnesses and potential hazards is needed to prevent the practice of data manipulation when reporting is carried out. This is also in accordance with previous research recommendations which stated that one of the efforts to control the practice of manipulating work accident and occupational disease data is to utilize a digitalization-based reporting information system.⁶ Based on existing problems, research is needed to introduce and explore the capabilities of the SIKAP application to prevent data manipulation practices in coal mining in East Kalimantan, Indonesia.

Methods

This research uses an exploratory design by utilizing the SIKAP application in the reporting system for work accidents, work-related diseases and potential hazards in coal mining. This research involved thirty coal mining workers from several coal companies operating in Samarinda City, East Kalimantan Province, Indonesia. The workers involved were given the SIKAP reporting application which had previously been tested on three information systems experts using the black box testing method to test that the SIKAP application could be used properly. Every worker is given the opportunity to use the SIKAP application to report work accidents, work-related illnesses, and potential dangers that have been experienced or seen while working at a coal company. The research was carried out in October 2023 based on a permit to carry out research issued by the Faculty of Economics and Business, Mulawarman University Number: 202/UN17.1.36.1/DT/2023. The results of trials using the SIKAP application on coal mining workers were then analyzed to assess the ability of the SIKAP application to prevent the practice of manipulating work accident reporting data, work-related diseases and potential hazards in coal mining.

Result

The SIKAP application is an acronym for Information System for Reporting Work Accidents, Occupational Diseases and Potential Hazards. The initial display when opening the SIKAP application consists of information about SIKAP, information about OSH activities, OSH policies, and other information related to OSH. There is an access menu at the bottom of the information consisting of worker, company and government accounts.

The worker account is preceded by creating a username, name, work unit, position, cellphone number, uploading a photo, and account password. Next, the worker clicks Log in by entering the username and password for the account that has been created. After logging in,

several application menus will appear consisting of Home, Announcements, Notifications, and Profile. The Home menu consists of Today's Data, Work Accident Report, Occupational Disease Report, Potential Hazard Report, and Report History.

Today's data includes data on work accidents, occupational disease, and potential dangers that have been reported by the worker and other workers at the company where he works. If the worker wants to report a work accident, then click the Report Work Accident menu. In this menu, workers enter a number of information, namely the type of work accident, victims, date of incident, time of incident, location of incident, body parts injured as a result of the accident, and actions taken by workers when experiencing an accident by attaching documentation/photos of the time of the incident. After completing all the required information, the worker then clicks Report.

For workers who want to report work-related illnesses, click Report of occupational disease. In the occupational disease report menu, a number of information will be entered by the worker, namely the type of occupational disease, details of the occupational disease, the complaint felt, the date and time of the complaint, and the actions taken by the worker when experiencing the complaint accompanied by documentation/photos. After completing all the required information, the worker then clicks Report. If the worker wants to report a potential hazard, then click Report Potential Hazard and fill in some information, namely location, description of the type of hazard, risks, actions taken, and attach documentation/photos of witnessing the potential hazard. After completing all the required information, the worker then clicks Report.

Worker accounts also have a Report History menu. This menu functions to check/monitor the status or reporting history of work accidents, occupational disease, and potential hazards that have been carried out by workers to the company's OSH unit. Workers can see the progress of the reports that have been carried out. The history of Work Accident Reports, occupational disease, and Potential Hazards consists of the statuses Sent, Processed, and Completed/Rejected. In this section, workers can see a description of the accident, occupational disease and potential hazard report information that has been filled in when reporting.

The report that has been sent by the worker can then be seen that the report was processed/received by officers in the OSH unit accompanied by the identity of the report recipient. Information was also obtained regarding suggestions/directions from officers in the OSH unit to workers/reporters. Workers will also receive information that the report has been completed by the OSH officer, accompanied by a description of the final action report from the

OSH officer. In this section there is also information about rejection of reports made by OSH officers if the report is deemed to have been repeatedly reported or has been completed by OSH officers.

Available Announcements menu, workers will see schedules or other important agendas informed by the company's OSH officers. On the Notification menu, workers will receive notifications automatically from reports that have been made previously. The profile will display a number of workers' personal information, including name, work unit and position. Details of the worker's SIKAP account can be seen in Figure 1 below:

Creating and using a company account is almost the same as a worker account, namely creating a username, name, work unit, position, cellphone number, uploading photos and account password. After registering the account, the company OSH officer then clicks Log in by entering the username and password for the account that has been created. When you log in, several application menus will appear consisting of Home, Announcements, Notifications and Profile.

The Home menu consists of Today's Data, Incoming Report, Process Report, and Completed Report. Today's Data Menu which includes data on work accidents, occupational disease, and potential hazards that have been reported by workers at the company. In this section, company OSH officers can search for accident, occupational disease and potential hazard data, both daily data and data according to the desired time period by clicking Search Data. The desired data can be displayed in graphical form by clicking the Graph button.

In the Incoming Reports, Process Reports and Completed Reports menu, company OSH officers can see a number of incoming report data from workers. OSH officers can also accept or reject reports from workers by providing directions/information addressed to the worker. In the Announcements menu, OSH officers can delete or add other important events/agenda which will be informed to workers. In the Notification menu, the company's OSH officer will receive notification automatically of each stage of incoming reporting and the profile will display a number of company OSH officer information. Details of the company's SIKAP account can be seen in Figure 2 below:

Creating a government account represented by certain service employees who are given the task of regarding accident, occupational disease, and potential danger data by creating a username, name, work unit, position, cellphone number, uploading photos and account password. After registering an account, employees in certain departments then click Log in by entering the username and password for the account that has been created. After logging in, several application menus will appear consisting of Home, Notifications and Profile.

The Home menu consists of Today's Data and Reports. Today's data includes data on work accidents, occupational disease, and potential dangers that have been reported by workers at certain companies. In this section, certain service employees can search for accident, occupational disease and potential hazard data, both daily data and data according to the specific time period desired by clicking Search Data. The desired data can be displayed in graphical form by clicking the Graph button. In the Reports menu, accident, occupational disease and potential hazard reports will be displayed in the Incoming, Processing and Completed Reports menu.

Certain service employees can view a number of incoming report data from workers at certain companies along with the history of each report. Service employees can also provide suggestions/input aimed at the company regarding incidents reported by their workers. In the Notification menu, certain service employees will receive automatic notification of each reporting process/stage that is entered into the SIKAP account belonging to a certain service. The profile will display a number of information about certain service employees who handle work accident data, occupational disease and potential hazards. Details of the government's SIKAP account can be seen in Figure 3 below:

Discussion

The Indonesian government continues to make improvements to improve the coal mining sector as one of the sectors at risk of work accidents and work-related diseases through regulations. In several countries such as China and Australia, regulations regarding coal mine safety have been developed.⁷ Regulations that are consistent in their implementation can influence workers' behavior at work.⁸

Reporting of work accidents, work-related diseases and potential hazards in the workplace, including in the coal mining sector, has been regulated in Minister of Manpower Regulation Number 3 of 1998 concerning Procedures for Reporting and Inspecting Accidents and has been used since 1998 until now using a written report format and an online reporting system. conventional from the company to the local regional government and the Ministry of Manpower of the Republic of Indonesia. Such reporting management has a number of weaknesses, such as the reporting process being slow, the possibility of incomplete information being obtained due to the disappearance of witnesses and loss of evidence when a work accident occurs, the unavailability of forms at the scene of the incident, and various other weaknesses.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Incomplete information regarding work accidents, work-related diseases and potential workplace hazards will have a negative impact on the company's OSH performance. It is difficult for company management to take action to prevent and handle work accidents and work-related diseases if they are not supported by detailed data and information regarding the root causes or potential hazards that result in work accidents and work-related diseases. On the other hand, having complete information about potential hazards in the work environment makes it easier for companies to identify work areas that are at risk of work accidents and proceed with corrective action so that similar incidents do not happen again.⁹

Reporting work accidents, work-related diseases and potential hazards is not only the company's responsibility to the government, especially when work accidents and work-related diseases occur, but also requires worker participation in reporting.¹⁰ This has also been regulated based on Decree of the Directorate General of Minerals and Coal Number 185.K/37.04/DJB/2019 that every mining worker is obliged to report any accidents or injuries caused by work or related to work. However, it is still found that mining workers think that reporting work accidents and potential dangers in the workplace is not important, so it is necessary to increase workers' perceptions about the importance of reporting.¹¹

Work accident reporting aims to make it easier for company management to identify work areas that allow work accidents,⁹ and at the same time identify the causes of accidents so that work accidents can be prevented early on.¹² In addition, Work accident reports can also estimate workers' unsafe actions.¹³ However, the K3 efforts carried out are still less effective in preventing work accidents, so a better investigation system is needed.¹⁴ Reporting coal mining accidents can also improve safety management in the sector mining.¹⁵

Reporting accidents and potential dangers by workers needs to receive support from the company.¹⁶ Workers' participation in reporting work accidents is not enough to create an K3 culture in the workplace, but the role of the company is also needed.¹⁷ Companies that have a high commitment to safety are positively correlated with workers' K3 behavior.¹⁸ Solutions to preventing work accidents and occupational diseases require partnerships and synergy between workers, government, companies and other parties.¹⁹

Challenge faced in reporting work accidents, work-related diseases and potential hazards in coal mining is the large number of cases that occur but are hidden or not reported.²⁰ The practice of hiding, deleting data, changing some or all of the data is an act of manipulation that will harm various parties, especially workers. Likewise, companies will experience difficulty in following up on OSH improvement efforts. Therefore, a digitalization-based information system is needed in the form of an application for reporting work accidents, work-

related illnesses and potential hazards in coal mining. The information produced from the reporting system becomes the basis for companies to make appropriate decisions,²¹ and has a positive impact on worker performance.²² The use of mining information technology has a positive impact on increasing safety and productivity.²³

The question is, is the SIKAP application able to prevent the practice of manipulating data on accidents, occupational disease and potential hazards in coal mining? The SIKAP application consists of three user groups, namely workers, companies and government. This is intended so that the reporting carried out does not only involve workers and companies for internal purposes, but also involves the government so that reports of work accidents, work-related illnesses and potential hazards in coal mining can be jointly accounted for by all parties related to accident data, work and potential hazards.

Every worker has the opportunity to report any work accidents, work-related illnesses and potential dangers experienced by themselves or their co-workers through the worker's account in the SIKAP application. Mining workers are able to find out in real-time the number of reports of work accidents, work-related illnesses and potential hazards that are entered into the application, whether reported by themselves or reports from other workers. Apart from that, workers have the opportunity to monitor the progress of each report made, so that it is not just reporting, but there is attention and follow-up from the company.

Reports of work accidents, work-related illnesses and potential hazards carried out by workers will be connected and synchronized with the SIKAP account owned by the company and government. Companies can follow up on reports of potential dangers reported by workers with various efforts or programs to prevent these potential dangers from causing work accidents and work-related diseases. Likewise, reports of work accidents and work-related illnesses reported in the SIKAP application are then followed up by the company in the form of immediate treatment so as not to cause more serious injuries.

The reporting data received by the government is also real-time so that the government can know the total data on work accidents, work-related diseases and potential hazards that have been carried out by workers. Apart from that, the government can also find out about the handling actions that have been taken by the company regarding work accidents and work-related illnesses experienced by workers. This means that there is no difference in the amount of data reported by workers and the data that will be received in company and government accounts. The compatibility of data received between workers, companies and the government will close the space for anyone to carry out data manipulation practices in the future, especially when companies make periodic reports and special reports to the government.

Conclusion

The SIKAP application is an innovative step in the reporting system for work accidents, work-related illnesses and potential hazards to prevent the practice of manipulating report data in coal mining in Indonesia.

References

1. Indonesia, R. Law of the Republic of Indonesia No.13 of 2003 concerning Employment. *Constitution. No.13 of 2003* (2003).
2. Indonesia, P. R. Republic of Indonesia Government Regulation Number 50 of 2012 concerning the Implementation of Occupational Health and Safety Management System. *President of Republik Indonesia* (2012).
3. Heraghty, D., Dekker, S. & Rae, A. Accident report interpretation. *Safety* **4**, 46 (2018).
4. Sneddon, A., Mearns, K. & Flin, R. Stress, fatigue, situation awareness and safety in offshore drilling crews. *Saf. Sci.* (2013) doi:10.1016/j.ssci.2012.05.027.
5. Molinero Ruiz, E., Pitarque, S., Fondevila-McDonald, Y. & Martin-Bustamante, M. How Reliable and Valid is the Coding of the Variables of the European Statistics on Accidents at Work (ESAW)? A Need to Improve Preventive Public Policies. *Saf. Sci.* (2015) doi:10.1016/j.ssci.2015.05.005.
6. Sultan, M. Collaboration Tripartite Elements Accident Reporting and Potential Hazards Management Coal Mining, Indonesia: Systematic Review. *J. Heal. Sci. Med. Res.* **41**, 1–5 (2023).
7. Quanlong Liu et al. Regulatory Regime on Coal Mine Safety in China and Australia: Comparative Analysis and Overall Findings. *Resour. Policy* **74**, (2021).
8. Wang, X., Zhang, C., Deng, J., Su, C. & Gao, Z. Analysis of Factors Influencing Miners Unsafe Behaviors in Intelligent Mines using a Novel Hybrid MCDM Model. *Int. J. Environ. Res. Public Health* **19**, (2022).

9. Weilli Fang et al. Automated text classification of near-misses from safety reports: An improved deep learning approach. *Adv. Eng. Informatics* **44**, (2020).
10. Winkler, M., Perlman, Y., & Westreich, S. Reporting Near-Miss Safety Events: Impacts and Decision-Making Analysis. *Saf. Sci.* **117**, 365–374 (2019).
11. Sultan, M., Setyadi, D. & Ramdan, I. M. Work Accident Reporting in Coal Mining, Indonesia: A Systematic Literature Review. *P.O.S.H Period. Occup. Saf. Heal.* **2**, 51–59 (2023).
12. Qiao, W., Li, X. & Liu, Q. Systemic approaches to incident analysis in coal mines: Comparison of the STAMP, FRAM and “2–4” models. *Resour. Policy* (2019) doi:10.1016/j.resourpol.2019.101453.
13. Kumara et al. Estimation of Human Error Rate in Underground Coal Mines Through Retrospective Analysis of Mining Accident Reports and Some Error Reduction Strategies. *Saf. Sci.* **123**, (2020).
14. Dash, A. K., Bhattacharjee, R. M., Ahmad, A., Sagesh, K. M. R. & Singh, C. S. Mine accident investigation in India: A system approach. *J. Mines, Met. Fuels* (2018).
15. Huang, Y. & Zhou, Q. Mine accident prediction and analysis based on multimedia big data. *Multimed. Tools Appl.* (2019) doi:10.1007/s11042-019-7175-6.
16. Westreich, M. W. Y. P. S. Reporting near-miss safety events: Impacts and decision-making analysis. *Saf. Sci.* **117**, 365–374 (2019).
17. Tetzlaff, Emily J., et al. Safety Culture: A Retrospective Analysis of Occupational Health and Safety Mining Reports. *Saf. Health Work* **12**, 201–208 (2021).
18. Xinfeng et al. The Mediating Role of Psychological Capital Between Perceived Management Commitment and Safety Behavior. *J. Safety Res.* **72**, 29–40 (2020).
19. Stewart, A. G. Mining is bad for health: a Voyage of Discovery. *Environ. Geochem. Health* **42**, 1153–1165 (2020).
20. Yang Xiuyun, Kees Krul, and D. S. Uncovering Coal Mining Accident Coverups: An Alternative Perspective on China’s New Safety Narrative. *Saf. Sci.* **148**, (2022).
21. Nugroho, A. S. *Management Information Systems: A Practitioner's Review of Information Technology*. (Teknosain, 2018).
22. Abualoush et al. The Role of Employees’ Empowerment as an Intermediary Variable Between Knowledge Management and Information Systems on Employees’ Performance. *VINE J. Inf. Knowl. Manag. Syst.* **48**, 217–237 (2018).
23. Yang, L., Birhane, G. E., Zhu, J. & Geng, J. Mining Employees Safety and the Application of Information Technology in Coal Mining: Review. *Front. Public Heal.* **9**, 1–12

(2021).




Figure 1. Display of worker accounts in the SIKAP application




Figure 2. Display of company accounts in the SIKAP application




Figure 3. Display of government accounts in the SIKAP application

**3. Belitung Nursing
Journal
Q3 : SJR 0,3
Indonesia**

[BNJ] Article Review Request



Joko Gunawan, S.Kep. Ners., PhD <editorbnj@gmail.com>
01/11/2022 16:10

To: Risnah

Risnah:

I believe that you would serve as an excellent reviewer of the manuscript, "Health Practice among Muslim Homebound Older Adults Living in the Southern Thai Community: Health Practice among Muslim Homebound Older Adults," which has been submitted to Belitung Nursing Journal. The submission's abstract is inserted below, and I hope that you will consider undertaking this important task for us.

Please log into the journal web site by 2022-11-29 to indicate whether you will undertake the review or not, as well as to access the submission and to record your review and recommendation.

The review itself is due 2022-11-29.

Submission URL: <https://www.belitungraya.org/BRP/index.php/bnj/reviewer/submission?submissionId=2406&reviewId=1190&key=TnPQe3>

If you review this manuscript you will be eligible for a 10% discount on one article processing charge for a manuscript submitted to BNJ. This must be claimed on submission of the manuscript and is available for one year from completion of your review. Only one discount can be claimed on a submitted manuscript and cannot be combined with any other discounts.

Belitung Nursing Journal is member of ReviewerCredits. Completing the peer-review you are invited to will automatically transfer your review data to ReviewerCredits adding it to your certified reviews and assign credits which you may use on ReviewerCredits online store. Register now a free ReviewerCredits account and be ready to earn credits immediately. Find out more [here](#).

Thank you for considering this request.

Joko Gunawan, S.Kep. Ners., PhD
jokogunawan@belitungraya.org

In 2024 Outlook is replacing Windows Mail and Calendar. [Learn more](#)

Try the new Outlook

Dismiss

☐ Try the new Outlook

[BNJ] Article Review Acknowledgement



Joko Gunawan, S.Kep. Ners., PhD <editorbnj@gmail.com>

02/11/2022 14:44

To: Risnah

Dear Risnah,

Thank you for completing the review of the submission, "Health Practice among Muslim Homebound Older Adults Living in the Southern Thai Community: Health Practice among Muslim Homebound Older Adults," for Belitung Nursing Journal. We appreciate your contribution to the quality of the work that we publish.

Please note that you are now eligible for a 15% discount on one article processing charge for a manuscript submitted to Belitung Nursing Journal. This must be claimed on submission of the manuscript and is available for one year from completion of your review. Only one discount can be claimed on a submitted manuscript and cannot be combined with any other discounts.

Belitung Nursing Journal is [member of Reviewer Credits](#). Completing the peer-review you are invited to, will qualify you to the third-party certification provided by this organization and assign you credits which may be used on Reviewer Credits online store. Register now a free Reviewer Credits account and get ready to start. Find out more here [HERE](#).

Best regards,

Joko Gunawan

Belitung Nursing Journal - Belitung Raya Editorial Office



Type here to search



IHSG -1.49%



21:38
30/05/2024

