

## Korespondensi Artikel

Nama Jurnal: **Jurnal Matematika UNAND**

Status Jurnal: TERAKREDITASI dengan Sinta 2 oleh Kementerian Riset, Teknologi, dan Pendidikan Tinggi (RistekDikti) Republik Indonesia

Judul Artikel: *ESSENTIAL PROPERTIES RELATED TO SHORT-TIME FRACTIONAL FOURIER TRANSFORM*

1. Submitted to the journal “Jurnal Matematika UNAND” (Jun 24, 2024)
2. Editor Response (Jun 25, 2024)
3. Editor Decision: Accept Submission without revision (Sep 17, 2024)
4. Copyediting Review Request (Sep 23, 2024)
5. Proofreading Request (Author) (Sep 23, 2024)
6. Copyediting Review Acknowledgement (Oct 2, 2024)
7. Proofreading Acknowledgement (Author) (Oct 2, 2024)
8. Invoice for ID-1249 (Oct 14, 2024)
9. Paper published (Oct 31, 2024)



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

## [JMUA] Submission Acknowledgement

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Mon, Jun 24, 2024 at 7:07 PM

sri Sri Sulasteri:

Thank you for submitting the manuscript, "Essential Properties Related to Short-Time Fractional Fourier Transform" to Jurnal Matematika UNAND. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL:

<https://jmua.fmipa.unand.ac.id/index.php/jmua/author/submission/1249>

Username: sri\_sulasteri

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Dr. Lyra Yulianti  
Jurnal Matematika UNAND

---

Jurnal Matematika UNAND

<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

**[JMUA]**

2 messages

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Tue, Jun 25, 2024 at 5:11 PM

Silahkan dilengkapi di bagian Supplementary dengan

- 1) full artikel dalam format \*.tex, dan
- 2) file gambar (jika ada) dalam format \*.jpg

Terima kasih

---

Jurnal Matematika UNAND

<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)

---

**SRI SULASTERI** <sri.sulasteri@uin-alauddin.ac.id>  
To: Zumardi Rahman <math.andalas.university.1996@gmail.com>

Tue, Jun 25, 2024 at 7:17 PM

Terimakasih atas informasinya. Artikel sudah saya lengkapi dengan file tex dan file cls.  
Terimakasih.

[Quoted text hidden]



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

**[JMUA] Editor Decision**

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Tue, Sep 17, 2024 at 8:23 AM

sri Sri Sulasteri:

We have reached a decision regarding your submission to Jurnal Matematika UNAND, "Essential Properties Related to Short-Time Fractional Fourier Transform".

Our decision is to: Accept Submission

Additionally, please note that it is still necessary to complete the metadata related to your submission through the journal application. Please ensure to complete this information as soon as possible.

Zumardi Rahman  
Direktorat Teknologi Informasi, Universitas Andalas  
[math.andalas.university.1996@gmail.com](mailto:math.andalas.university.1996@gmail.com)

-----  
Reviewer A:

The paper introduced the short-time fractional Fourier transform (STFrFT) and investigated its essential properties. The aspects of novelty are completely elaborated in the manuscript, so the results are technically sound. In my opinion the manuscript could be published in JMUA without any revision.

The paper introduced the short-time fractional Fourier transform (STFrFT) and investigated its essential properties. The aspects of novelty are completely elaborated in the manuscript, so the results are technically sound. In my opinion the manuscript could be published in JMUA without any revision.

-----  
Jurnal Matematika UNAND  
<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

**[JMUA] Copyediting Review Request**

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Mon, Sep 23, 2024 at 10:39 AM

sri Sri Sulasteri:

Your submission "Essential Properties Related to Short-Time Fractional Fourier Transform" for Jurnal Matematika UNAND has been through the first step of copyediting, and is available for you to review by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and click on the File that appears in Step 1.
3. Open the downloaded submission.
4. Review the text, including copyediting proposals and Author Queries.
5. Make any copyediting changes that would further improve the text.
6. When completed, upload the file in Step 2.
7. Click on METADATA to check indexing information for completeness and accuracy.
8. Send the COMPLETE email to the editor and copyeditor.

Submission URL:

<https://jmua.fmipa.unand.ac.id/index.php/jmua/author/submissionEditing/1249>

Username: sri\_sulasteri

This is the last opportunity to make substantial copyediting changes to the submission. The proofreading stage, that follows the preparation of the galleys, is restricted to correcting typographical and layout errors.

If you are unable to undertake this work at this time or have any questions, please contact me. Thank you for your contribution to this journal.

Zumardi Rahman  
Direktorat Teknologi Informasi, Universitas Andalas  
[math.andalas.university.1996@gmail.com](mailto:math.andalas.university.1996@gmail.com)

---

Jurnal Matematika UNAND

<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)

---

**2 attachments**

**317-324-SRI-SULASTERI-1249.pdf**  
271K



**Artikel STFrFT.tex**  
23K



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

**[JMUA] Proofreading Request (Author)**

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Mon, Sep 23, 2024 at 10:41 AM

sri Sri Sulasteri:

Your submission "Essential Properties Related to Short-Time Fractional Fourier Transform" to Jurnal Matematika UNAND now needs to be proofread by following these steps. Additionally, please ensure that you complete the metadata for your journal submission.

1. Click on the Submission URL below.
2. Log into the journal and view PROOFING INSTRUCTIONS
3. Click on VIEW PROOF in Layout and proof the galley in one or more formats used.
4. Enter corrections (typographical and format) in Proofreading Corrections (Please write the corrections in the attached tex file).
5. Save and email corrections to the Layout Editor and Proofreader.
6. Complete the metadata for your submission.
7. Send the COMPLETE email to the editor.

Submission URL:

<https://jmua.fmipa.unand.ac.id/index.php/jmua/author/submissionEditing/1249>

Username: sri\_sulasteri

Zumardi Rahman  
Direktorat Teknologi Informasi, Universitas Andalas  
[math.andalas.university.1996@gmail.com](mailto:math.andalas.university.1996@gmail.com)

---

Jurnal Matematika UNAND  
<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)

---

**2 attachments**

 **Artikel STFrFT.pdf**  
271K

 **Artikel STFrFT.tex**  
23K



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

**[JMUA] Copyediting Review Acknowledgement**

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Wed, Oct 2, 2024 at 3:24 PM

sri Sri Sulasteri:

Thank you for reviewing the copyediting of your manuscript, "Essential Properties Related to Short-Time Fractional Fourier Transform," for Jurnal Matematika UNAND. We look forward to publishing this work.

Zumardi Rahman  
Direktorat Teknologi Informasi, Universitas Andalas  
[math.andalas.university.1996@gmail.com](mailto:math.andalas.university.1996@gmail.com)

---

Jurnal Matematika UNAND  
<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)



SRI SULASTERI &lt;sri.sulasteri@uin-alauddin.ac.id&gt;

---

**[JMUA] Proofreading Acknowledgement (Author)**

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: sri Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Wed, Oct 2, 2024 at 3:26 PM

sri Sri Sulasteri:

Thank you for proofreading the galley for your manuscript, "Essential Properties Related to Short-Time Fractional Fourier Transform," in Jurnal Matematika UNAND. We are looking forward to publishing your work shortly.

If you subscribe to our notification service, you will receive an email of the Table of Contents as soon as it is published. If you have any questions, please contact me.

Zumardi Rahman  
Direktorat Teknologi Informasi, Universitas Andalas  
[math.andalas.university.1996@gmail.com](mailto:math.andalas.university.1996@gmail.com)

---

Jurnal Matematika UNAND  
<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)





SRI SULASTERI <sri.sulasteri@uin-alauddin.ac.id>

---

## [JMUA] Invoice for ID-1249

1 message

---

**Dr. Lyra Yulianti** <math.andalas.university.1996@gmail.com>  
Reply-To: Zumardi Rahman <math.andalas.university.1996@gmail.com>  
To: Sri Sulasteri <sri.sulasteri@uin-alauddin.ac.id>

Mon, Oct 14, 2024 at 6:14 PM

---

Jurnal Matematika UNAND  
<http://jmua.fmipa.unand.ac.id/index.php/jmua/>

Contact me: [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id) (Lyra Yulianti)



---

**EDISI-4-2024-Invoice-9-ID-1249.pdf**  
166K

## Essential Properties Related to Short-Time Fractional Fourier Transform

SRI SULASTERI<sup>a,\*</sup>, NASRULLAH BACHTIAR<sup>b</sup>, WAHYUNI EKASASMITA<sup>c</sup>

<sup>a</sup> *Department of Mathematics Education, Universitas Islam Negeri Alauddin Makassar,*

<sup>b</sup> *Department of Mathematics, Hasanuddin University,*

<sup>c</sup> *Department of Mathematics, Institut Teknologi Bacharuddin Jusuf Habibie.*

*email : sri.sulasteri@uin-alauddin.ac.id, nasrullahmipa.013@gmail.com,*

*wahyuni.ekasasmita@ith.ac.id*

Accepted ..... Revised ..... Published .....

**Abstract.** *We start by defining the short-time fractional Fourier transform in this paper, which is a natural generalization of the fractional Fourier transform. We then investigate its essential properties and explore an uncertainty principle related to this proposed transformation.*

**Keywords:** fractional Fourier transform, short-time fractional Fourier transform, uncertainty principle

### 1. Introduction

It is well known that the fractional Fourier transform (FrFT) is a generalized of the usual Fourier transform (FT), has several applications in optics, signal processing, and quantum mechanics [1,2,3,4,5,6,7]. The fractional Fourier transform depends on a parameter  $\vartheta$  and can be considered as a counterclockwise rotation of the function to arbitrary angles in the spatial frequency domain. Since the FrFT is an extension of the FT, we may transfer some properties of the FT to the FrFT domain like shifting, scaling, modulation, convolution, and inequalities with some changes [8,9,10].

On the other hand, the fractional Fourier domain-frequency contents, which are needed in some applications, are not found by FrFt. This problem can be solved using short-time fractional Fourier transform (STFrFT) [11]. A crucial time–frequency analysis tool is the STFrFT [12]. It has been extensively applied in numerous signal processing domains including speech and acoustics.

The uncertainty principle (UP) is a key concept in signal processing and quantum mechanics. In many changes, one of the well-liked outcomes is likewise the uncertainty principle. A number of works, including [13,14], have recently addressed

\*corresponding author

the uncertainty principles associated with the conventional short-time Fourier transform. The STFrFT, which may be thought of as one rotation of the conventional short-time Fourier transform, was proposed by the authors in [15]. The authors of [16] conclude that there is a connection between the FrFT parameter and the uncertainty principle for the STFrFT by reducing a fundamental inequality. Nevertheless, there are currently no publications that include the crucial characteristics and underlying principles of uncertainty for the short-time fractional Fourier transform (STFrFT).

This paper's remaining sections are arranged as follows. The fundamentals of the fractional Fourier transform (FrFT) are presented in Section 2, along with a summary of several results that are crucial to understanding the conclusions drawn in this work. The definition of the short-time fractional Fourier transform (STFrFT) is presented in Section 3. This section also includes the derivation of certain fundamental properties, such as Moyal's and the inversion formula. The attributes will yield significant benefits for the advancement of our primary findings. We also investigate a principle of uncertainty associated with this suggested transformation. Lastly, we present our conclusion in Section 4.

## 2. Basic Facts on Fractional Fourier Transform

In this part, we mainly review the basic notations on the fractional Fourier transform (FrFT). We start with the well known definition below.

For  $1 \leq p \leq \infty$ , we use the notation  $L^s(\mathbb{R})$  to represent the space of measurable functions on real numbers  $\mathbb{R}$  with norm

$$\|h\|_{L^s(\mathbb{R})} = \left( \int_{\mathbb{R}} |h(x)|^s dt \right)^{1/s} < \infty, \quad 1 \leq s < \infty. \quad (2.1)$$

In particular, when  $s = \infty$  we obtain  $L^\infty(\mathbb{R})$ -norm

$$\|h\|_{L^\infty(\mathbb{R})} = \operatorname{ess\,sup}_{x \in \mathbb{R}} |h(x)|. \quad (2.2)$$

Especially, if  $h \in L^\infty(\mathbb{R})$  is continuous then

$$\|h\|_{L^\infty(\mathbb{R})} = \sup_{x \in \mathbb{R}} |h(x)|. \quad (2.3)$$

The usual inner product of  $L^2(\mathbb{R})$  is then defined as

$$\langle h, \phi \rangle_{L^2(\mathbb{R})} = \int_{\mathbb{R}} h(x) \overline{\phi(x)} dx. \quad (2.4)$$

where  $\overline{\phi(x)}$  is complex conjugate of  $\phi(x)$ .

We remind that the Fourier transform of a function  $h \in L^2(\mathbb{R})$  is defined by

$$\mathcal{F}\{h\}(\psi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} h(x) e^{-ix\psi} dx. \quad (2.5)$$

Definition 2.1. For any function  $h \in L^2(\mathbb{R})$ , the fractional Fourier transform (FrFT) with parameter  $\vartheta$  is described by the formula

$$\mathcal{F}_\vartheta\{h\}(\psi) = \int_{\mathbb{R}} h(x) K_\vartheta(\psi, x) dx, \quad (2.6)$$

where the kernel transform  $K_\vartheta(\psi, x)$  is defined by

$$K_\vartheta(\psi, x) = \begin{cases} C_\vartheta e^{i(|x|^2 + |\psi|^2) \frac{\cot \vartheta}{2} - it \cdot \psi \csc \vartheta}, & \vartheta \neq n\pi \\ \delta(x - \psi), & \vartheta = 2n\pi \\ \delta(x + \psi), & \vartheta = (2n + 1)\pi, n \in \mathbb{Z}, \end{cases} \quad (2.7)$$

for  $\delta$  is Dirac delta function. Here

$$C_\vartheta = \sqrt{\frac{1 - i \cot \vartheta}{2\pi}}, \quad \overline{C_\vartheta} = \sqrt{\frac{1 + i \cot \vartheta}{2\pi}}, \quad (2.8)$$

and it is evident that

$$|C_\vartheta| = \frac{1}{\sqrt{2\pi \sin \vartheta}}. \quad (2.9)$$

It is straightforward to verify that the FrFT kernel fulfills the following basic properties.

$$\overline{K_\vartheta(\psi, x)} = K_{-\vartheta}(\psi, x), \quad (2.10)$$

and

$$\int_{\mathbb{R}} K_\vartheta(\psi, x) \overline{K_\vartheta(\psi', t)} dt = \delta(\psi - \psi'), \quad (2.11)$$

where  $\overline{K_\vartheta(\psi, x)}$  is the complex conjugate of  $K_\vartheta(\psi, x)$ .

Definition 2.2. For any function  $h \in L^2(\mathbb{R})$  with  $\mathcal{F}_\vartheta\{h\} \in L^2(\mathbb{R})$ . The inversion formula of the fractional Fourier transform (FrFT) of the function  $h$  is described as

$$\begin{aligned} h(x) &= \mathcal{F}_\vartheta^{-1}[\mathcal{F}_\vartheta\{h\}](x) \\ &= \int_{\mathbb{R}} \mathcal{F}_\vartheta\{h\}(\psi) \overline{K_\vartheta(\psi, x)} d\psi \\ &= \int_{\mathbb{R}} \mathcal{F}_\vartheta\{h\}(\psi) \overline{C_\vartheta} e^{-i(|x|^2 + |\psi|^2) \frac{\cot \vartheta}{2} + ix \cdot \psi \csc \vartheta} d\psi. \end{aligned} \quad (2.12)$$

Equation (2.12) tell us to recover the original function from its FrFT.

**Lemma 2.3 (FrFT Parseval [17]).** For two functions  $h, \phi \in L^2(\mathbb{R})$ , it holds

$$\langle h, \phi \rangle_{L^2(\mathbb{R})} = \langle \mathcal{F}_\vartheta\{h\}, \mathcal{F}_\vartheta\{\phi\} \rangle_{L^2(\mathbb{R})}, \quad (2.13)$$

and

$$\|h\|_{L^2(\mathbb{R})}^2 = \|\mathcal{F}_\vartheta\{h\}\|_{L^2(\mathbb{R})}^2. \quad (2.14)$$

### 3. Short-Time Fractional Fourier Transform and Its Essential Properties

This part begins by defining the short-time fractional Fourier transform (STFrFT) which is a generalized of the classical fractional Fourier transform. We then collect its important properties which will be needed later.

**Definition 3.1.** For any function  $\phi \in L^\infty(\mathbb{R})$  and  $h \in L^2(\mathbb{R})$ , the short-time fractional Fourier transform (STFrFT) with respect to  $\phi$  is described by

$$\begin{aligned} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) &= C_\vartheta \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} e^{i\left(\left(\frac{x^2+\psi^2}{2}\right) \cot \vartheta - ix\psi \csc \vartheta\right)} dx \\ &= \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} K_\vartheta(x, \psi) dx. \end{aligned} \quad (3.1)$$

For a fixed  $t$ , equation (3.1) may be expressed in the form

$$\mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) = \mathcal{F}_\vartheta\{\check{A}_{f,\bar{\phi}}\}(\psi), \quad (3.2)$$

where

$$\check{A}_{f,\bar{\phi}}(x, t) = h(x) \overline{\phi(x-t)}. \quad (3.3)$$

**Remark 3.2.** It should be observed that when  $\vartheta = \frac{\pi}{2}$ , equation (3.1) becomes

$$\begin{aligned} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \check{A}_{h,\phi}(x, t) e^{-ix\psi} dx \\ &= \mathcal{F}_\vartheta\{\check{A}_{h,\phi}\}(\psi). \end{aligned} \quad (3.4)$$

The fundamental relationship of the STFrFT with classical FT is given by

$$\begin{aligned} C_\vartheta^{-1} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) e^{-i\psi^2 \frac{\cot \vartheta}{2}} &= \int_{\mathbb{R}} \check{A}_{h,\bar{\phi}}^\vartheta(x, t) e^{-ix\psi \csc \vartheta} dx \\ &= \mathcal{F}\left\{\check{A}_{h,\bar{\phi}}^\vartheta(x, t)\right\}(\psi \csc \vartheta), \end{aligned} \quad (3.5)$$

where

$$\check{A}_{h,\phi}^\vartheta(x, t) = h(x) \phi(x-t) e^{\frac{it^2 \cot \vartheta}{2}}. \quad (3.6)$$

Some of the essential properties of the STFrFT are collected in the following theorems.

**Theorem 3.3 (Moyal's formula ).** Let  $h, \phi \in L^2(\mathbb{R})$ , we have

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) \right|^2 d\psi dt = \|f\|_{L^2(\mathbb{R})}^2 \|\phi\|_{L^2(\mathbb{R})}^2. \quad (3.7)$$

**Proof.** In view of Parseval's formula for the FrFT in equation (2.14) and relation

(3.4), we have

$$\begin{aligned}
\int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi &= \int_{\mathbb{R}} \left| \mathcal{F}\{\check{A}_{h, \phi}\}(\psi) \right|^2 d\psi \\
&= \int_{\mathbb{R}} \left| \check{A}_{h, \phi}(\psi) \right|^2 dx \\
&= \int_{\mathbb{R}} \check{A}_{h, \phi}(\psi) \overline{\check{A}_{h, \phi}(\psi)} d\psi \\
&= \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} \overline{h(x)} \overline{\phi(x-t)} dx \\
&= \int_{\mathbb{R}} h(x) \overline{h(x)} \overline{\phi(x-t)} \phi(x-t) dx. \tag{3.8}
\end{aligned}$$

We integrate the both sides of equation (3.8) with respect to  $dt$  and get

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} \int_{\mathbb{R}} h(x) \overline{h(x)} \overline{\phi(x-t)} \phi(x-t) dx dt. \tag{3.9}$$

Performing the change of variables  $x - t = u$ , we infer that

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} \int_{\mathbb{R}} h(x) \overline{h(x)} \overline{\phi(u)} \phi(u) dx du. \tag{3.10}$$

Using Fubini's theorem on the right side of equation (3.10), we further get

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} |h(x)|^2 dx \int_{\mathbb{R}} |\phi(u)|^2 du, \tag{3.11}$$

or,

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \|h\|_{L^2(\mathbb{R})}^2 \|\phi\|_{L^2(\mathbb{R})}^2,$$

which completes the proof.  $\square$

**Theorem 3.4 (Inversion formula).** Let  $h, \phi \in L^2(\mathbb{R})$ , then inversion formula of the STFrFT is given by

$$h(x) = \frac{1}{\phi(0)} \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.12}$$

**Proof.** It follows with relation (3.1) that

$$\begin{aligned}
\mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) &= C_{\vartheta} \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} e^{i\left(\left(\frac{x^2+\psi^2}{2}\right) \cot \vartheta - ix\psi \csc \vartheta\right)} dx \\
&= \int_{\mathbb{R}} \check{A}_{f, \phi}(x, t) K_{\vartheta}(x, \psi) dx, \tag{3.13}
\end{aligned}$$

where  $\check{A}_{f, \phi}(x, t)$  is defined by equation (3.3). Using inversion of FrFT in (2.12), then

$$\check{A}_{f, \phi}(x, t) = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.14}$$

Equation (3.14) above can be rewritten in the form

$$h(x) \overline{\phi(x-t)} = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.15}$$

Choosing  $x = t$  in equation (3.15), we obtain

$$h(x)\overline{\phi(0)} = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \quad (3.16)$$

It means that

$$h(x) = \frac{1}{\overline{\phi(0)}} \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx.$$

Thus, the desired result is obtained.  $\square$

Next, we will build Sharp Housdorff-Young inequality for the short-time fractional Fourier transform (STFrFT), which generalizes Sharp Hausdorff-Young inequality for the FT in StFrFT domains.

**Theorem 3.5.** Let  $p(1 \leq r \leq 2)$  and  $q$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ , then for any  $h, \phi \in L^p(\mathbb{R})$ , we have

$$\left\| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right\|_{L^q(\mathbb{R} \times \mathbb{R})} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q}^{\frac{1}{2q}}} \|h\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})} \quad (3.17)$$

**Proof.** With aid of Sharp Housdorff-Young inequality for the FT, we find that

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{h(x)\}(\psi)|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p dx \right)^{\frac{1}{p}}. \quad (3.18)$$

We replace  $h(x)$  by  $\check{A}_{h,\phi}^{\vartheta}(x, t)$  by (3.6) into both sides of equation (3.18) above, we get

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{\check{A}_{h,\phi}^{\vartheta}\}(\psi)|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |\check{A}_{h,\phi}^{\vartheta}(x, t)|^p dx \right)^{\frac{1}{p}}. \quad (3.19)$$

We setting  $\psi = \psi \csc \vartheta$  in equation (3.19) above, we obtain that

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{\check{A}_{h,\phi}^{\vartheta}\}(\psi \csc \vartheta)|^q d(\psi \csc \vartheta) \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |\check{A}_{h,\phi}^{\vartheta}(x, t)|^p dx \right)^{\frac{1}{p}}. \quad (3.20)$$

Applying (3.5) to left hand side of equation (3.20), we find

$$\left( (2\pi)^{\frac{q}{2}} |\sin \vartheta|^{\frac{q}{2} - 1} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{1}{p}}. \quad (3.21)$$

Hence,

$$\left( \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^q d\psi \right)^{\frac{1}{q}} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q}^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{1}{p}}, \quad (3.22)$$

and

$$\int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^q d\psi \leq \left( \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q}^{\frac{1}{2q}}} \right)^q \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{q}{p}}. \quad (3.23)$$

Integrating both sides equation (3.23) with respect to  $dt$  we get

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta} \{h\}(t, \psi) \right|^q d\psi dt \leq \left( \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \right)^q \left( \int_{\mathbb{R}} \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx dt \right)^{\frac{1}{p}}. \quad (3.24)$$

Equation (3.24) may be expressed as

$$\begin{aligned} \left( \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta} \{h\}(t, \psi) \right|^q d\psi dt \right)^{\frac{1}{q}} &\leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p dx \int_{\mathbb{R}} |\overline{\phi(x-t)}|^p dt \right)^{\frac{1}{p}} \\ &= \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \|f\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}. \end{aligned} \quad (3.25)$$

The above equation is the same as

$$\left\| \mathcal{STF}_{\phi}^{\vartheta} \{h\}(t, \psi) \right\|_{L^q(\mathbb{R} \times \mathbb{R})} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \|h\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}.$$

We obtained the desired result.  $\square$

#### 4. Conclusion

In this paper, we have introduced the short-time fractional Fourier transform (STFrFT) and investigated its essential properties. We have investigated Sharp Housdorff-Young inequality related to this transformation by applying several relations and properties that have been obtained for the proposed transformation.

#### 5. Acknowledgments

The authors are thankful to St. Nurhilmah Busrah for assistance in writing the manuscript.

#### Daftar Pustaka

- [1] Namias, V., 1980, The fractional order Fourier transform and its application to quantum mechanics, *J. Inst. Math. Appl.*, Vol. **25**: 241–265.
- [2] McBride, A.C., Kerr, F.H., 1987, On Namias's fractional Fourier transforms, *IMA J. Appl. Math.*, Vol **39**: 159–175.
- [3] Tao, R., Xin, Y., and Wang, Y., 2007, Double image encryption based on random phase encoding in the fractional Fourier domain, *Opt. Exp.*, Vol. **15**: 16067–16079.
- [4] Tao, R., Lang, J., and Wang, Y., 2008, Optical image encryption based on the multiple-parameter fractional Fourier transform, *Opt. Lett.*, Vol. **33**: 581–583.
- [5] Qi, L., Tao, R., Zhou, S.Y., and Wang, Y., 2004, Detection and parameter estimation of multicomponent LFM signal based on the fractional Fourier transform, *Science in China (Ser. F. Information Sciences)*, Vol. **47**: 184–198.
- [6] Pei, S.C., and Ding, J.J., 2000, Closed-form discrete fractional and affine Fourier transforms, *IEEE Transactions on Signal Processing*, Vol. **48**(5): 1338–1353.



- [7] Durak, L., and Arikan, O., 2003, Short-time Fourier transform: Two fundamental properties and an optimal implementation, *IEEE Transactions on Signal Processing*, Vol. **51**(5): 1231–1242.
- [8] Bahri, M., 2017, Logarithmic uncertainty principle, convolution theorem related to continuous fractional wavelet transform and its properties on a generalized Sobolev space, *International Journal of Wavelets, Multiresolution and Information Processing*, Vol. **15** (5): 1750050.
- [9] Anh, P.K., Castro, L.P., Thao, P.T., 2017, Two new convolutions for the fractional Fourier transform, *Wireless Personal Communications*, Vol. **92**: 623–637.
- [10] Shi, J., Sha, X., Shong, X., and Zhang, N., 2012, Generalized convolution theorem associated with fractional Fourier transform, *Wireless Communication and Mobile Computing*, Vol. **14**: 1340–1351.
- [11] Tao, R., Li, Y. L., Wang, Y., 2009, Short-time fractional Fourier transform and its applications, *IEEE Transactions on Signal Processing*, Vol. **58**(5): 2568–2580.
- [12] Tao, R., Qi L., and Wang, Y. 2004, *Theory and Application of the Fractional Fourier Transform*, Tsinghua University Press, Beijing.
- [13] Cohen, L., 2000, The uncertainty principles of windowed wave functions, *Opt. Communication*, **179**(2000): 221–229.
- [14] Wódkiewicz, K., 1984, Operational approach to phase-space measurements in quantum mechanics, *Phys. Rev. Lett.*, Vol. **52**(13): 1064–1067.
- [15] Stankovic, L., Alieva, T., Bastiaans, M.J., 2003, Time–frequency signal analysis based on the windowed fractional Fourier transform, *Signal Processing*, Vol. **8**: 2459–2468.
- [16] Guanlei, X., Xiaotong, W., Xiaogang, X., 2009, The logarithmic, Heisenberg’s and short-time uncertainty principles associated with fractional Fourier transform, *Signal Processing*, Vol. **89**(3): 339–343.
- [17] Bahri, M., and Karim, S.A.A., 2023, Fractional Fourier transform: main properties and inequalities, *Mathematics*, Vol. **11**(5): 1234.

## ESSENTIAL PROPERTIES RELATED TO SHORT-TIME FRACTIONAL FOURIER TRANSFORM

SRI SULASTERI<sup>a,\*</sup>, NASRULLAH BACHTIAR<sup>b</sup>, WAHYUNI EKASASMITA<sup>c</sup>

<sup>a</sup> *Department of Mathematics Education, Universitas Islam Negeri Alauddin Makassar,*

<sup>b</sup> *Department of Mathematics, Hasanuddin University,*

<sup>c</sup> *Department of Mathematics, Institut Teknologi Bacharuddin Jusuf Habibie,*

*email : sri.sulasteri@uin-alauddin.ac.id, nasrullahmipa.013@gmail.com,*

*wahyuni.ekasasmita@ith.ac.id*

Accepted June 24, 2024    Revised September 17, 2024    Published October 21, 2024

**Abstract.** *We start by defining the short-time fractional Fourier transform in this paper, which is a natural generalization of the fractional Fourier transform. We then investigate its essential properties and explore an uncertainty principle related to this proposed transformation.*

**Keywords:** fractional Fourier transform, short-time fractional Fourier transform, uncertainty principle

### 1. Introduction

It is well known that the fractional Fourier transform (FrFT) is a generalized of the usual Fourier transform (FT), has several applications in optics, signal processing, and quantum mechanics [1,2,3,4,5,6,7]. The fractional Fourier transform depends on a parameter  $\vartheta$  and can be considered as a counterclockwise rotation of the function to arbitrary angles in the spatial frequency domain. Since the FrFT is an extension of the FT, we may transfer some properties of the FT to the FrFT domain like shifting, scaling, modulation, convolution, and inequalities with some changes [8,9,10].

On the other hand, the fractional Fourier domain-frequency contents, which are needed in some applications, are not found by FrFt. This problem can be solved using short-time fractional Fourier transform (STFrFT) [11]. A crucial timefrequency analysis tool is the STFrFT [12]. It has been extensively applied in numerous signal processing domains including speech and acoustics.

The uncertainty principle (UP) is a key concept in signal processing and quantum mechanics. In many changes, one of the well-liked outcomes is likewise the uncertainty principle. A number of works, including [13,14], have recently addressed

\*Corresponding author

the uncertainty principles associated with the conventional short-time Fourier transform. The STFrFT, which may be thought of as one rotation of the conventional short-time Fourier transform, was proposed by the authors in [15]. The authors of [16] conclude that there is a connection between the FrFT parameter and the uncertainty principle for the STFrFT by reducing a fundamental inequality. Nevertheless, there are currently no publications that include the crucial characteristics and underlying principles of uncertainty for the short-time fractional Fourier transform (STFrFT).

This paper's remaining sections are arranged as follows. The fundamentals of the fractional Fourier transform (FrFT) are presented in Section 2, along with a summary of several results that are crucial to understanding the conclusions drawn in this work. The definition of the short-time fractional Fourier transform (STFrFT) is presented in Section 3. This section also includes the derivation of certain fundamental properties, such as Moyal's and the inversion formula. The attributes will yield significant benefits for the advancement of our primary findings. We also investigate a principle of uncertainty associated with this suggested transformation. Lastly, we present our conclusion in Section 4.

## 2. Basic Facts on Fractional Fourier Transform

In this part, we mainly review the basic notations on the fractional Fourier transform (FrFT). We start with the well known definition below.

For  $1 \leq p \leq \infty$ , we use the notation  $L^s(\mathbb{R})$  to represent the space of measurable functions on real numbers  $\mathbb{R}$  with norm:

$$\|h\|_{L^s(\mathbb{R})} = \left( \int_{\mathbb{R}} |h(x)|^s dt \right)^{1/s} < \infty, \quad 1 \leq s < \infty. \quad (2.1)$$

In particular, when  $s = \infty$  we obtain  $L^\infty(\mathbb{R})$ -norm:

$$\|h\|_{L^\infty(\mathbb{R})} = \operatorname{ess\,sup}_{x \in \mathbb{R}} |h(x)|. \quad (2.2)$$

Especially, if  $h \in L^\infty(\mathbb{R})$  is continuous then:

$$\|h\|_{L^\infty(\mathbb{R})} = \sup_{x \in \mathbb{R}} |h(x)|. \quad (2.3)$$

The usual inner product of  $L^2(\mathbb{R})$  is then defined as:

$$\langle h, \phi \rangle_{L^2(\mathbb{R})} = \int_{\mathbb{R}} h(x) \overline{\phi(x)} dx. \quad (2.4)$$

where  $\overline{\phi(x)}$  is complex conjugate of  $\phi(x)$ .

We remind that the Fourier transform of a function  $h \in L^2(\mathbb{R})$  is defined by:

$$\mathcal{F}\{h\}(\psi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} h(\psi) e^{-ix\psi} dx. \quad (2.5)$$

**Definition 2.1.** For any function  $h \in L^2(\mathbb{R})$ , the fractional Fourier transform (FrFT) with parameter  $\vartheta$  is described by the formula

$$\mathcal{F}_\vartheta\{h\}(\psi) = \int_{\mathbb{R}} h(x) K_\vartheta(\psi, x) dx, \quad (2.6)$$

where the kernel transform  $K_{\vartheta}(\psi, x)$  is defined by:

$$K_{\vartheta}(\psi, x) = \begin{cases} C_{\vartheta} e^{i(|x|^2 + |\psi|^2) \frac{\cot \vartheta}{2} - it \cdot \psi \csc \vartheta}, & \vartheta \neq n\pi, \\ \delta(x - \psi), & \vartheta = 2n\pi, \\ \delta(x + \psi), & \vartheta = (2n + 1)\pi, n \in \mathbb{Z}, \end{cases} \quad (2.7)$$

for  $\delta$  is Dirac delta function. Here

$$C_{\vartheta} = \sqrt{\frac{1 - i \cot \vartheta}{2\pi}}, \quad \overline{C_{\vartheta}} = \sqrt{\frac{1 + i \cot \vartheta}{2\pi}}, \quad (2.8)$$

and it is evident that:

$$|C_{\vartheta}| = \frac{1}{\sqrt{2\pi \sin \vartheta}}. \quad (2.9)$$

It is straightforward to verify that the FrFT kernel fulfills the following basic properties.

$$\overline{K_{\vartheta}(\psi, x)} = K_{-\vartheta}(\psi, x), \quad (2.10)$$

and

$$\int_{\mathbb{R}} K_{\vartheta}(\psi, x) \overline{K_{\vartheta}(\psi', t)} dt = \delta(\psi - \psi'), \quad (2.11)$$

where  $\overline{K_{\vartheta}(\psi, x)}$  is the complex conjugate of  $K_{\vartheta}(\psi, x)$ .

**Definition 2.2.** For any function  $h \in L^2(\mathbb{R})$  with  $\mathcal{F}_{\vartheta}\{h\} \in L^2(\mathbb{R})$ . The inversion formula of the fractional Fourier transform (FrFT) of the function  $h$  is described as:

$$\begin{aligned} h(x) &= \mathcal{F}_{\vartheta}^{-1}[\mathcal{F}_{\vartheta}\{h\}](x), \\ &= \int_{\mathbb{R}} \mathcal{F}_{\vartheta}\{h\}(\psi) \overline{K_{\vartheta}(\psi, x)} d\psi, // \\ &= \int_{\mathbb{R}} \mathcal{F}_{\vartheta}\{h\}(\psi) \overline{C_{\vartheta}} e^{-i(|x|^2 + |\psi|^2) \frac{\cot \vartheta}{2} + ix \cdot \psi \csc \vartheta} d\psi. \end{aligned} \quad (2.12)$$

Equation (2.12) tell us to recover the original function from its FrFT.

**Lemma 2.3 (FrFT Parseval [17]).** For two functions  $h, \phi \in L^2(\mathbb{R})$ , it holds

$$\langle h, \phi \rangle_{L^2(\mathbb{R})} = \langle \mathcal{F}_{\vartheta}\{h\}, \mathcal{F}_{\vartheta}\{\phi\} \rangle_{L^2(\mathbb{R})}, \quad (2.13)$$

and

$$\|h\|_{L^2(\mathbb{R})}^2 = \|\mathcal{F}_{\vartheta}\{h\}\|_{L^2(\mathbb{R})}^2. \quad (2.14)$$

### 3. Short-Time Fractional Fourier Transform and Its Essential Properties

This part begins by defining the short-time fractional Fourier transform (STFrFT) which is a generalized of the classical fractional Fourier transform. We then collect its important properties which will be needed later.

**Definition 3.1.** For any function  $\phi \in L^\infty(\mathbb{R})$  and  $h \in L^2(\mathbb{R})$ , the short-time fractional Fourier transform (STFrFT) with respect to  $\phi$  is described by

$$\begin{aligned} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) &= C_\vartheta \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} e^{i\left(\left(\frac{x^2+\psi^2}{2}\right) \cot \vartheta - ix\psi \csc \vartheta\right)} dx \\ &= \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} K_\vartheta(x, \psi) dx. \end{aligned} \quad (3.1)$$

For a fixed  $t$ , equation (3.1) may be expressed in the form

$$\mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) = \mathcal{F}_\vartheta\{\check{A}_{f,\bar{\phi}}\}(\psi), \quad (3.2)$$

where

$$\check{A}_{f,\bar{\phi}}(x, t) = h(x) \overline{\phi(x-t)}. \quad (3.3)$$

**Remark 3.2.** It should be observed that when  $\vartheta = \frac{\pi}{2}$ , equation (3.1) becomes:

$$\begin{aligned} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \check{A}_{h,\phi}(x, t) e^{-ix\psi} dx \\ &= \mathcal{F}\{\check{A}_{h,\phi}\}(\psi). \end{aligned} \quad (3.4)$$

The fundamental relationship of the STFrFT with classical FT is given by:

$$\begin{aligned} C_\vartheta^{-1} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) e^{-i\psi^2 \frac{\cot \vartheta}{2}} &= \int_{\mathbb{R}} \check{A}_{h,\bar{\phi}}^\vartheta(x, t) e^{-ix\psi \csc \vartheta} dx \\ &= \mathcal{F}\left\{\check{A}_{h,\bar{\phi}}^\vartheta(x, t)\right\}(\psi \csc \vartheta), \end{aligned} \quad (3.5)$$

where

$$\check{A}_{h,\phi}^\vartheta(x, t) = h(x) \phi(x-t) e^{\frac{it^2 \cot \vartheta}{2}}. \quad (3.6)$$

Some of the essential properties of the STFrFT are collected in the following theorems.

**Theorem 3.3 (Moyal's formula).** Let  $h, \phi \in L^2(\mathbb{R})$ , we have:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) \right|^2 d\psi dt = \|f\|_{L^2(\mathbb{R})}^2 \|\phi\|_{L^2(\mathbb{R})}^2. \quad (3.7)$$

**Proof.** In view of Parseval's formula for the FrFT in equation (2.14) and relation (3.4), we have:

$$\begin{aligned} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) \right|^2 d\psi &= \int_{\mathbb{R}} \left| \mathcal{F}\{\check{A}_{h,\phi}\}(\psi) \right|^2 d\psi, \\ &= \int_{\mathbb{R}} \left| \check{A}_{h,\phi}(\psi) \right|^2 dx, \\ &= \int_{\mathbb{R}} \check{A}_{h,\bar{\phi}}(\psi) \overline{\check{A}_{h,\bar{\phi}}(\psi)} d\psi, \\ &= \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} h(x) \overline{\phi(x-t)} dx, \\ &= \int_{\mathbb{R}} h(x) \overline{h(x) \phi(x-t)} \phi(x-t) dx. \end{aligned} \quad (3.8)$$

We integrate the both sides of equataion (3.8) with respect to  $dt$  and get:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt, = \int_{\mathbb{R}} \int_{\mathbb{R}} h(x) \overline{h(x)\phi(x-t)} \phi(x-t) dx dt. \quad (3.9)$$

Performing the change of variables  $x - t = u$ , we infer that:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} \int_{\mathbb{R}} h(x) \overline{h(x)\phi(u)} \phi(u) dx du. \quad (3.10)$$

Using Fubini's theorem on the right side of equation (3.10), we further get:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} |h(x)|^2 dx \int_{\mathbb{R}} |\phi(u)|^2 du, \quad (3.11)$$

or,

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \|h\|_{L^2(\mathbb{R})}^2 \|\phi\|_{L^2(\mathbb{R})}^2, \quad (3.12)$$

which completes the proof.  $\square$

**Theorem 3.4 (Inversion formula).** *Let  $h, \phi \in L^2(\mathbb{R})$ , then inversion formula of the STFrFT is given by:*

$$h(x) = \frac{1}{\phi(0)} \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \quad (3.13)$$

**Proof.** It follows with relation (3.1) that:

$$\begin{aligned} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) &= C_{\vartheta} \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} e^{i\left(\left(\frac{x^2+\psi^2}{2}\right) \cot \vartheta - ix\psi \csc \vartheta\right)} dx \\ &= \int_{\mathbb{R}} \check{A}_{f, \overline{\phi}}(x, t) K_{\vartheta}(x, \psi) dx, \end{aligned} \quad (3.14)$$

where  $\check{A}_{f, \overline{\phi}}(x, t)$  is defined by equation (3.3). Using inversion of FrFT in (2.12), then:

$$\check{A}_{f, \overline{\phi}}(x, t) = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \quad (3.15)$$

Equation equation (3.15) above can be rewritten in the form:

$$h(x) \overline{\phi(x-t)} = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \quad (3.16)$$

Choosing  $x = t$  in equation (3.16), we obtain

$$h(x) \overline{\phi(0)} = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \quad (3.17)$$

It means that

$$h(x) = \frac{1}{\phi(0)} \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx.$$

Thus, the desired result is obtained.  $\square$

Next, we will build Sharp Housdorff-Young inequality for the short-time fractional Fourier transform (STFrFT), which generalizes Sharp Hausdorff-Young inequality for the FT in StFrFT domains.

**Theorem 3.5.** *Let  $p(1 \leq r \leq 2)$  and  $q$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ , then for any  $h, \phi \in L^p(\mathbb{R})$ , we have:*

$$\left\| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right\|_{L^q(\mathbb{R} \times \mathbb{R})} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \|h\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}. \quad (3.18)$$

**Proof.** With aid of Sharp Housdorff-Young inequality for the FT, we find that:

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{h(x)\}(\psi)|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p dx \right)^{\frac{1}{p}}. \quad (3.19)$$

We replace  $h(x)$  by  $\check{A}_{h,\phi}^\vartheta(x, t)$  by (3.6) into both sides of equation (3.19) above, we get:

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{\check{A}_{h,\phi}^\vartheta\}(\psi)|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |\check{A}_{h,\phi}^\vartheta(x, t)|^p dx \right)^{\frac{1}{p}}. \quad (3.20)$$

We set  $\psi = \psi \csc \vartheta$  in equation (3.20) above, we obtain that:

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{\check{A}_{h,\phi}^\vartheta\}(\psi \csc \vartheta)|^q d(\psi \csc \vartheta) \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |\check{A}_{h,\phi}^\vartheta(x, t)|^p dx \right)^{\frac{1}{p}}. \quad (3.21)$$

Applying (3.5) to left hand side of equation (3.21), we find:

$$\left( (2\pi)^{\frac{q}{2}} |\sin \vartheta|^{\frac{q}{2}-1} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{1}{p}}. \quad (3.22)$$

Hence,

$$\left( \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi \right)^{\frac{1}{q}} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{1}{p}}, \quad (3.23)$$

and

$$\int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi \leq \left( \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \right)^q \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{q}{p}}. \quad (3.24)$$

Integrating both sides equation (3.24) with respect to  $dt$  we get:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi dt \leq \left( \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \right)^q \left( \int_{\mathbb{R}} \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx dt \right)^{\frac{q}{p}}. \quad (3.25)$$

Equation (3.25) may be expressed as:

$$\begin{aligned} \left( \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta} \{h\}(t, \psi) \right|^q d\psi dt \right)^{\frac{1}{q}} &\leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q}^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p dx \int_{\mathbb{R}} |\overline{\phi(x-t)}|^p dt \right)^{\frac{1}{p}} \\ &= \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q}^{\frac{1}{2q}}} \|f\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}. \end{aligned} \quad (3.26)$$

The above equation is the same as:

$$\left\| \mathcal{STF}_{\phi}^{\vartheta} \{h\}(t, \psi) \right\|_{L^q(\mathbb{R} \times \mathbb{R})} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q}^{\frac{1}{2q}}} \|h\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}.$$

We obtained the desired result.  $\square$

#### 4. Conclusion

In this paper, we have introduced the short-time fractional Fourier transform (STFrFT) and investigated its essential properties. We have investigated Sharp Housdorff-Young inequality related to this transformation by applying several relations and properties that have been obtained for the proposed transformation.

#### 5. Acknowledgments

The authors are thankful to St. Nurhilmah Busrah for assistance in writing the manuscript.

#### Bibliography

- [1] Namias, V., 1980, The fractional order Fourier transform and its application to quantum mechanics, *J. Inst. Math. Appl.*, Vol. **25**: 241–265
- [2] McBride, A.C., Kerr, F.H., 1987, On Namias's fractional Fourier transforms, *IMA J. Appl. Math.*, Vol **39**: 159 – 175
- [3] Tao, R., Xin, Y., and Wang, Y., 2007, Double image encryption based on random phase encoding in the fractional Fourier domain, *Opt. Exp.*, Vol. **15**: 16067 - 16079
- [4] Tao, R., Lang, J., and Wang, Y., 2008, Optical image encryption based on the multiple-parameter fractional Fourier transform, *Opt. Lett.*, Vol. **33**: 581 - 583
- [5] Qi, L., Tao, R., Zhou, S.Y., and Wang, Y., 2004, Detection and parameter estimation of multicomponent LFM signal based on the fractional Fourier transform, *Science in China (Ser. F. Information Sciences)*, Vol. **47**: 184 - 198
- [6] Pei, S.C., and Ding, J.J., 2000, Closed-form discrete fractional and affine Fourier transforms, *IEEE Transactions on Signal Processing*, Vol. **48**(5): 1338 - 1353
- [7] Durak, L., and Arikan, O., 2003, Short-time Fourier transform: Two fundamental properties and an optimal implementation, *IEEE Transactions on Signal Processing*, Vol. **51**(5): 1231 - 1242
- [8] Bahri, M., 2017, Logarithmic uncertainty principle, convolution theorem related to continuous fractional wavelet transform and its properties on a generalized Sobolev space, *International Journal of Wavelets, Multiresolution and Information Processing*, Vol. **15**(5): 1750050



- [9] Anh, P.K., Castro, L.P., Thao, P.T., 2017, Two new convolutions for the fractional Fourier transform, *Wireless Personal Communications*, Vol. **92**: 623 – 637
- [10] Shi, J., Sha, X., Shong, X., and Zhang, N., 2012, Generalized convolution theorem associated with fractional Fourier transform, *Wireless Communication and Mobile Computing*, Vol. **14**: 1340 – 1351
- [11] Tao, R., Li, Y. L., Wang, Y., 2009, Short-time fractional Fourier transform and its applications, *IEEE Transactions on Signal Processing*, Vol. **58**(5): 2568 – 2580
- [12] Tao, R., Qi L., and Wang, Y. 2004, *Theory and Application of the Fractional Fourier Transform*, Tsinghua University Press, Beijing.
- [13] Cohen, L., 2000, The uncertainty principles of windowed wave functions, *Opt. Communication*, **179**(2000): 221 - 229
- [14] Wódkiewicz, K., 1984, Operational approach to phase-space measurements in quantum mechanics, *Phys. Rev. Lett.*, Vol. **52**(13): 1064 - 1067
- [15] Stankovic, L., Alieva, T., Bastiaans, M.J., 2003, Timefrequency signal analysis based on the windowed fractional Fourier transform, *Signal Processing*, Vol. **8**: 2459 - 2468
- [16] Guanlei, X., Xiaotong, W., Xiaogang, X., 2009, The logarithmic, Heisenberg's and short-time uncertainty principles associated with fractional Fourier transform, *Signal Processing*, Vol. **89**(3): 339 – 343
- [17] Bahri, M., and Karim, S.A.A., 2023, Fractional Fourier transform: main properties and inequalities, *Mathematics*, Vol. **11**(5): 1234



## JURNAL MATEMATIKA UNAND

**Published by:** Departemen Matematika dan Sains Data FMIPA Universitas Andalas

**Website:** <http://jmua.fmipa.unand.ac.id>

**Email:** [matematika@sci.unand.ac.id](mailto:matematika@sci.unand.ac.id), [lyra@sci.unand.ac.id](mailto:lyra@sci.unand.ac.id)

National Accredited in SINTA 2 by Direktur Jenderal Pendidikan Tinggi, Riset dan Teknologi Republik Indonesia No 152/E/KPT/2023

**- PAID -**

Received from : SRI SULASTERI, NASRULLAH BACHTIAR, WAHYUNI EKASASMITA  
Affiliation: : UNIVERSITAS ISLAM NEGERI ALAUDDIN MAKASSAR  
Invoice number : 009/JMUA/INV/ED-4/2024  
Invoice date : October 31<sup>st</sup>, 2024  
Article ID : 1249  
In payment : Article publication Vol 13 Issue 4

NO	DESCRIPTION	QUANTITY	PRICE	TOTAL
1	Open access charge of publication fee	1 article	IDR 1.000.000	IDR 1.000.000

Payment info:

**LYRA YULIANTI**

**Bank Syariah Indonesia,**

**Account number 7237558728**

Regards,

A handwritten signature in black ink, appearing to read 'Lyra Yulianti', is written over a horizontal line.

**Lyra Yulianti**  
**Chief Editor**

Library



← Back to Submissions

Submission

Review

Copyediting

Production

Submission Files

Search

▶		3236	<a href="#">redaksi, Artikel STFrFT.pdf</a>	24 June 2024	Article Text
---	---	------	---	--------------------	--------------

Download All Files

Pre-Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

1249 / **SULASTERI et al.** / ESSENTIAL PROPERTIES RELATED TO SHORT-TIME FRACTIONAL FC

Library

Workflow

Publication

Submission

Review

Copyediting

Production

Round 1

Reviewer's Attachments

Search

Jurnal Matematika UNAND



← Back to Submissions

No Files

Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
▶ <a href="#">Editor Decision</a>	sri_sulasteri	-	0	<input type="checkbox"/>
	17-09-2024			
	07:23			

Library



Jurnal Matematika UNAND



← Back to Submissions

Copyediting Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
<a href="#">Copyediting</a>	— 23-09- 2024 09:40	-	0	<input type="checkbox"/>

Copyedited

Search

No Files



Submissions

[My Queue](#)

[Archives](#) 1

Help

Archived Submis

Search

Filters

New Submission

1249

**SULASTERI et al.**  
ESSENTIAL PROPERTIES RELATED TO SHORT-TIME FRACT...

1

Published

[View](#)