



Original Article

## Prevalence of human papillomavirus type 6 and type 11 in pregnant women

Nadyah Haruna <sup>1✉</sup>, Mochammad Hatta <sup>2</sup>, Firdaus Hamid <sup>2</sup>, Andi Rofian Sultan <sup>2</sup>, Safri <sup>2</sup>, Monika Fitria Farid <sup>3</sup>, Esa Lestari <sup>4</sup>

<sup>1</sup> Department of Microbiology, Faculty of Medicine and Health Sciences, UIN Alauddin Makassar, Makassar, South Sulawesi, Indonesia

<sup>2</sup> Department of Microbiology, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

<sup>3</sup> Department of Obstetrics and Gynecology, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

<sup>4</sup> Obstetrics and Gynecology Polyclinic, Tajuddin Chalid Hospital, Makassar, South Sulawesi, Indonesia

### ARTICLE INFORMATION

Received: July 14, 2024

Revised: October 14, 2024

Accepted: October 17, 2024

### KEYWORDS

Human Papillomavirus; Prenatal Care; Pregnancy;

### CORRESPONDENCE

Phone: +62 85255747546

E-mail: nadyaharuna@uin-alauddin.ac.id

### ABSTRACT

**Background:** Pregnancy reduces the effectiveness of cell-mediated immunity, making the body more vulnerable to Human Papilloma Virus (HPV) infection. HPV comes in multiple variants, including non-pro-oncogenic varieties like 6 and 11. Less research has been done on low-risk HPV infections than on high-risk HPV since they are more frequently linked to genital warts and rarely result in cancer. Examples of these infections are types 6 and 11. It is necessary to conduct this study to ascertain whether asymptomatic pregnant women have been exposed to HPV types 6 and 11.

**Purpose:** This study aims to determine the prevalence of HPV types 6 and 11 in pregnant women.

**Methods:** A descriptive cross-sectional survey was carried out in Makassar in May 2023. The study population consisted of pregnant women who underwent antenatal care at Tajuddin Chalid Hospital Makassar during the specified research period. After providing clarification and obtaining consent from eligible mothers, we collected a swab from the vaginal area. A PCR examination was conducted on the 167 maternal swabs that were received. Statistical analyses included frequency distribution using SPSS.

**Results:** According to the study, 121 (72.5%) of the 167 samples tested positive for HPV type 6 DNA, while 46 (27.5%) tested negative. Meanwhile, 67% of 167 patients tested positive for HPV type 11 but were negative for the virus. Only 21 respondents with negative identification were found to be harbouring both low-risk Human Papillomavirus types. In addition, 78 respondents received positive results for HPV types 6 and 11.

**Conclusion:** Pregnant women can be infected with HPV types 6 and 11 even if they are asymptomatic. The high percentage of positive results suggests that routine maternal screening is recommended, especially during pregnancy.

### INTRODUCTION

Human papillomavirus (HPV) is an epitheliotropic virus capable of infecting the skin and mucous membranes,<sup>1-3</sup> and is categorized into high-risk and low-risk types.<sup>4</sup> High-risk types exhibit a heightened propensity for oncogenic transformation and progression to cervical cancer. At the same time, low-risk HPV can lead to the development of warts in both men and women, including genital warts referred to as condyloma acuminata.<sup>5-8</sup> Transmission primarily happens through sexual activity, predominantly vaginal and anal intercourse.<sup>7,9-11</sup> Data from the World

Health Organization (WHO) indicates that HPV infection leading to cervical cancer is the fourth most prevalent malignancy among women.<sup>12</sup> Approximately 90% of the 342,000 cervical cancer fatalities transpire in low- and middle-income nations. The incidence of cervical cancer in Indonesia continues to rise.<sup>13</sup> Data from the Ministry of Health indicates that in 2021, the incidence of suspected cervical cancer reached 5,847 cases, reflecting an increase of 832 cases from the prior year. In 2021, data from the Ministry of Health of the Republic of Indonesia indicated that the incidence of cervical cancer was 23.4 per 100,000 Population, accompanied by an average

<https://doi.org/10.30595/medisains.v23i1.23103>

©(2025) by the Medisains Journal. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at [Attribution-NonCommercial 4.0 International](#).

mortality rate of 13.9 per 100,000 population,<sup>14</sup> while less data about genital warts, especially in pregnancy.

Genital warts, while not lethal, can induce considerable psychological and physical distress in affected individuals. Despite being categorized as low-risk, patients with Condyloma acuminata should receive early notification due to the potential for concurrent infection with numerous HPV serotypes.<sup>7-9,15-17</sup> Condyloma acuminata is the primary manifestation of HPV infection in pregnancy. In this phase, there is a fast proliferation in both the quantity and dimensions of warts, with lesions frequently resolving spontaneously during the postpartum period. Novice clinicians may erroneously classify these disappearing warts as warts associated with pregnancy. The elevation of local estrogen and glycogen concentrations in the female genitalia during pregnancy, along with the accompanying immunological alterations, facilitates the proliferation of HPV.<sup>15,18,19</sup>

Ongoing research and surveillance efforts are crucial in elucidating the complex dynamics of HPV transmission and the differential impact of high-risk and low-risk strains on public health. By understanding the nuances of HPV epidemiology, healthcare providers and policymakers can develop more targeted strategies for prevention, screening, and management, ultimately reducing the burden of HPV-related diseases. The reported incidence of HPV infection is considerably higher in high-risk HPV as individuals typically seek medical attention only when their disease has progressed to a severe phase.

Conditions that are classified as low-risk HPV infections are uncommon. It is not uncommon for many individuals to disregard the importance of taking tests, particularly early examinations. The purpose of this study is to ascertain the prevalence of low-risk HPV strains 6 and 11 in asymptomatic expectant women, as there is a dearth of studies that specifically address the prevalence of low-risk HPV during pregnancy.

## METHOD

### Study Design

This is a descriptive observational study with a cross-sectional design.

### Setting and Respondents

This study was conducted at Tajuddin Chalid Hospital in May 2023. The study population included pregnant women who visited the Antenatal Care unit at the hospital and met the specified eligibility criteria. A total of 167 vaginal swab samples were collected. The inclusion criteria for the study were: no history of sexually transmitted infections, not currently undergoing treatment for HPV infection, and willingness to provide signed informed consent. The exclusion criterion was women who were unwilling to participate in the study. The sampling technique used was non-random.

## Variable, Instrument, and Measurement

The variables examined in this study were HPV type 6 and HPV type 11, which were analyzed using Polymerase Chain Reaction (PCR). The DNA of the target organism for HPV was amplified using specific primers synthesized by Geno Tech Corp (Korea) through PCR. This process involves amplifying a specific portion of the human papillomavirus type 11 genome, which has a length of 528 base pairs. The operation encompasses this area. The amplification in this study utilized the forward primer sequence "agttccgtagatgccaaagggca." Human Papillomavirus type 6 (HPV-6) DNA sequence comprises 757 base pairs. The sequence acgtggcctgtgcgtacagtc corresponds to the forward formula. The order is reversed as follows: agagacgagtcaggceaagtc.<sup>20</sup> The examination was carried out in the Molecular Biology laboratory of Hasanuddin University Teaching Hospital.

## Data Analysis

Descriptive statistics (percentages) were used to determine the participants' prevalence of human papillomavirus infection.

## Ethical Consideration

The research team adhered to ethical standards, with ethical approval granted under No. 426/UN.4.6.4.5.31/PP36/2023 by the Health Research Ethics Committee (KEPK) at UNHAS, Protocol UH232010. Participants provided written informed consent before enrollment, and all personal data was kept confidential, with no identifiable information disclosed in any publications.

## RESULTS

### Characteristics of Participants

The majority of participants were women who had given birth multiple times, were aged between 20 and 30 years, had been pregnant for a period of 12 to 24 weeks, and had never been pregnant before. History of abortion and had completed secondary education. No participants had received previous HPV vaccination (Table 1).

### HPV type 6 and type 11 DNA results distribution

According to the PCR examination results of vaginal swabs from asymptomatic pregnant women, 121 mothers tested positive for HPV type 6, whereas 103 mothers tested positive for HPV type 11 (Table 2). The study included 78 moms with HPV types 6 and 11 DNA, while 21 mothers tested negative for both variations. This section consists of What/How the presented data were produced; no raw data should be present in the article. The produced data are presented in tables or figures with an explanation of the results/findings from the work.

**Table 1.** Characteristics of Respondents Who Undertook Ante-Natal Care (n=167)

Characteristics	Result
<b>Age</b> (years old)	
< 20	10 (5.98%)
20-30	101 (60.47 %)
>30-40	44 (26.34%)
>40	12 (7.18 %)
<b>Balance</b>	
Nulliparous	14 (8.38%)
Primipara	64 (38.23%)
Multipara	73 (43.71%)
Grande Multipara	16 (9.58%)
<b>Gestational Age</b> (in weeks)	
0 to 12	10 (5.98%)
>12 to 24	101 (60.47 %)
>24 to 36	44 (26.34 %)
>36	12 (7.18 %)
<b>Marriage History</b>	
No	0
1x	147 (88.2 %)
>1x	26 (15.57 %)
<b>History of abortion</b>	
No	135 (80.84%)
1x	20 (17.96 %)
>1x	2 (1.197 %)
<b>Level of Education</b>	
Basic	43 (25.75 %)
Middle	100 (59.88%)
High Education	24 (14.37%)
<b>History of HPV Vaccination</b>	
No	167 (100%)
Yes	0

**Table 2.** Proportion of HPV Type-11 and HPV type-6 co-infection

HPV Type 6/11	Positive	Negative	Total
<b>Positive</b>	78 (46.7%)	43 (25.7%)	121 (72.5%)
<b>Negative</b>	25 (15 %)	21 (12.6%)	46 (27.5%)
<b>Total</b>	103 (61.7%)	64 (38.3%)	167 (100%)

## DISCUSSION

This study involved 167 patients who participated to determine the prevalence of HPV types 6 and 11. Findings showed that most respondents were 20-30 years old, with the study participants being pregnant women undergoing ANC check-ups. However, other studies focusing on patients with genital warts have included participants from similar age demographics.<sup>21,22</sup>

This proportion is in line with existing studies showing that low-risk HPV infections often co-occur.<sup>23</sup> Previous epidemiologic studies have shown that HPV types 6 and 11 are frequently recurrent cases.<sup>24</sup> This study is also in line with studies showing that multiple infections are common in patients with genital warts and other manifestations of lesions, mainly associated with HPV types 6 and 11.<sup>25</sup>

The significant reduction in genital wart prevalence seen following vaccination underscores the critical role of vaccination initiatives in diminishing the incidence of these low-risk HPV infections. In this context, our data indicating a 46.7% rate of dual infection underscores the necessity of immunization that encompasses both HPV strains. The results from our table suggest that approximately 75.7% of individuals positive for HPV type 11 were also infected with HPV type 6, demonstrating a significant prevalence of dual infection. This finding aligns with the study, which indicated that infection with a kind of HPV frequently precedes infection with a different type in high-risk populations, particularly among younger individuals exhibiting high sexual activity.<sup>22</sup> Many studies examining the frequency of HPV in pregnant women reveal a higher susceptibility to HPV infection compared to non-pregnant women. A past study found that the total prevalence of HPV in pregnant women was 16.82%, compared with 12.25% in non-pregnant women.<sup>11,32</sup>

This conclusion is based on a thorough analysis of data from studies investigating the prevalence of HPV in pregnant patients. Although our study identified Low-Risk HPV types 6 and 11, another study found High-Risk HPV type 16 in the same sample of pregnant women.<sup>32</sup> Another study showed that HPV levels decrease during pregnancy. Still, high viral load contributes to its persistence. These data support the hypothesis that altered immune responses characterize pregnancy.<sup>31,32</sup> However, this pattern was seen in the first two trimesters, while there was an increase in HPV clearance during the postpartum phase.

Given that HPV clearance rates over the 12 months were comparable between pregnant and non-pregnant women, this increase was sufficient to offset the impact of the first two trimesters. Additionally, studies have shown that pregnant people who test positive for HPV show increased levels of anti-HPV IgA antibodies in the early trimester and early postpartum period. However, the limited production of antibodies at the local level may be mainly due to the suppression of the local immune response.<sup>32</sup>

A comprehensive study,<sup>26</sup> revealed that before the introduction of the HPV vaccine, around 1% of the sexually active population was infected with HPV types 6 and 11. The significant reduction in genital wart prevalence seen following vaccination underscores the critical role of vaccination initiatives in diminishing the incidence of these low-risk HPV infections. In this context, our data indicating a 46.7% rate of dual infection underscores the necessity of immunization that encompasses both HPV strains. The results from our table suggest that approximately 75.7% of individuals positive for HPV type 11 were also infected with HPV type 6, demonstrating a significant prevalence of dual infection.

This finding aligns with the study, which indicated that infection with a kind of HPV frequently precedes infection with a different type in high-risk populations, particularly among younger individuals exhibiting high sexual

activity.<sup>22</sup> Past study,<sup>22</sup> found that in populations lacking vaccination access, the incidence of dual infection exceeded 60% among individuals with genital warts. This finding corroborates our data indicating dual infection with the two HPV types, with nearly half of the total sample exhibiting infection by both types. Several countries that have implemented extensive HPV vaccination programs have reported decreased rates of dual infection, highlighting the significance of vaccination interventions in reducing the prevalence of these infections.

Our data indicated that 67.2% of individuals negative for HPV type 11 tested positive for HPV type 6. This shows that individuals not infected with HPV type 11 may still have a significant likelihood of being infected with HPV type 6, which indicates that HPV type 6 was predominantly identified as a solitary infection in individuals with anogenital warts. HPV type 6 is more frequently linked to external warts and is associated with more prominent lesions, in contrast to HPV type 11, which occurs less often as a solitary infection.

The HPV vaccination, which protects against types 6 and 11, has demonstrated efficacy in decreasing the prevalence of these illnesses. Numerous studies have indicated a significant reduction in the prevalence of genital warts in nations that have adopted immunization programs. Another study conducted in Europe, 23, 28, demonstrated that a quadrivalent vaccine targeting HPV types 6 and 11 decreased infection prevalence by 90% among the vaccinated cohort.

The data indicating a 46.7% prevalence of dual infection with HPV types 6 and 11 underscores the necessity for expanded vaccine coverage, particularly in areas with elevated infection rates. Although most HPV infections resolve spontaneously, some cases can result in significant health complications such as genital warts and other types of cancer. The efficacy of HPV vaccination in reducing the occurrence of vaccine-targeted genotypes has been proven. However, individuals may still be at risk of contracting viral strains not covered by the vaccine.

This study emphasizes the importance of HPV vaccination in reducing HPV infections. These findings underscore the need for targeted educational initiatives to promote HPV vaccination and safer sexual practices, particularly among young populations who may be susceptible to high-risk HPV types. While this research offers a general understanding of the occurrence of low-risk HPV in pregnancy, it has limitations and lacks comprehensive reporting. However, it does pave the way for future studies on the potential risk of high-risk HPV infection in pregnant women. Additionally, it suggests the inclusion of additional variables to gather data that can aid in preventing and managing the impact of infection—human papillomavirus (HPV), particularly in pregnant women.

## CONCLUSIONS AND RECOMMENDATION

Despite immune system modulation during pregnancy, not all pregnant women are infected with low-risk HPV types 6 and 11. This study found that over 50% of asymptomatic pregnant women were exposed to these strains. Regular screenings and HPV vaccinations for women of childbearing age are recommended to reduce infection risk. Future research should focus on larger, longitudinal studies to evaluate the long-term effects of HPV infection during pregnancy on maternal and fetal health.

## REFERENCES

1. Vashisht S, Mishra H, Mishra PK, Ekielski A, Talegaonkar S. Structure, Genome, Infection Cycle and Clinical Manifestations Associated with Human Papillomavirus. *Curr Pharm Biotechnol*. 2019;20(15):1260-1280. doi:10.2174/1389201020666190802115722
2. Bonde J, Bottari F, Iacobone AD, et al. Human Papillomavirus Same Genotype Persistence and Risk: A Systematic Review. *J Low Genit Tract Dis*. 2021;25(1):27-37. doi:10.1097/LGT.0000000000000573
3. Wilting SM, Steenbergen RDM. Molecular events leading to HPV-induced high-grade neoplasia. *Papillomavirus Research*. 2016;2:85-88. doi:10.1016/j.pvr.2016.04.003
4. Trottier H, Mayrand MH, Coutlée F, et al. Human papillomavirus (HPV) perinatal transmission and risk of HPV persistence among children: Design, methods, and preliminary results of the HERITAGE study. *Papillomavirus Research*. 2016;2:145-152. doi:10.1016/j.pvr.2016.07.001
5. Hutter JN, Decker CF. Human papillomavirus infection. *Disease-a-Month*. 2016;62(8):294-300. doi:10.1016/j.disamonth.2016.03.014
6. Bonde J, Bottari F, Iacobone AD, et al. Human Papillomavirus Same Genotype Persistence and Risk: A Systematic Review. *J Low Genit Tract Dis*. 2021;25(1):27-37. doi:10.1097/LGT.0000000000000573
7. Sehna B, Zikan M, Nipcova M, Dusek L, Cibula D, Slama J. The association among cervical, anal, and oral HPV infections in high-risk and low-risk women. *Eur J Obstet Gynecol Reprod Biol X*. 2019;4:100061-4. doi:10.1016/j.eurox.2019.100061
8. Condrat CE, Filip L, Gherghel M, Cretoiu D, Suciu N. Maternal HPV Infection: Effects on Pregnancy Outcome. *Viruses*. 2021;13(12):2455. doi:10.3390/v13122455
9. Zhang J, Zha T, Wang X, He W. Prevalence and genotype distribution of HPV infections among women in Chengdu, China. *Virology*. 2024;21(1):52. doi:10.1186/s12985-024-02317-x
10. Liu P, Xu L, Sun Y, Wang Z. The prevalence and risk of human papillomavirus infection in pregnant women. *Epidemiol Infect*. 2014;142(8):1567-1578. doi:10.1017/S0950268814000636

11. WHO 2021. Sexually Transmitted Diseases Guideline. Accessed October 27, 2022. who.int
12. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394-424. doi:10.3322/caac.21492
13. Kemenkes. Profil Kesehatan Indonesia 2020. 2020.
14. Kaur H, Schmidt-Grimminger D, Chen B, et al. HPV Prevalence and its Association with Perinatal Outcomes among Singleton Mothers: Analysis of Pregnancy Risk Assessment and Monitoring System (PRAMS) Data, 2004-2011. *Current Women s Health Reviews*. 2019;15(2):143-149. doi:10.2174/1573404814666180517085022
15. Saito M, Rajesh A, Innes C, et al. The High-Risk Human Papillomavirus Type Influences the Tissue Microenvironment in Cervical Intraepithelial Neoplasia Grade 2. *Viruses*. 2023;15(9):1953. doi:10.3390/v15091953
16. Mosbah A, Barakat R, Nabel Y, Barakat G. High-risk and low-risk human papillomavirus in association with spontaneous preterm labor: a case-control study in a tertiary center, Egypt. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2018;31(6):720-725. doi:10.1080/14767058.2017.1297403
17. Jain M, Yadav D, Jarouliya U, et al. Epidemiology, Molecular Pathogenesis, Immuno-Pathogenesis, Immune Escape Mechanisms and Vaccine Evaluation for HPV-Associated Carcinogenesis. *Pathogens*. 2023;12(12):1380. doi:10.3390/pathogens12121380
18. Sugai S, Nishijima K, Enomoto T. Management of Condyloma Acuminata in Pregnancy: A Review. *Sex Transm Dis*. 2021;48(6):403-409. doi:10.1097/OLQ.0000000000001322
19. Romero-Pastrana F. Detection and Typing of Human Papilloma Virus by Multiplex PCR with Type-Specific Primers. *ISRN Microbiology*. 2012;2012:1-5. doi:10.5402/2012/186915
20. Aleebrahim F, Vahdat K, Nasirian M. Estimating the Number of Genital Warts Cases in Bushehr City (Southwest Iran) in 2019 Using the Multiplier Method. *Arch Clin Infect Dis*. 2021;15(6). doi:10.5812/archcid.105620
21. Machalek DA, Garland SM, Brotherton JML, et al. Very Low Prevalence of Vaccine Human Papillomavirus Types Among 18- to 35-Year Old Australian Women 9 Years Following Implementation of Vaccination. *J Infect Dis*. 2018;217(10):1590-1600. doi:10.1093/infdis/jiy075
22. Garland SM, Kjaer SK, Muñoz N, et al. Impact and Effectiveness of the Quadrivalent Human Papillomavirus Vaccine: A Systematic Review of 10 Years of Real-world Experience. *Clinical Infectious Diseases*. 2016;63(4):519-527. doi:10.1093/cid/ciw354
23. Fortes HR, von Ranke FM, Escuissato DL, et al. Recurrent respiratory papillomatosis: A state-of-the-art review. *Respir Med*. 2017;126:116-121. doi:10.1016/j.rmed.2017.03.030
24. Hartwig S, Syrjänen S, Dominiak-Felden G, Brotons M, Castellsagué X. Estimation of the epidemiological burden of human papillomavirus-related cancers and non-malignant diseases in men in Europe: a review. *BMC Cancer*. 2012;12(1):30. doi:10.1186/1471-2407-12-30
25. Silverberg M. Condyloma in pregnancy is strongly predictive of Juvenile-Onset recurrent respiratory papillomatosis. *Obstetrics & Gynecology*. 2003;101(4):645-652. doi:10.1016/S0029-7844(02)03081-8
26. Khopkar US, Rajagopalan M, Chauhan AR, et al. Prevalence and Burden Related to Genital Warts in India. *Viral Immunol*. 2018;31(5):346-351. doi:10.1089/vim.2017.0157
27. Kjaer SK, Nygård M, Dillner J, et al. A 12-Year Follow-up on the Long-Term Effectiveness of the Quadrivalent Human Papillomavirus Vaccine in 4 Nordic Countries. *Clinical Infectious Diseases*. 2018;66(3):339-345. doi:10.1093/cid/cix797
28. Doorbar J. The human Papillomavirus twilight zone – Latency, immune control and subclinical infection. *Tumour Virus Research*. 2023;16:200268. doi:10.1016/j.tvr.2023.200268
29. Doorbar J. Host control of human papillomavirus infection and disease. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2018;47:27-41. doi:10.1016/j.bpobgyn.2017.08.001
30. Wu Y, Ye H, Liu J, et al. Prevalence of anemia and sociodemographic characteristics among pregnant and non-pregnant women in southwest China: A longitudinal observational study. *BMC Pregnancy Childbirth*. 2020;20(1) : 535. doi:10.1186/s12884-020-03222-1
31. Værnesbranden MR, Wiik J, Sjøborg K, Staff AC, Carlsen KCL, Haugen G et al. Maternal human papillomavirus infections at mid-pregnancy and delivery in a Scandinavian mother-child cohort study. *International Journal of Infectious Diseases*. 2021;1(108):574-581. Doi:10.1016/j.ijid.2021.05.064
32. Egawa N, Doorbar J. The low-risk papillomaviruses. *Virus Res*. 2017;231:119-127. doi:10.1016/j.virusres.2016.12.017