



Prodi Kemas

Anxiety and Depression Among Female Workers in the Formal And Informal Sectors After the Covid-19 Pandemic

 Fatmawaty Mallapiang

 Skripsi Mahasiswa

 Universitas Islam Negeri Alauddin Makassar

Document Details

Submission ID**trn:oid:::1:3250477778****Submission Date****May 15, 2025, 10:41 AM GMT+7****Download Date****May 15, 2025, 10:46 AM GMT+7****File Name****1913_Mallapiang_e_al_ENG_1.pdf****File Size****156.0 KB****7 Pages****4,591 Words****23,501 Characters**

22% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- Bibliography




Exclusions

- 1 Excluded Source

Match Groups

-  **51 Not Cited or Quoted 22%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 21%  Internet sources
- 8%  Publications
- 3%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 51 Not Cited or Quoted 22%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 21% Internet sources
- 8% Publications
- 3% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	www.socialmedicine.info	12%
2	Internet	ejournal.umm.ac.id	3%
3	Internet	www.researchgate.net	1%
4	Internet	revabeno.emnuvens.com.br	<1%
5	Publication	Masagus M. Ridhwan, Asep Suryahadi, Jahen F. Rezki, Dinda Thalia Andariesta. "T...	<1%
6	Internet	theicph.com	<1%
7	Internet	esdw.eu	<1%
8	Internet	www.coursehero.com	<1%
9	Internet	www.cide.mx	<1%
10	Publication	Rohollah Fallah Madvari, Mohadeseh Bagheshahi, Reyhane Sefidkar, Ali Dehghan...	<1%

11	Internet	chasqui.univalle.edu.co	<1%
12	Internet	digitallibrary.tulane.edu	<1%
13	Internet	p.pdfkul.com	<1%
14	Internet	sifisheriessciences.com	<1%
15	Internet	wiredspace.wits.ac.za	<1%
16	Internet	www.cdc.gov	<1%
17	Internet	www.cedoc.inmujeres.gob.mx	<1%
18	Internet	www.frontiersin.org	<1%
19	Internet	www.mdpi.com	<1%
20	Internet	www.scitepress.org	<1%
21	Publication	"Proceedings of the Andalus International Public Health Conference 2017", BMC P...	<1%



ORIGINAL RESEARCH

Anxiety and Depression Among Female Workers in the Formal And Informal Sectors After the Covid-19 Pandemic

Ansiedad y depresión entre trabajadoras de los sectores formal e informal tras la pandemia de Covid-19 en Macasar, Indonesia

Fatmawaty Mallapiang. Public Health Department, Universitas Islam Negeri Alauddin Makassar, Indonesia.
Email: fatmawaty.mallapiang@uin-alauddin.ac.id, <https://orcid.org/0000-0003-2432-570X>

Nurhayati. Public Health Department, Universitas Islam Negeri Alauddin Makassar, Indonesia.
Email: nurhayatibaharuddin05@gmail.com, <https://orcid.org/0000-0001-9875-5011>

Ranti Ekasari. Public Health Department, Universitas Islam Negeri Alauddin Makassar, Indonesia.
Email: ranti.ekasari@uin-alauddin.ac.id, <https://orcid.org/0000-0003-2914-7308>

Received: October 30, 2024.

Accepted: February 11, 2025.

Conflict of interest: none.

DOI: <https://doi.org/10.71164/socialmedicine.v18i2.2025.1913>

Abstract

After the pandemic, women in both formal and informal sectors found themselves at home, balancing work and domestic responsibilities. In addition to their jobs, they took on the role of primary caregivers, assisting their children with home learning while managing household tasks. This dual burden has led many women to experience psychological challenges, such as anxiety and depression. This study aimed to assess the levels of anxiety and depression among women working in both the formal and informal sectors, and to identify factors potentially related to these mental health issues. The research employed a quantitative, cross-sectional study design, with primary data collected between September 10 and October 10, 2021. The participants were women working in the formal sector (e.g., office workers) and the informal sector (e.g., traders, tailors, etc.) across fourteen sub-districts in Makassar. A total of 100 women were included in the sample, selected through purposive sampling. The results showed that most respondents had normal levels of anxiety and depression. The study found a significant correlation between the length of employment and anxiety levels among formal sector workers, but no significant correlation with depression levels or other factors. This suggests that both anxiety and depression are common among working women, who face the added challenge of balancing family and work responsibilities.

Keywords: Depression; anxiety; female workers

Resumen

Tras la pandemia, las mujeres de los sectores formal e informal se encontraron en casa, compaginando su trabajo con las responsabilidades domésticas. Además de sus empleos, asumieron el papel de cuidadoras principales, apoyando a sus hijos en su escolarización en línea y en casa mientras gestionaban sus demás tareas. Esta doble carga ha llevado a muchas a experimentar problemas psicológicos, como ansiedad y depresión. Este estudio pretende evaluar los niveles de ansiedad y depresión entre las que trabajan tanto en el sector formal como en el informal, para identificar factores potencialmente relacionados con estos problemas de salud mental. La investigación empleó un diseño cuantitativo transversal, con datos primarios recabados entre el 10 de septiembre y el 10 de octubre de 2021. Las participantes trabajaban en el sector formal (por ejemplo, oficinistas) y en el sector informal (por ejemplo, comerciantes, costureras, etc.) en catorce subdistritos de Macasar, Indonesia. Se conformó la muestra mediante un muestreo intencionado, seleccionando un total de 100 mujeres. Los resultados mostraron que la mayoría presentaba niveles normales de ansiedad y depresión, entre las trabajadoras del sector formal con una correlación significativa entre la duración de su empleo y sus niveles de ansiedad, pero ninguna correlación significativa con los niveles de depresión u otros factores. Esto sugiere que tanto la ansiedad como la depresión son comunes entre las mujeres trabajadoras que se enfrentan al reto añadido de equilibrar las responsabilidades familiares y laborales.

Palabras clave: depresión; ansiedad; trabajadoras

Introduction

Female workers face unique biological differences compared to their male counterparts, due to reproductive functions such as menstruation, pregnancy, childbirth, and breastfeeding. In addition to these biological factors, women often bear additional responsibilities, such as childcare and eldercare, particularly when disruptions in facilities, services, and schools occur. These extra pressures, which have intensified during the COVID-19 pandemic, further complicate women's roles as they strive to balance work and caregiving duties.¹

Women make up 75% of the workforce in the health and social care sectors globally, and the percentages are much higher in some occupations like nursing and midwifery.² Healthcare personnel who are women experience the effects of crises to a greater extent. At this time, it is unknown how the physical and mental health of healthcare professionals is being impacted, and how this affects the supply of healthcare professionals, access to healthcare, the caliber of treatment delivered, and decision-making regarding health systems.¹

Female workers are more susceptible to stress because of the high psychological demands and lack of control associated with women's tasks at home and in the family.³ Women experience conflicting responsibilities, so that female workers have a tendency to get stressed more easily than men. Depression is also a potential problem. Depression is the most common non-fatal illness and the fourth most common cause of disability.^{4,5}

According to the previous research, levels of stress, anxiety, and depression were higher in female health workers than in male workers, and they were also higher in single workers than in married workers, and were higher in nurses in comparison to other health workers. These levels are higher for those working in city hospitals, when compared to other kinds of hospitals. In comparison to personnel working in other services, these levels are once again much higher for health workers in pandemic clinics and emergency services, where workers come into direct contact with patients who have COVID-19 symptoms.⁶ Professions with a predominance of women are also paid less and have poorer social standing.²

Based on those previous data and research, this study aims to discover the relation between certain variables to stress and depression in female workers within the formal and informal sector after the Covid-19 pandemic, taking into consideration that female workers often have multiple responsibilities and tend to experience stress and depression more often.

Materials and Methods

Participants

This study was conducted from September 10 until October 10, 2021. It was a cross-sectional study where all variables were collected at once. The study used primary data that was collected with a questionnaire, in which the respondents were female workers.

The respondents were females who work in formal sector (office workers) and informal sector (traders, tailors, and etc), and who work in fourteen sub-districts in Makassar. The total consisted of 100 people. Purposive sampling was used in this study. Inclusion criteria were (1) female workers in both formal and informal sectors; (2) a willingness to be a respondent on this research. Students and those that refused to participate were excluded.

Procedures and Measures

The instrument used was a questionnaire containing statements to be filled out by respondents. Variable stress is measured with an instrument developed by the Depression Anxiety Stress Scale 42 (DASS-42) with a 42-item self-reporting instrument designed by the Psychology Foundation of Australia. There are different questions for reporting anxiety and for depression. Each question had four options with its score.

0 = Did not apply to me at all

1 = Applied to me to some degree, or some of the time

2 = Applied to me to a considerable degree, or a good part of the time

3 = Applied to me very much, or most of the time

After that, all scores from every question were calculated and divided into five categories. The categories for anxiety consist of normal (0-7), mild (8-9), moderate (10-14), severe (15-19), and extremely severe (>20). While the categories for depression consist of normal (0-9), mild (10-13), moderate (14-20), severe (21-27), and extremely severe (>28).

Statistical Analysis

Descriptive analysis was calculated through frequency for univariate analysis and chi-square for bivariate analysis using statistical software. The independent variables were age, marital status, working period, and education. The dependent variable was the level of anxiety and depression experienced post the COVID-19 pandemic.

Ethical Clearance

The study was declared to be ethically appropriate in accordance to 7 (seven) WHO 2011 standards. The number of ethical approval was B.354/KEPK/FKIK/XII/2022.

Result

The study provides a comprehensive analysis of anxiety and depression levels among female workers in the formal and informal sectors in Makassar during the post-COVID-19 pandemic period. It categorizes female workers' mental health status by their respective sectors, reflecting the effects of sector-specific working conditions on psychological well-being. The respondents consist of 50 workers in formal sector and 50 workers in informal sectors.

Based on the result, it was showed that most both formal (58.0%) and informal (64.0%) workers were ≥ 35 years old, meaning they were in the older age group of workers. For formal workers, most of them were married (58.0%), while for informal workers many were not married yet (48.0%). The working period was also found to be different between the two groups, in which most formal workers had already worked above four years (56%), while most informal workers

had worked under or equal to four years (78%)(Table 1).

Looking at psychological conditions in term of anxiety, both formal and informal workers tended to have normal level of anxiety. However, the percentage level of anxiety in the normal category was higher for formal sector workers (66.0%) than for informal sector workers (46.0%). For moderate levels of anxiety, the percentage for informal workers (12.0%) was higher than for formal workers (6.0%).

For the depression variable, both formal and informal workers also reported normal levels. All workers in informal sectors reported normal levels of depression (100.0%). While informal sector workers reported similarly, with 96.0%. Within the formal sectors, 4.0% workers reported mild depression levels (Table 1).

According to bivariate analysis, only working period had significant correlation to anxiety level in formal workers. Based on the age variable, female workers under 35 years old who worked in informal sectors were higher in moderate level of anxiety than those workers in the formal sectors. In line with marital status, married female workers in the informal sector show higher level of anxiety than those who work in formal sectors (Table 2).

Table 1. Frequency Distribution of Independent and Dependent Variables on Formal and Informal Worker in Makassar, 2022

Variables	Formal Worker (n=50)		Informal Worker (n=50)	
	n	%	n	%
Age				
≤ 35 years old	29	58.0	32	64.0
> 35 years old	21	42.0	18	36.0
Marital Status				
Not Married	17	34.0	24	48.0
Married	29	58.0	16	32.0
Divorced	4	8.0	10	20.0
Working period				
≤ 4 years	22	44.0	39	78.0
> 4 years	28	56.0	11	22.0
Education				
Elementary School	0	0	5	10.0
Junior High School	0	0	4	8.0
Senior High School	28	44.0	33	66.0
Bachelor Degree	22	56.0	8	16.0
Anxiety				
Normal	33	66.0	23	46.0
Mild	14	28.0	21	42.0
Moderate	3	6.0	6	12.0
Severe	0	0	0	0
Extremely Severe	0	0	0	0
Depression				
Normal	48	96.0	50	100.0
Mild	2	4.0	0	0
Moderate	0	0	0	0
Severe	0	0	0	0
Extremely Severe	0	0	0	0

FatmawatyMallapiang, Nurhayati, Ranti Ekasari

Table 2. Factors Associated to Anxiety Level on Formal and Informal Worker in Makassar, 2022

Variables	Formal						Informal					
	Normal	Mild	Moderate	Severe	Extremely Severe	P value	Normal	Mild	Moderate	Severe	Extremely Severe	P value
Age						0.215						0.539
≤ 35 years old	22 (75.9%)	6 (20.7%)	1 (3.4%)	0 (0.0%)	0 (0.0%)		14 (43.8%)	13 (40.6%)	5 (15.6%)	0 (0.0%)	0 (0.0%)	
> 35 years old	11 (52.4%)	8 (38.1%)	2 (9.5%)	0 (0.0%)	0 (0.0%)		9 (50.0%)	8 (44.4%)	1 (5.6%)	0 (0.0%)	0 (0.0%)	
Marital Status						0.658						0.792
Not Married	13 (76.5%)	3 (17.6%)	1 (5.9%)	0 (0.0%)	0 (0.0%)		9 (37.5%)	12 (50.0%)	3 (12.5%)	0 (0.0%)	0 (0.0%)	
Married	18 (62.1%)	9 (31.0%)	2 (6.9%)	0 (0.0%)	0 (0.0%)		8 (50.0%)	6 (37.5%)	2 (12.5%)	0 (0.0%)	0 (0.0%)	
Divorced	2 (50.0%)	2 (50.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		6 (60.0%)	3 (30.0%)	1 (10.0%)	0 (0.0%)	0 (0.0%)	
Working period						0.041						0.929
≤ 4 years	18 (81.8%)	4 (18.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		18 (46.2%)	16 (41.0%)	5 (12.8%)	0 (0.0%)	0 (0.0%)	
> 4 years	15 (53.6%)	10 (35.7%)	3 (10.7%)	0 (0.0%)	0 (0.0%)		5 (45.5%)	5 (45.5%)	1 (9.1%)	0 (0.0%)	0 (0.0%)	
Education						0.915						0.385
Elementary School	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		3 (60.0%)	2 (40.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Junior High School	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		2 (50.0%)	2 (50.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Senior High School	18 (64.3%)	8 (28.6%)	2 (7.1%)	0 (0.0%)	0 (0.0%)		15 (45.5%)	12 (36.4%)	6 (18.2%)	0 (0.0%)	0 (0.0%)	
Bachelor Degree	15 (68.3%)	6 (27.3%)	1 (4.5%)	0 (0.0%)	0 (0.0%)		3 (37.5%)	5 (62.5%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	

Table 3. Factors Associated to Depression Level on Formal and Informal Worker in Makassar, 2022

Variables	Formal						Informal					
	Normal	Mild	Moderate	Severe	Extremely Severe	P value	Normal	Mild	Moderate	Severe	Extremely Severe	P value
Age						0.135						-
≤ 35 years old	27 (93.1%)	2 (6.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		32 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
> 35 years old	21 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		18 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Marital Status						0.783						-
Not Married	16 (94.1%)	1 (5.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		24 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Married	28 (96.6%)	1 (3.4%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		16 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Divorced	4 (100.0%)	0 (0.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		10 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Working period						0.066						-
≤ 4 years	20 (90.9%)	2 (9.1%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		39 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
> 4 years	28 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		11 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Education						0.123						-
Elementary School	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		5 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Junior High School	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		4 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Senior High School	26 (92.9%)	2 (1.1%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		33(100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Bachelor Degree	22 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		50 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	

Based on the working period variable, those who have worked under or equal to four years in theinformal sector tend to have higher level of anxiety, on a moderate level, than in the formal sector. In education, female workers who have completed their senior year of high school and work in informal sectors were higher in moderate levels of anxiety, than those within the formal sectors. Overall, however, most formal and informal workers reported normal level of anxiety (Table 2).

In the informal sector, all categories of all variables showed a normal level of depression. In the formal sector, female workers who were below or equal to 35 years old, were not married,

workedless than or equal to four years, and had completed senior high school showed higher level of mild depression, than the other categories (Table 3).

Discussion

Following the pandemic, socioeconomic disparities caused disparities in lifestyles and housing conditions. Low-income individuals' living and working circumstances need to be changed. Through technology and online learning, it is essential to raise public knowledge of social and housing policy. Seminars and workshops in the underprivileged neighborhoods of urban areas can raise awareness.⁸

The informal sector refers to economic activities carried out by individuals, families, or groups based on mutual trust and agreement, without formal legal incorporation. Many women continue to prefer the informal sector as a means of earning a living, as the number of female workers in this sector is significantly higher compared to others. The informal sector offers opportunities for women, especially those with limited education, to engage in work more easily. An example of this is women working as traders.⁹

In contrast, women employed in the formal sector typically work professionally, in roles such as factory workers, technicians, doctors, teachers, or nurses. These women face stress that stems from both their home and work environments. Factors contributing to stress include long working hours, excessive workloads, unclear job roles, anxiety, and fatigue.¹¹

Anxiety is a subjective sensation of unsettling mental tension that occurs as a response to either internal or external insecurities, or to both. Physical and psychological symptoms, such as shaking, sweating, an elevated heart rate, and an inability to concentrate, are typically associated with anxiety.¹²

In this study, both formal and informal workers tended to have normal level of anxiety. However, the percentage level of anxiety in the normal category was higher for formal sector workers than for informal sector workers.

This research suggests that the anxiety experienced by formal and informal sector women workers is still within normal limits, but that the pandemic added additional stressors, and that both formal and informal sector workers sometimes feel anxious about the Covid-19 virus outbreak. Formal sector women workers whose work is directly related to the community are especially afraid of contracting the Covid-19 virus. Meanwhile, women workers in the informal sector (traders and employees) worried about their reduced income due to the Covid-19 virus outbreak, which caused a lack of customers and consumers.

Work From Office (WFO) was founded by workers who were worried about being exposed to the corona virus when news emerged that a new cluster occurred within their office. Among other factors, this condition arises when employees do not comply with the implementation of health protocols.¹³

According to prior research, 26.2% of nurses reported having mild anxiety, while 45.4% of nurses had work experience of more than three years. During the pandemic, nurses with short tenure ran a higher risk of developing anxiety.¹⁴ More seasoned nurses are more accustomed to the potential risks of the job, and can more easily adjust to changing hospital conditions.¹¹

Depression is a mental condition marked by lowered mood, anhedonia, and can include abnormalities in numerous brain networks and areas.^{15,16} Depression can cause low productivity, loss of initiative and apathy at work¹⁷, as well as many health concerns such as obesity. Both formal and informal workers showed normal depression levels, however in the formal sectors, only a small number of workers reported mild depression.

Research suggests that difficulties working conditions, long hours, high job expectations, poor management, and low earnings can contribute to depression. This study is the first to examine variations in working and living conditions according to household income (\$20,000 or less) and job title (administrators vs. staff).¹⁸

Previous research has shown that the pandemic and the adoption of stringent metrics had a significant negative influence on the general population's psychological health and quality of life (QOL). The risk factors for increased anxiety and depressive symptoms were gender, younger age, student status, unemployment, and past psychiatric history. High Precautionary Measures (PM) may be protective against depression.¹⁹

The Covid-19 pandemic has given rise to numerous manifestations of mental health illnesses, including xenophobia, post-traumatic stress disorder, moral panic, overthinking, worry, and

terror. In the post-pandemic era, contextual interpretation of Qur'anic verses relating to mental health can be a normal guideline actualized in daily life.²⁰

Conclusion

Most of the respondents either in formal or informal sectors tended to have a normal level of both anxiety and depression. Working period had a significant correlation with anxiety level, while many other variables were not shown to be significant. No variables had a significant correlation with depression levels. Both anxiety and depression can occur in workers, especially women, often because they have double and conflicting responsibilities at work and at home.

Addressing the psychological well-being of female workers requires cooperation from multiple parties, including employers and family members. Employers can play a vital role in supporting mental health in the workplace, while family members can provide the essential emotional support. Additionally, workers must be equipped to manage and process information about their own mental health. Although the proportion of women workers experiencing moderate levels of anxiety and depression is relatively small, it is important for companies to recognize this as a potential workplace hazard and take steps to support mental health.

Acknowledgement

The authors would like to thank all of the workers who were willing to become respondents within our research.

References

1. Morgan R, Tan HL, Oveisi N, Memmott C, Korzuchowski A, Hawkins K, et al. Women healthcare workers' experiences during COVID-19 and other crises: A scoping review. *Int J Nurs Stud Adv* [Internet]. 2022;4(December 2021):100066. Available from: <https://doi.org/10.1016/j.ijnsa.2022.100066>
2. World Health Organization. Delivered by women, led by men: a gender and equity analysis of the global health and social workforce [Internet]. Human Resources for Health Observer. 2019. 60 p. Available from: <https://www.who.int/hrh/resources/health-observer24/en/>
3. Chela-Alvarez X, Bulilete O, García-Buades ME, Ferrer-Perez VA, Llobera-Canaves J. Perceived factors of stress and its outcomes among hotel housekeepers in the Balearic Islands: A qualitative approach from a gender perspective. *Int J Environ Res Public Health*. 2021;18(1):1–22.
4. Brinda EM, Rajkumar AP, Attermann J, Gerdtham UG, Enemark U, Jacob KS. Health, Social, and Economic Variables Associated with Depression Among Older People in Low and Middle Income Countries: World Health Organization Study on Global AGEing and Adult Health. *Am J Geriatr Psychiatry* [Internet]. 2016;24(12):1196–208. Available from: <https://www.sciencedirect.com/science/article/pii/S1064748116301841>
5. Sriharan A, Ratnapalan S, Tricco AC, Lupea D. Women in healthcare experiencing occupational stress and burnout during COVID-19: A rapid review. *BMJ Open*. 2021;11(4).
6. Tengilimoğlu D, Zekioglu A, Tosun N, Işık O, Tengilimoğlu O. Impacts of COVID-19 pandemic period on depression, anxiety and stress levels of the healthcare employees in Turkey. *Leg Med*. 2021;48(November 2020).
7. Psychology Foundation of Australia. Depression Anxiety Stress Scales - DASS [Internet]. Psychology Foundation of Australia. 2018 [cited 2020 Sep 5]. Available from: <http://www2.psy.unsw.edu.au/dass/>
8. Gür M. Post-pandemic lifestyle changes and their interaction with resident behavior in housing and neighborhoods: Bursa, Turkey [Internet]. Vol. 37, *Journal of Housing and the Built Environment*. Springer Netherlands; 2022. 823–862 p. Available from: <https://doi.org/10.1007/s10901-021-09897-y>
9. Chen M, Harvey J. The Informal Economy in Arab Nations: a comparative perspective. 2017;69. Available from: <https://www.wiego.org/sites/default/files/resources/files/Informal-Economy-Arab-Countries-2017.pdf>
10. Malmström M, Wincent J. Bank lending and financial discrimination from the formal economy: How women entrepreneurs get forced into involuntary bootstrapping. *J Bus Ventur Insights* [Internet]. 2018;10:e00096. Available from: <https://www.sciencedirect.com/science/article/pii/S2>

352673418300568

11. Lulli LG, Giorgi G, Pandolfi C, Foti G, Finstad GL, Arcangeli G, et al. Identifying psychosocial risks and protective measures for workers' mental wellbeing at the time of covid-19: A narrative review. *Sustain*. 2021;13(24).
12. Pardede JA, Simanjuntak GV, Manalu N. Effectiveness of Deep Breath Relaxation and Lavender Aromatherapy Against Preoperative Patient Anxiety. *Res Artic Divers Equal Heal Care*. 2020;17(4):168–73.
13. Muslim M. Stress Management during the Covid-19 Pandemic. *J Manaj Bisnis*. 2020;23(2):192–201.
14. Li R, Chen Y, Lv J, Liu L, Zong S, Li H, et al. Anxiety and related factors in frontline clinical nurses fighting COVID-19 in Wuhan. *Medicine (Baltimore)*. 2020;99(30).
15. Dai L, Zhou H, Xu X, Zuo Z. Brain structural and functional changes in patients with major depressive disorder: A literature review. *PeerJ*. 2019;2019(11):1–17.
16. Shumet S, Azale T, Angaw DA, Tesfaw G, Wondie M, Alemu WG, et al. Help-seeking preferences to informal and formal source of care for depression: A community-based study in northwest ethiopia. *Patient Prefer Adherence*. 2021;15:1505–13.
17. Montoya-Murillo G, Ibarretxe-Bilbao N, Peña J, Ojeda N. The impact of apathy on cognitive performance in the elderly. *Int J Geriatr Psychiatry*. 2019;34(5):657–65.
18. Linnan L, Arandia G, Bateman LA, Vaughn A, Smith N, Ward D. The health and working conditions of women employed in child care. *Int J Environ Res Public Health*. 2017;14(3).
19. Solomou I, Constantinidou F. Prevalence and predictors of anxiety and depression symptoms during the COVID-19 pandemic and compliance with precautionary measures: Age and sex matter. *Int J Environ Res Public Health* [Internet]. 2020;17(14):1–19. Available from: <https://www.mdpi.com/1660-4601/17/14/4924>
20. Nurani S, Pekalongan I, Pahlawan J, Kajen K, Pekalongan K. The Qur ' an and Mental Health in Post-Pandemic Era. 2021;6(2). Available from: <https://ejournal.uinsaizu.ac.id/index.php/maghza/article/view/5711>



Social Medicine

Health For All

ISSN: 1557-7112