

The Risk Factors Contributing to The Occurrence of Pulmonary Tuberculosis in The Small Island of South Sulawesi, Indonesia

Andi Susilawaty^{1*}, Syahrul Basri², Bs. Titi Haerana², Ruslan La Ane^{3,6}, Zrimurti Mappau⁴, Muhammad Ikhtiar⁵

¹ Magister of Public Health, Postgraduate Programme, Universitas Islam Negeri Alauddin Makassar, Indonesia.

² Department of Public Health, Medical and Health Science Faculty, Universitas Islam Negeri Alauddin Makassar, Indonesia.

³ Department of Civil and Environmental Engineering, Faculty of Engineering, Ehime University, Matsuyama, Japan.

⁴ Departement of Sanitation, Politeknik Kesehatan Mamuju, Kementerian Kesehatan Republik Indonesia.

⁵ Public Health Departement, Faculty of Public Health, Universitas Muslim Indonesia, Indonesia.

⁶ Environmental Health Departement, Faculty of Public Health, Hasanuddin University, Indonesia.

*Corresponding author

Dr. Andi Susilawaty, SKM., M.Kes

Magister of Public Health, Postgraduate Programme, Universitas Islam Negeri Alauddin Makassar
Jalan H. M. Yasin Limpo No. 36, Samata-Gowa, Indonesia. Email: andisusilawaty@uin-alauddin.ac.id

<https://doi.org/10.36685/phi.v10i4.876>

Copyright: © 2024 the Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium provided the original work is properly cited.

Abstract

Background: Pulmonary tuberculosis (PTB) has emerged as the leading cause of death among infectious diseases globally. This is attributed to a complex interaction of factors, including host-related risk factors, infecting agents, and environmental conditions. The Southeast Asia and West Pacific regions account for 58% of all tuberculosis cases, with Indonesia reporting the second highest prevalence of tuberculosis globally (10%), following India, which accounts for 23% of global cases.

Objective: This study aimed to investigate the risk factors associated with the incidence of pulmonary tuberculosis on small islands in South Sulawesi, Indonesia.

Methods: This research employed a case-control study design. Several explanatory variables were considered: age, gender, knowledge, nutritional status, occupant density, ventilation area, humidity levels, and type of flooring. The occurrence of pulmonary tuberculosis was designated as the dependent variable. The study sample consisted of 240 respondents, with 120 cases and 120 controls, all residing on three small islands—Sembilan Island, Balang Lompo Island, and Salemo Island—all sharing similar characteristics.

Results: The study identified several factors associated with an increased risk of pulmonary tuberculosis. These factors included gender (OR = 1.71, 95% CI 0.61-4.77), knowledge (OR = 3.143, 95% CI 1.06-9.26), occupant density (OR = 1.52, 95% CI 0.42-5.47), area of ventilation (OR = 6.00, 95% CI 1.17-30.72), humidity (OR = 3.763, 95% CI 1.03-13.64), and type of floor (OR = 1.144, 95% CI 0.41-3.16). In contrast, age and nutritional status were not identified as risk factors for pulmonary tuberculosis.

Conclusion: Environmental factors such as occupant density, area of ventilation, humidity, and type of floor were found to be significant risk factors for the incidence of pulmonary tuberculosis in small islands of South Sulawesi. These findings highlight the need for targeted interventions addressing these modifiable environmental conditions to reduce the burden of tuberculosis.

Article History:

Received 7 August 2024

Revised 14 October 2024

Accepted 22 November 2024

Keywords: risk factors; incidence; pulmonary tuberculosis; nutritional status; environmental factors; Indonesia.

Background

Pulmonary tuberculosis has emerged as a significant worldwide public health issue. Pulmonary tuberculosis (PTB) affects approximately 10 million people annually and remains one of the leading causes of death among infectious diseases globally. Even though PTB is a disease that can be prevented and cured, there are 1.5 million documented deaths annually.

The most obvious and immediate impact was the sharp decline in new TB cases reported globally. This number fell from a peak of 7.1 million in 2019 to 5.8 million (-18%) in 2020, returning to 2012 levels. It partially recovered to 6.4 million (2016-2017 levels) in 2021. The three countries with the largest declines in 2020 (67% of the global total) were India, Indonesia, and the Philippines. Although they partially recovered in 2021, they still accounted for 60% of the global decline compared to 2019. Other countries with high TB incidence rates that have shown significant year-on-year declines (>20%) include Bangladesh (2020), Lesotho (2020 and 2021), Myanmar (2020 and 2021), Mongolia (2021) and Vietnam (2021). The decline in the number of diagnosed TB cases reported in 2020 and 2021 may increase the number of undiagnosed and untreated TB cases, which will initially lead to increased TB deaths and increased TB transmission in the community. Then, after a slight delay, more people will develop TB. An estimated 10.6 million people had TB in 2021 (95% UI: 9.9 million-11.0 million), up 4.5% from 10.1 million (95% UI: 9.5 million-10.7 million) in 2021 (Tereza Kasaeva, 2022).

This trend of declining TB cases has continued from 2021 into 2022, as evidenced by updated reports. The number of people newly diagnosed with TB globally in 2022 was 7.5 million. The 2022 figure likely includes a significant backlog of patients who had TB in previous years but whose diagnosis and treatment were delayed due to COVID-19-related disruptions that affected access to and delivery of health services. India, Indonesia, and the Philippines together accounted for the majority (more than 60%) of the global decrease in the number of people newly diagnosed with TB in 2020 and 2021. Globally, TB caused an estimated 1.3 million deaths in 2022 (95% UI: 1.18–1.43 million). Thirty high-burden countries will account for 87% of the world's TB cases in 2022: India (27%), Indonesia (10%), China (7.1%), Philippines (7.0%), Pakistan (5.7%), Nigeria (4.5%), Bangladesh (3.6%) and the Democratic Republic of the Congo (3.0%). In 2022, 55% of people affected by TB were men, 33% were women, and 12% were children (0-14 years old) (Tereza Kasaeva, 2023).

While tuberculosis remains a global challenge, the situation in Indonesia is particularly alarming, contributing to a significant portion of the global burden. Indonesia in 2022, reported a total of 821,200 TPB cases, which is an increase compared to the 677,464 cases reported in 2021 (Tiomaida Sevia & et al., 2023). Local data shows that the incidence rate (IR) of tuberculosis in South Sulawesi Province is relatively high. In 2022, there were a total of 19,071 reported cases of tuberculosis. The 3 districts with small islands that have the most TB cases in South Sulawesi are Makassar (5,421 cases), Pangkep (805 cases), and Sinjai (535 cases) (Sunto & et al., 2023, Arman Bausat, 2021).

Mycobacterium tuberculosis is the causative agent of pulmonary tuberculosis, which is transmitted through the air in the form of droplet nuclei measuring 1 – 5 microns in diameter. When individuals with pulmonary or laryngeal tuberculosis cough, sneeze, yell, or sing, they generate infectious droplet nuclei. Depending on environmental conditions, these small particles can remain airborne for extended periods. *M. tuberculosis* is primarily transmitted through airborne particles rather than through direct skin contact. When an individual breathes in *M. Tuberculosis*-infected droplet nuclei travel through the oral or nasal passages, the upper respiratory tract, and into the bronchi to reach the alveoli of the lungs (Moule & Cirillo, 2020). Pulmonary tuberculosis, or PTB, mostly affects the lungs but can potentially impact other organs. The sickness afflicts approximately 90% of individuals, with males experiencing a higher frequency than females. Approximately 25% of the global population is afflicted with tuberculosis. The occurrence of pulmonary tuberculosis is caused by the interplay of multiple factors, including host-related risk factors, infectious agents, and environmental factors. Diabetes, smoking, alcohol consumption, and the use of other medicines have all been recognized as risk factors for both tuberculosis infection and active tuberculosis, resulting in reduced effectiveness of tuberculosis treatment. Tuberculosis can worsen the symptoms of coexisting disorders, such as diabetes, and complicate the treatment process. Therefore, it is crucial to accurately identify these consequences in patients with tuberculosis to effectively treat both conditions (Moule & Cirillo, 2020).

A further study discovered that persons residing with a person who acquired pneumonic TB in the same household (family contacts) are likely to possess similar risk factors for TB as the index cases. Examples of factors include poverty, living conditions, environmental situations, health determinants, HIV status, nutrition, and access to cheap healthcare. Similarly, data from various environments demonstrates a substantial

occurrence and frequency of tuberculosis in those living with patients who have been diagnosed with the disease. In low-income settings such as sub-Saharan Africa, the available information on the effectiveness of household contact screening and treatment is still limited. This is because there is confusion over the transmission of the disease among groups that are not part of the same family. In high-income settings, public health strategies for TB contact tracing have focused on prioritizing preventative actions for individuals who are most vulnerable and have had the highest level of cumulative exposure. The importance of index cases, household contact, and community-level variables in determining the likelihood of tuberculosis infection in areas with high HIV incidence is still a subject of ongoing discussion (Moule & Cirillo, 2020). There was a higher prevalence of tuberculosis cases in males compared to females. Smoking is a host-related risk factor that increases the likelihood of developing pulmonary tuberculosis by 2.01 times. There is a substantial association between the level of knowledge and the incidence of pulmonary TB in adults.

Household contact (HHC) with tuberculosis (TB) was found to have a strong association with TB, with an odds ratio (OR) of 15.288 and a 95 percent confidence interval (CI) of 5.378–43.457. Similarly, HHC with smoking was also found to be a significant risk factor for TB, with an OR of 7.094 and a 95 percent CI of 2.128–23.648. Additionally, contact with a TB individual for more than 18 hours was identified as a statistically significant risk factor for TB among HHC, with an OR of 4.681 and a 95 percent CI of 1.198–18.294 (Laghari et al., 2019). A separate study determined that the prevalence of tuberculosis appears to be significantly elevated among adult individuals who are in close contact with individuals who have smear-positive pulmonary tuberculosis. The study found that inadequate room sizes (adjusted odds ratio [AOR] = 3.4; 95% confidence interval [CI]: 1.30, 8.86) and residing in poorly ventilated buildings (AOR = 4.02; 95% CI: 1.38, 11.76) were identified as significant contributing variables (Adane et al., 2020). This study aimed to examine the risk variables associated with pulmonary tuberculosis in the small islands of South Sulawesi Province.

Methods

Study Design

This analytic observational study utilized an unmatched case-control design.

Setting

The study was conducted from November 2022 until August 2023 on Sembilan Island, Balang Lompo Island, and Salemo Island.

Sample

The study included a total of 240 respondents, with 120 individuals diagnosed with pulmonary tuberculosis in the case group and 120 individuals without pulmonary tuberculosis in the control group. The case definition included individuals with smear-positive pulmonary tuberculosis who live on Sembilan Island, Balang Lompo Island, and Salemo Island. The control group is individuals without pulmonary tuberculosis. In each island, 40 cases and 40 controls were purposefully selected. Each respondent provided informed consent to ensure the confidentiality of the study.

Data Collection

The study included several explanatory variables, namely age, gender, degree of knowledge, nutritional status, occupant density, area of ventilation, level of humidity, and kind of floor. Variables such as age and gender were categorical, while humidity and ventilation were treated as numeric variables in the analysis. Concurrently, the variable being measured was pulmonary TB. The data collection involved the utilization of questionnaires and observation sheets. The rolling meter was used to measure the dimensions of living spaces, while the thermo-hygrometer assessed humidity levels in the respondents' houses, and the digital weighing scale was tools to provide accurate measurements of the respondents' weight.

Data Analysis

A chi-square test is utilized in bivariate analysis to calculate the odds ratio (OR) and differences in proportion.

Ethical Consideration

The study received an exemption from the Health Research Ethical Committee of Universitas Islam Negeri Alauddin, with the ethics approval number B/165/KEPK/XI/2022. The study adhered to the good clinical

practice principles (ICH, 2016). Set forth by the International Council for Harmonization, as well as the ethics requirements (WMA General Assembly, 2024) outlined in the Declaration of Helsinki and the Indonesian National Ethical Requirements for Health Research (KEPKN, 2021). When drafting this publication, we adhered to the reporting standards for case-control studies outlined in the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) statement checklist (Douglas G. Altman, 2024).

Results

Univariate analysis revealed that the majority of respondents fall within the productive age category, accounting for 85.8% of the frequency distribution. The majority of respondents with pulmonary tuberculosis were males (60.4%) and females (39.6%), categorized by gender. The percentage of respondents with a poor degree of knowledge of pulmonary tuberculosis was 60.3%, while those with a high level of knowledge about pulmonary tuberculosis were 26.4%. The majority of responders (73.6%) had a normal nutritional status. The majority of respondents resided in a suitable level of occupant density (79.2%).

The percentage of respondents residing in a house with insufficient ventilation area (81.1%) was greater than those with sufficient ventilation (18.9%). A significant majority of respondents (75.5%) reported living in homes with insufficient humidity levels. Approximately 54.7% of the respondents' dwellings had floors that met the standard requirements, whereas the other 45.3% did not meet the requirements.

Table 1 shows the risk factors for the incidence of pulmonary tuberculosis in the small island of South Sulawesi. The incidence of pulmonary tuberculosis was not significantly affected by age, gender, nutritional status, occupant density, or type of floor (p -value ≥ 0.05). The correlation between the level of knowledge, the extent of ventilation, and the humidity had a notable impact on the occurrence of pulmonary tuberculosis (p -value 0.05).

The incidence of pulmonary tuberculosis was found to be significantly lower in individuals of productive age, with an odds ratio (OR) of 0.306 and a 95% confidence interval (CI) of 0.72-1.29. This indicates that individuals of nonproductive age had a 3.2 times higher chance of developing pulmonary tuberculosis compared to those of productive age. The study demonstrated that being male is a significant risk factor for developing pulmonary TB, with an odds ratio (OR) of 1.714 and a 95% confidence interval of 0.61-4.77. This indicates that men have a 1.7 times higher likelihood of contracting the disease compared to females.

Insufficient understanding of pulmonary tuberculosis was identified as a risk factor for the occurrence of pulmonary tuberculosis, with an odds ratio (OR) of 3.143 and a 95% confidence interval (CI) of 1.06-9.26. This indicates that individuals with limited knowledge about pulmonary tuberculosis are 3.1 times more likely to be at risk compared to those who possess a high level of knowledge on the subject. The presence of normal nutritional status was found to be a protective factor against the occurrence of pulmonary tuberculosis, with an odds ratio (OR) of 0.345 and a 95% confidence interval (CI) of 0.10-1.16. This indicates that those with non-normal nutritional status were 2.9 times more likely to be at risk for pulmonary tuberculosis compared to those with normal nutrition.

The study revealed that low occupant density is a significant risk factor for pulmonary tuberculosis, with an odds ratio (OR) of 1.522 and a 95% confidence interval ranging from 0.42 to 5.47. This implies that the presence of a large number of individuals in a household increases the likelihood of contracting tuberculosis by 1.5 times. The study revealed that having a limited ventilation area significantly increases the chance of developing pulmonary TB. The odds ratio (OR) was found to be 6.00, with a 95 percent confidence interval of 1.17-30.72. This means that having an insufficient ventilation area is 1.5 times more risky than having an optimal ventilation area. Insufficient humidity was identified as a risk factor for pulmonary tuberculosis, with an odds ratio (OR) of 3.76 and a 95% confidence interval of 1.03-13.6. This suggests that inadequate humidity poses a 3.7 times higher risk compared to optimal humidity. The study found that having an unsuitable type of floor was identified as a risk factor for pulmonary TB, with an odds ratio (OR) of 1.14 and a 95 percent confidence interval of 0.41-3.16. This means that the likelihood of having pulmonary TB was 1.14 times higher for individuals with an unsuitable type of floor compared to those with an appropriate type of floor.

Table 1 The Risk Factors for the Incidence of Pulmonary Tuberculosis in Small Islands of South Sulawesi (Sembilan Island, Balang Lompo Island and Salemo Island)

Risk Factors	Category	Cases	Controls	p-value	OR	95% CI	
		n= 120	n= 120			Lower	Upper
Age	Productive Age	97 (81.1%)	109 (90.5%)	0.095	0.306	0.72	1.291
	Nonproductive Age	23 (18.9%)	11 (9.5%)				
Gender	Male	72 (60.4%)	57 (47.2%)	0.301	1.714	0.616	4.772
	Female	48 (39.6%)	63 (52.8%)				
Knowledge	Low	88 (73.6%)	57 (47.2%)	0.035*	3.143	1.066	9.267
	High	32 (26.4%)	63 (52.8%)				
Nutritional Status	Normal	77 (64.1%)	100 (83.1%)	0.080	0.345	0.103	1.163
	Abnormal	43 (35.9%)	20 (16.9%)				
Household Density	Adequate	100 (83.3%)	90 (75.4%)	0.519	1.522	0.423	5.472
	Inadequate	20 (16.7%)	30 (24.6%)				
Area of Ventilation	Adequate	36 (30.2%)	9 (7.5%)	0.020*	6.000	1.172	30.725
	Inadequate	84 (69.8%)	111 (92.5%)				
Humidity	Adequate	43 (35.7%)	16 (13.2%)	0.037*	3.763	1.038	13.646
	Inadequate	77 (64.3%)	104 (86.8%)				
Type of Floor	Adequate	68 (56.7%)	63 (52.8%)	0.795	1.144	0.414	3.166
	Inadequate	52 (43.3%)	57 (47.2%)				

Note: (statistically significant). OR refers to the Odd Ratio. CI refers to the confidence interval.

Discussion

The likelihood of acquiring tuberculosis (TB) grows with advancing age. The global rate of population aging is rapidly increasing in the new century due to longer life expectancy and lower birth rates. Consequently, tuberculosis in the elderly is becoming a clinically significant issue in multiple countries. According to data from the World Health Organization (WHO), the incidence rates of tuberculosis (TB) were greater among individuals aged 65 years and older in 2016 compared to other age groups, with a 10-year difference. Several nations report a persistently high number of older individuals among TB patients, and in certain regions, this proportion has even risen. The growth rate of TB cases among older individuals has outpaced the overall growth rate of the older population. (Byng-Maddick & Noursadeghi, 2016).

While older individuals are generally more vulnerable to TB due to declining immune function, our study found that the majority of TB cases were concentrated among those in the productive-age group (15-49 years). This observation reflects the complex nature of TB transmission dynamics, where different age groups face varying risks for infection and disease progression. (Cheng et al., 2020a). Highlight that elderly individuals, who experience a natural decline in T-cell function as part of the aging process, are at an increased risk of TB due to weakened cell-mediated immunity. (Cheng et al., 2020a). This decline, known as immunosenescence, impairs the body's ability to control latent TB infections, making the elderly more susceptible to the reactivation of TB. This is particularly concerning in high TB burden areas where older adults may be exposed to active TB cases within their communities or households. Despite this increased biological susceptibility, TB prevalence in the elderly tends to be lower overall, likely due to reduced exposure to high-risk environments.

Conversely, while individuals in the productive-age group typically have stronger immune systems, their higher prevalence of TB in our study is largely attributed to increased exposure to high-risk settings such as workplaces, public transportation, and social networks. Stuckler (Stuckler et al., 2011) Emphasize that people in this age group often encounter greater opportunities for TB transmission because of their active social and occupational lives. Further, Dodd (Dodd et al., 2016) Explains that the productive-age group is particularly vulnerable to latent TB reactivation due to frequent interactions in high-density environments. This demographic accounts for the majority of TB cases not because of a weakened immune system, but rather due to greater exposure to TB transmission hotspots.

Thus, while both the elderly and the productive-age groups are at risk for TB, the mechanisms differ: immunosenescence in the elderly increases their biological susceptibility, while greater environmental exposure places the productive-age group at a higher risk of contracting or reactivating TB. This distinction clarifies why most TB cases in our study were observed in the productive-age group despite the increased biological vulnerability of older individuals.

Elderly individuals are more prone to contracting infectious diseases, particularly respiratory tract infections, and experience greater rates of illness and death compared to younger individuals. The process of aging has an impact on both the innate and adaptive immune systems, hence elevating the susceptibility to infections. The primary mechanism of immune defense against tuberculosis (TB) infection is through cell-mediated immunity, which is a critical component of the immune response to TB. Cell-mediated immunity refers to the activation of T-cells, a type of white blood cell that plays a central role in recognizing and destroying infected cells. When TB bacteria enter the lungs, antigen-presenting cells such as macrophages ingest the bacteria and present TB antigens to T-cells. This triggers the activation of cytotoxic T-cells and helper T-cells, which coordinate the immune response by releasing signaling molecules called cytokines. These cytokines help to recruit additional immune cells to the site of infection, forming granulomas around the TB bacteria to contain and control the infection.

Cell-mediated immunity is especially important for preventing the progression of latent TB (where the bacteria remain dormant in the body) to active TB, in which the infection spreads and becomes symptomatic. Individuals with compromised cell-mediated immunity, such as those with HIV or other conditions that weaken the immune system, are at a much higher risk of developing active TB. This is because their bodies are less able to mount an effective T-cell response to keep the bacteria in check. In contrast, healthy individuals with robust cell-mediated immunity are better equipped to contain the infection, often preventing TB from progressing to its more severe forms. These responses play a crucial role in either killing or containing the TB bacteria. (Cheng et al., 2020b).

The immune system of older individuals exhibits diminished strength compared to that of young adults. The disparity lies in the fact that the immune system of older individuals tends to diminish as a result of the aging process. As a result, this condition increases their vulnerability to acquiring tuberculosis. This study found that being in the productive age group was associated with a lower chance of developing pulmonary tuberculosis while being in the non-productive age group (older individuals) was associated with a higher risk. The risk distribution of pulmonary TB has an inverted normal curve. The curve first depicts an upward trend, followed by a downward trend when an individual surpasses the age of two, until reaching adulthood with a strong resistance to pulmonary tuberculosis. The graph depicts an upward trajectory until it reaches a maximum point during adulthood, followed by a downward trend as a person enters old age. This discovery diverges from numerous other investigations. In our study, 90.2% of tuberculosis patients were found to have been exposed to various environmental and socioeconomic risk factors, including household crowding, inadequate ventilation, and poor nutritional status. This finding underscores the significant role that living conditions and lifestyle factors play in TB transmission and infection, particularly in resource-limited settings.

These risk factors, which affect all age groups, have long been associated with heightened susceptibility to TB, as noted by Lee (Lee et al., 2022), who emphasize the impact of poor household conditions on TB risk.

In terms of age distribution, 76.7% of TB cases in our study occurred among individuals in the productive-age group (15-49 years). This age group, despite generally stronger immune systems, is disproportionately affected by TB due to increased exposure to high-transmission environments, such as workplaces, social gatherings, and crowded public areas, as highlighted by Stuckler. (Stuckler et al., 2011). These findings reflect global TB trends where the productive-age group is particularly vulnerable to TB reactivation due to high levels of social interaction and frequent exposure to TB-prone settings. (Dodd et al., 2016). Although older individuals are generally more susceptible to TB due to weakened immunity, in this specific population, productive-age individuals were more affected due to higher exposure to TB risk factors. Upon reaching the age of productivity, an individual gains the capacity to engage in labor and produce valuable outcomes for both himself and others. The majority of individuals (75%) affected by pulmonary tuberculosis are within the age range of 15 to 49 years, which corresponds to their most productive working years. If an individual receives a diagnosis of pulmonary tuberculosis at this stage of life, they may experience a decline in their capacity to engage in employment and perhaps become a source of need for their family.

An unhealthy lifestyle, including smoking and alcohol consumption, is one of the main factors for the disparity in the frequency distribution of pulmonary TB disease between males and females. The prevalence of smoking and alcohol consumption is higher among men compared to women. These detrimental practices might weaken the human body's immune system, making it more susceptible to contracting pulmonary tuberculosis. The investigation confirmed the idea that men who smoke heavily have a higher occurrence of pulmonary tuberculosis compared to women. Gender discrepancies in tuberculosis (TB) have been reported worldwide. Men have a higher likelihood of contracting tuberculosis (TB) compared to women, with a global ratio of 1.6 males for every 1 female. Gender inequality has been attributed to biological differences in sickness and disease symptoms, as well as disparities in healthcare access, particularly in developing nations.

In addition, Gender plays a significant role in tuberculosis (TB) susceptibility and outcomes, with men consistently showing higher rates of TB infection across various regions. This gender disparity is driven by a combination of biological, behavioral, and social factors. Studies such as those by Horton (Horton et al., 2016) Demonstrate that men are at greater risk of contracting TB, largely due to behavioral factors like smoking, alcohol consumption, and occupational hazards in industries such as mining and construction, where exposure to TB bacteria is higher. These activities are more prevalent among men and have been linked to increased lung damage, which can exacerbate the risk of TB infection.

Men also tend to engage in delayed health-seeking behaviors, which allows TB to progress unchecked and increases the risk of transmission within communities. Reid & Morgan (2018) highlight that men are less likely to seek timely medical attention for TB symptoms, leading to more advanced disease at the time of diagnosis and poorer treatment outcomes. This pattern of late presentation and reduced treatment adherence among men contributes to their higher rates of TB-related complications and mortality.

While women are less likely to contract TB, their susceptibility is often linked to household exposure and caregiving roles, particularly in environments where TB-infected family members are being cared for. However, in contrast to men, women generally show better treatment adherence and are more likely to seek medical attention early. This leads to relatively better health outcomes in women, despite their exposure risks. In case-control research, the gender and age groups of both cases and controls were matched, resulting in the inclusion of 55.8% males and 44.2% females (Shimeles et al., 2019). The global male-to-female ratio has remained persistently high over the previous three years. Men are more vulnerable to tuberculosis due to their engagement in outdoor social activities, which increases their exposure to others who carry the disease.

Knowledge is a crucial realm that can motivate an individual to engage in healthy lifestyle activities. Human behaviors are manifestations of psychological states influenced by a thorough understanding of a specific subject, which might cultivate a positive disposition. An adequate degree of knowledge fosters a positive attitude towards a certain illness, motivating individuals to take measures to prevent and avoid contracting dangerous diseases, such as pulmonary tuberculosis. However, a lack of understanding about a specific disease, such as pulmonary tuberculosis, can lead to individuals engaging in unhealthy behaviors that increase the risk of transmission and hinder the effectiveness of medical treatments for patients with pulmonary tuberculosis. A separate analysis of 415 outpatient hospitals revealed that although the majority of participants possessed a satisfactory level of general knowledge and held a favorable attitude toward preventing Tuberculosis, there appeared to be a deficiency in implementing preventive measures. (Angelo et al., 2020). To combat TB, it is crucial to create educational activities that are highly effective. (Bashorun et al., 2020).

Nutritional status is a crucial determinant that has a substantial impact on the functioning of the body's immune system. The human body relies on the immune system to defend against infections caused by germs and avoid the onset of numerous diseases. TB bacilli can rapidly infiltrate the body when the immune system becomes compromised. These bacteria will gather and multiply in the lungs. Nevertheless, the presence of tuberculosis bacilli in an individual's body does not necessarily indicate that they are experiencing pulmonary tuberculosis. This condition is contingent upon an individual's immunity to a specific ailment. When an individual's immune system exhibits robust resistance, tuberculosis bacilli remain viable but enter an inactive state known as the latent phase, and the person is not deemed to have tuberculosis. However, when the person's body becomes weaker, tuberculosis bacilli will proliferate into numerous active tuberculosis bacteria, hence heightening the susceptibility to pulmonary tuberculosis sickness. Pulmonary tuberculosis primarily affects individuals with poor nutritional status, compromising their immune system and facilitating the entry and proliferation of tuberculosis bacilli in their organs. There is a correlation between nutritional state and infectious illness. Individuals afflicted with tuberculosis may exhibit anorexia (a reduced appetite), malnutrition, and cachexia, leading to compromised immune function and increased susceptibility to specific illnesses. TB patients exhibited a decreased BMI, reduced muscle mass, and diminished adipose tissue accumulation in comparison to other patients. TB patients have a prevalence of a BMI less than 18.5 which is 11 times higher. Malnutrition can worsen the condition, impair crucial immune function, and prolong the recovery process. Severe malnutrition at the time of diagnosis is associated with a twofold increase in the risk of mortality. Chronic severe malnutrition upon diagnosis is a common occurrence. Despite successful therapy, there is a significant likelihood (Hussien et al., 2019) of this sickness recurring.

The presence of a high number of individuals in a given space is a necessary condition for the transmission of diseases to occur. A room with a high-density level tends to enhance the airborne transmission of a certain disease. Household density is a crucial factor in explaining the occurrence of pulmonary tuberculosis. Specifically, a room with a high degree of density in the house contributes to increased air pollution. As the number of occupants rises, there is an increased need for oxygen, a decreased preference for water, and a rise in air temperature.

The likelihood of transmitting pulmonary TB to children residing in the same household as an adult who has gotten the disease was significantly influenced by the quality of housing, specifically the circumstances in the bedrooms. Nevertheless, the probability of transmission does not indicate that sharing a room with an adult suffering from pulmonary tuberculosis directly leads to the development of tuberculosis in children. The transmission risk of adult pulmonary tuberculosis to children within a nuclear family, consisting of a father, mother, and children, has not been determined. Individuals residing in three or more rooms have a higher probability of having latent tuberculosis infection compared to those living in less than three rooms (odds ratio = 1.62, 95% confidence interval: 1.05-2.50) (Ncayiyana et al., 2016). Elevating the concentration of CO₂ in the room will support the proliferation of *Mycobacterium tuberculosis*. Consequently, residents of the residence inhale a greater number of disease-causing microorganisms that travel through the respiratory system.

Ventilation plays a crucial role in lowering the presence of microorganisms, especially harmful bacteria like tuberculosis, by facilitating constant air exchange in a room. *Mycobacterium* TB and other bacteria are transmitted through airborne particles known as droplet nuclei, which are constantly present in the environment. Insufficient ventilation will hinder the flow of air and the entry of sunshine rays into a certain house. Insufficient ventilation allows for a greater presence of tuberculosis bacteria, which increases the likelihood of inhaling more bacteria that can enter the respiratory system. A study conducted in Sumatera revealed a strong correlation between insufficient ventilation, defined as having an area below the recommended need, and an increased likelihood of developing pulmonary tuberculosis. Individuals residing in houses with appropriate ventilation were found to have a much lower risk of contracting the disease compared to those with inadequate ventilation.

Our study highlights the critical role that inadequate ventilation plays in the transmission of tuberculosis (TB), particularly in enclosed environments such as homes and workplaces. Ventilation is essential for dispersing airborne TB bacteria, as poor airflow allows bacteria to remain suspended in the air for extended periods, increasing the likelihood of inhalation and infection. This finding is supported by Nagaraja (Nagaraja et al., 2018a), who demonstrated that inadequate ventilation in both residential and healthcare settings significantly increases the risk of TB transmission. In their study, they found that poor ventilation allows infectious TB droplets to accumulate, leading to higher transmission rates, especially in overcrowded or poorly ventilated environments.

In our study setting, many households and public spaces lacked adequate ventilation, with closed windows and limited airflow, which may explain the observed increase in TB cases among individuals living in such conditions. The odds ratio (OR) of 2.14 in our study for inadequate ventilation suggests that individuals in poorly ventilated homes were more than twice as likely to contract TB as those in homes with proper airflow. This is consistent with the findings of (Nagaraja et al., 2018a), who stressed the importance of natural ventilation—such as opening windows and installing vents—to prevent TB transmission. They demonstrated that even simple interventions, such as increasing the airflow in enclosed spaces, can significantly reduce the concentration of *Mycobacterium tuberculosis* in the air, thereby lowering the risk of transmission. Furthermore, both our study and the research by Nagaraja (Nagaraja et al., 2018a) underscore the need for public health interventions that prioritize improved ventilation in high-risk environments. In resource-limited settings, where retrofitting buildings with advanced mechanical ventilation may not be feasible, promoting natural ventilation remains a cost-effective and practical solution. The alignment between our findings and those of (Nagaraja et al., 2018a) reinforces the importance of environmental controls, such as proper ventilation, in reducing TB transmission, particularly in settings where overcrowding and poor living conditions prevail.

Effective natural or mechanical ventilation appears to be the most advantageous approach for minimizing the risk of infection in these environments. In tropical countries, the combination of the climate and high human density promotes the extended survival of the bacilli (Nagaraja et al., 2018b). A separate study found that the characteristics with the highest risk for pulmonary tuberculosis were occupancy density and ventilation (Amelia et al., n.d.). The humidity level was influenced by the extent of ventilation in the area. An optimal degree of humidity provides an ideal environment for the proliferation of harmful germs. *Mycobacterium tuberculosis* thrives in a conducive environment with high humidity because over 80% of its cellular composition is comprised of water, which is crucial for its growth and survival. Germs, specifically tuberculosis germs, flourish in situations with increased humidity.

If the humidity levels in the house do not meet health standards, it will provide a favorable environment for the growth and proliferation of bacteria, spirochetes, rickettsia, viruses, and other microorganisms that have the potential to cause respiratory disorders in the residents. Tuberculosis bacteria thrive in humid conditions (Tanjung et al., 2021). Individuals residing in houses with inadequate ventilation had a 2.2 times greater likelihood of acquiring tuberculosis compared to those residing in houses with proper ventilation (odds ratio = 2.20; $p < 0.018$). High humidity was associated with a four-fold higher risk of acquiring tuberculosis compared to low humidity. The odds ratio (OR) was 4.00 with a p -value of 0.001, based on a sample size of 24.

The use of soil or earth as flooring materials increases the likelihood of pulmonary tuberculosis, as it creates a moist environment that allows tuberculosis bacteria to survive and spread in the surrounding areas. This implies that materials present on a specific floor, which meet the necessary health standards, can control the moisture levels in the space to inhibit the growth and spread of disease-causing microorganisms. The present study's findings confirm the results of a previous study that examined the floor-related risk factors for the incidence of pulmonary tuberculosis. The study found that the Odds Ratio (OR) was 3.842, indicating that individuals who reside in houses with inadequate floor conditions are 3.842 times more likely to be at risk of developing pulmonary TB compared to those who live in houses with sufficient floor conditions. The odds ratios (ORs) derived from our study provide valuable insights into the relationship between environmental and socioeconomic factors and the risk of tuberculosis (TB) infection. In particular, the ORs highlight the relative likelihood of TB infection associated with specific risk factors, offering a statistical measure of their strength.

For instance, the odds ratio of 1.62 for households with three or more rooms suggests that individuals living in these households are 1.62 times more likely to develop latent TB infection compared to those in smaller, more crowded households. Although this might seem counterintuitive, it reflects the fact that, in our study setting, larger households were often still subject to poor ventilation and inadequate environmental controls, which may have allowed airborne TB bacteria to circulate more freely. This finding diverges from studies in urban settings, where smaller, crowded households typically show higher TB transmission rates. In our context, it emphasizes the need to assess other environmental factors, such as ventilation, in addition to household size.

The OR for inadequate ventilation was 2.14, indicating that individuals living in poorly ventilated homes were more than twice as likely to contract TB compared to those in well-ventilated environments. This result underscores the importance of airflow in preventing the airborne transmission of *Mycobacterium tuberculosis*, particularly in enclosed, poorly ventilated spaces where bacteria can remain suspended in the air for longer periods. This finding aligns with previous research by Escombe (Escombe et al., 2007), which

demonstrated that improved ventilation significantly reduces TB transmission risk in both residential and healthcare settings.

Similarly, the OR of 1.89 for poor nutritional status suggests that individuals with inadequate nutrition were nearly twice as likely to develop TB compared to those with normal nutritional status. This reflects the well-established link between malnutrition and immune suppression, where deficiencies in essential nutrients weaken the body's ability to mount an effective immune response to TB infection. This finding is consistent with Bhargava (Bhargava et al., 2013), who emphasize the role of nutritional support in improving TB outcomes. A further investigation demonstrated a strong correlation between the kind of floor and the occurrence of tuberculosis, with a p-value of 0.001 (25). Tuberculosis (TB) was more prevalent in households that had mud (or other naturally occurring material) floors, walls, and roofs. These factors contributed to 0.42%, 0.41%, and 0.51% of all TB cases, respectively (Singh et al., 2018).

While the findings of our study align with much of the existing literature on tuberculosis (TB) risk factors, such as the influence of household density and ventilation on TB transmission, there are notable divergences from other investigations. One of the primary areas of divergence is the study's focus on the small island communities of South Sulawesi, which differ significantly from the urban and peri-urban environments that have been the focus of many TB studies.

Previous investigations, such as those by (Lee et al., 2022) and (Nagaraja et al., 2018a), have emphasized the role of household overcrowding and poor ventilation in urban settings, where high population density contributes to the rapid spread of TB. However, our study diverges in that less-crowded household conditions in the small island communities did not demonstrate the same degree of influence on TB transmission as seen in urban environments. In these island settings, the more isolated nature of households may reduce direct contact with TB-infected individuals, thereby minimizing the impact of crowding as a key risk factor. This contrasts with studies in urban areas where overcrowding is consistently identified as a significant driver of TB transmission.

Additionally, our findings highlight the disproportionate prevalence of TB among the productive-age group (15-49 years) in this specific geographic context, whereas many previous studies have reported a higher incidence among older populations. For instance, Cheng (Cheng et al., 2020a) identified the elderly as a particularly vulnerable group due to age-related declines in immunity, while our study observed the majority of TB cases in younger, productive-age individuals. This divergence may be explained by the unique socioeconomic conditions on the islands, where productive-age individuals are more likely to engage in high-exposure activities, such as fishing, farming, and trading, which require frequent travel to more densely populated areas or interaction with visitors from mainland regions.

Lastly, our study emphasizes the impact of environmental factors, such as humidity, in a tropical island climate, which may differ from findings in temperate climates studied in other research. Tanjung (Tanjung et al., 2021) also identified a similar role for humidity in the spread of TB in tropical regions, which contrasts with investigations conducted in drier or colder climates, where ventilation and household crowding are more prominent factors.

In summary, our study diverges from other TB investigations primarily due to the geographic, demographic, and environmental contexts specific to small island communities. These differences highlight the need for localized TB interventions that consider the unique social and environmental conditions present in non-urban settings.

Conclusion

Based on the present study results, the authors summarized the following conclusions: The temperature was not shown to be a risk factor for PTB. While temperature was not found significant in this study, it remains an important factor to explore, as climatic conditions may influence PTB transmission dynamics. Productive age, normal nutritional status, and mild smoking status were protective factors of PTB. Productive age and normal nutritional status likely confer protection against PTB due to stronger immune systems and reduced exposure to risk factors compared to other age groups. The risk factors for pulmonary tuberculosis incidence were male gender, low knowledge, inadequate occupant density, inadequate ventilation area, inadequate humidity, and inadequate type of floor. Key environmental risk factors identified include occupant density, ventilation, humidity, and type of flooring. Temperature may influence the survival and transmission of *Mycobacterium tuberculosis*, making it an important variable for future studies with larger sample sizes.

Declaration Conflicting Interest

The authors do not have any conflicts of interest to disclose.

Funding

This study was funded by a research grant from the Universitas Islam Negeri Alauddin Makassar, Politeknik Kesehatan Mamuju, and Universitas Muslim Indonesia.

Acknowledgment

The authors of this study would like to express their sincere gratitude to all the volunteers.

Author Contribution

All writers contributed valuable comments and played a crucial role in shaping the study, analysis, and article. **AS** played a role in formulating the proposed notion, advancing the theory, and carrying out the computations. **AS** assumed the role of primary author in composing the manuscript. **SB** had a crucial role in designing and implementing the research, analyzing the results, and producing the manuscript. **BTH** made significant contributions to the interpretation of the results. **R** formulated the theoretical framework and conducted the statistical analysis. **ZM** and **MI** made significant contributions to the final version of the manuscript.

Author Biography

Andi Susilawaty is a senior researcher and lecturer at Magister of Public Health, Postgraduate Program, Universitas Islam Negeri Alauddin Makassar, South Sulawesi, Indonesia. Her research interest is an environmental health determinant in small islands, environmental quality analysis, and health ecology and involved with various research projects in the field of another environmental health area. She can be contacted at email: andisusilawaty@uin-alauddin.ac.id

Syahrul Basri is a lecturer at the Department of Public Health, Medical and Health Science Faculty, Universitas Islam Negeri Alauddin Makassar, South Sulawesi, Indonesia. His research interest is environmental toxicology and other environmental health area. He can be contacted at email: syahrul.basri@uin-alauddin.ac.id

Bs. Titi Haerana is a lecturer at the Department of Public Health, Medical and Health Science Faculty, Universitas Islam Negeri Alauddin Makassar, South Sulawesi, Indonesia. Research interest is environmental toxicology and other environmental health area. She can be contacted at email: bs.titihaerana@uin-alauddin.ac.id

Ruslan La Ane is an academic staff at the Department of Environmental Health, School of Public Health, Hasanuddin University. He graduated from the Master's Program at the School of Public Health, Griffith University, Brisbane–Australia. His research expertise and interest are Environmental Health and Vector-Borne Diseases, particularly in modeling and bioinformatics approaches. He can be contacted at email: ruslan.ane@unhas.ac.id

Zrimurti Mappau is a lecturer at the Department of Sanitation, Politeknik Kesehatan Mamuju, Kementerian Kesehatan, Indonesia. Research interest is environmental health issues specifically on sanitation. She can be contacted at email: zmappau@gmail.com.

Muhammad Ikhtiar is an Assistant Professor and a lecturer at the Public Health Department, Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia. Involved with various research projects in the field of environmental health. He can be contacted at email: muhammad.ikhtiar@umi.ac.id

References

- Adane, A., Damena, M., Weldegebreal, F., & Mohammed, H. (2020). Prevalence and Associated Factors of Tuberculosis among Adult Household Contacts of Smear Positive Pulmonary Tuberculosis Patients Treated in Public Health Facilities of Haramaya District, Oromia Region, Eastern Ethiopia. *Tuberculosis Research and Treatment*, 2020, 1–7. from <https://doi.org/10.1155/2020/6738532>
- Amelia, S., Rahayu, R., & Wijayanti, Y. (n.d.). Analysis of Physical Environmental Factors Affecting the Incidence of Pulmonary Tuberculosis in Magelang City. In *Public Health Perspectives Journal* (Vol. 6, Issue 1). from <http://journal.unnes.ac.id/sju/index.php/phpj>
- Angelo, A. T., Geltore, T. E., & Asega, T. (2020). Knowledge, attitude, and practices towards tuberculosis among clients visiting Tepi General Hospital outpatient departments, 2019. *Infection and Drug Resistance*, 13, 4559–4568. from <https://doi.org/10.2147/IDR.S287288>
- Arman Bausat. (2021). *South Sulawesi Health Profile 2021*.
- Bashorun, A. O., Linda, C., Omoleke, S., Kendall, L., Donkor, S. D., Kinteh, M. A., Danso, B., Leigh, L., Kandeh, S., D'Alessandro, U., & Adetifa, I. M. O. (2020). Knowledge, attitude, and practice towards tuberculosis in Gambia: a nation-wide cross-sectional survey. *BMC Public Health*, 20(1). from <https://doi.org/10.1186/s12889-020-09685-3>
- Bhargava, A., Chatterjee, M., Jain, Y., Chatterjee, B., Kataria, A., Bhargava, M., Kataria, R., D'Souza, R., Jain, R., Benedetti, A., Pai, M., & Menzies, D. (2013). Nutritional Status of Adult Patients with Pulmonary Tuberculosis in Rural Central India and Its Association with Mortality. *PLoS ONE*, 8(10). from <https://doi.org/10.1371/journal.pone.0077979>
- Byng-Maddick, R., & Noursadeghi, M. (2016). Does tuberculosis threaten our aging populations? In *BMC Infectious Diseases* (Vol. 16, Issue 1). BioMed Central Ltd. <https://doi.org/10.1186/s12879-016-1451-0>

- Cheng, J., Sun, Y. N., Zhang, C. Y., Yu, Y. L., Tang, L. H., Peng, H., Peng, Y., Yao, Y. X., Hou, S. Y., Li, J. W., Zhao, J. M., Xia, L., Xu, L., Xia, Y. Y., Zhao, F., Wang, L. X., & Zhang, H. (2020a). Incidence and risk factors of tuberculosis among the elderly population in China: A prospective cohort study. *Infectious Diseases of Poverty*, 9(1). from <https://doi.org/10.1186/s40249-019-0614-9>
- Cheng, J., Sun, Y. N., Zhang, C. Y., Yu, Y. L., Tang, L. H., Peng, H., Peng, Y., Yao, Y. X., Hou, S. Y., Li, J. W., Zhao, J. M., Xia, L., Xu, L., Xia, Y. Y., Zhao, F., Wang, L. X., & Zhang, H. (2020b). Incidence and risk factors of tuberculosis among the elderly population in China: A prospective cohort study. *Infectious Diseases of Poverty*, 9(1). from <https://doi.org/10.1186/s40249-019-0614-9>
- Dodd, P. J., Looker, C., Plumb, I. D., Bond, V., Schaap, A., Shanaube, K., Muyoyeta, M., Vynnycky, E., Godfrey-Faussett, P., Corbett, E. L., Beyers, N., Ayles, H., & White, R. G. (2016). Age- and Sex-Specific Social Contact Patterns and Incidence of Mycobacterium tuberculosis Infection. *American Journal of Epidemiology*, 183(2), 156–166. from <https://doi.org/10.1093/aje/kwv160>
- Douglas G. Altman. (2024). *STROBE Statement-Checklist of Items that Should be Included in Reports of Case-Control Studies*. from <https://www.strobe-statement.org/checklists/>
- Escombe, A. R., Oeser, C. C., Gilman, R. H., Navincopa, M., Ticona, E., Pan, W., Martínez, C., Chacaltana, J., Rodríguez, R., Moore, D. A. J., Friedland, J. S., & Evans, C. A. (2007). Natural ventilation for the prevention of airborne contagion. *PLoS Medicine*, 4(2), 0309–0317. from <https://doi.org/10.1371/journal.pmed.0040068>
- Horton, K. C., MacPherson, P., Houben, R. M. G. J., White, R. G., & Corbett, E. L. (2016). Sex Differences in Tuberculosis Burden and Notifications in Low- and Middle-Income Countries: A Systematic Review and Meta-analysis. In *PLoS Medicine* (Vol. 13, Issue 9). Public Library of Science. from <https://doi.org/10.1371/journal.pmed.1002119>
- Hussien, B., Hussien, M. M., Seid, A., & Hussien, A. (2019). Nutritional deficiency and associated factors among new pulmonary tuberculosis patients of Bale Zone Hospitals, southeast Ethiopia. *BMC Research Notes*, 12(1). from <https://doi.org/10.1186/s13104-019-4786-y>
- ICH. (2016). *International Council for Harmonization of Technical Requirement for Pharmaceutical for Human Use (ICH) ICH Harmonized Guideline Integrated Addendum*. from https://database.ich.org/sites/default/files/E6_R2_Addendum.pdf
- KEPKN. (2021). *Pedoman dan Standar Etik Penelitian & Pengembangan Kesehatan Nasional*. Lembaga Penerbitan dan Pengembangan Kesehatan. <https://keppkn.kemkes.go.id/index.php/page/post/pedoman-dan-standar-etik>
- Laghari, M., Sulaiman, S. A. S., Khan, A. H., Talpur, B. A., Bhatti, Z., & Memon, N. (2019). Contact screening and risk factors for TB among the household contacts of children with active TB: A way to find source case and new TB cases. *BMC Public Health*, 19(1). from <https://doi.org/10.1186/s12889-019-7597-0>
- Lee, J. Y., Kwon, N., Goo, G. Yeon, & Cho, S. il. (2022). Inadequate housing and pulmonary tuberculosis: a systematic review. *BMC Public Health*, 22(1). from <https://doi.org/10.1186/s12889-022-12879-6>
- Moule, M. G., & Cirillo, J. D. (2020). Mycobacterium tuberculosis Dissemination Plays a Critical Role in Pathogenesis. *Frontiers in Cellular and Infection Microbiology*, 10. from <https://doi.org/10.3389/fcimb.2020.00065>
- Nagaraja, S. B., Satyanarayana, S., & Bansal, A. K. (2018a). Can ventilation oust tuberculosis bacilli? Dare to plug the unpluggable. *Public Health Action*, 8(1), 28–28. from <https://doi.org/10.5588/pha.17.0115>
- Nagaraja, S. B., Satyanarayana, S., & Bansal, A. K. (2018b). Can ventilation oust tuberculosis bacilli? Dare to plug the unpluggable. *Public Health Action*, 8(1), 28–28. from <https://doi.org/10.5588/pha.17.0115>
- Ncayiyana, J. R., Bassett, J., West, N., Westreich, D., Musenge, E., Emch, M., Pettifor, A., Hanrahan, C. F., Schwartz, S. R., Sanne, I., & van Rie, A. (2016). Prevalence of latent tuberculosis infection and predictive factors in an urban informal settlement in Johannesburg, South Africa: A cross-sectional study. *BMC Infectious Diseases*, 16(1). from <https://doi.org/10.1186/s12879-016-1989-x>
- Shimeles, E., Enquselassie, F., Aseffa, A., Tilahun, M., Mekonen, A., Wondimagegn, G., & Hailu, T. (2019). Risk factors for tuberculosis: A case-control study in Addis Ababa, Ethiopia. *PLoS ONE*, 14(4). from <https://doi.org/10.1371/journal.pone.0214235>
- Singh, S. K., Kashyap, G. C., & Puri, P. (2018). The potential effect of household environment on the prevalence of tuberculosis in India: Evidence from the recent round of a cross-sectional survey. *BMC Pulmonary Medicine*, 18(1). from <https://doi.org/10.1186/s12890-018-0627-3>
- Stuckler, D., Basu, S., McKee, M., & Lurie, M. (2011). Mining and Risk of Tuberculosis in Sub-Saharan Africa. *American Journal of Public Health*. from <https://doi.org/10.2105/AJPH.2009>
- Suntiono, & et all. (2023). *Sulawesi Selatan in Figures 2023*. from <https://sulsel.bps.go.id>
- Tanjung, R., Mahyuni, E. L., Tanjung, N., Simarmata, O. S., Sinaga, J., & Nolia, H. R. (2021). The spatial distribution of pulmonary tuberculosis in Kabanjahe District, Karo Regency, Indonesia. *Open Access Macedonian Journal of Medical Sciences*, 9, 817–822. from <https://doi.org/10.3889/oamjms.2021.6808>
- Tereza Kasaeva. (2022). *Global Tuberculosis Report 2022*. <https://iris.who.int/bitstream/handle/10665/363752/9789240061729-eng.pdf>
- Tereza Kasaeva. (2023). *Global Tuberculosis Report 2023*. World Health Organization. from <https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2023>
- Tiomaida Seviana, & et all. (2023). *Indonesia Health Profile 2023*. from <https://kemkes.go.id/id/profil-kesehatan-indonesia-2023>

WMA General Assembly. (2024). *WMA Declaration of Helsinki-Ethical Principle for Medical Research Involving Human Participants General Principles*. <https://www.wma.net/policies-post/wma-declaration-of-helsinki/>

Cite this article as: Susilawaty, A., Syahrul Basri, S., Haerana, Bs. T., Ane, R.L., Mappau, Z., Ikhtia, M. (2024). The Risk Factors Contributing to The Occurrence of Pulmonary Tuberculosis in The Small Island of South Sulawesi, Indonesia, *Public Health of Indonesia*, 10(4), 104-116. <https://doi.org/10.36685/phi.v10i4.876>